

Complexity Theory

Georg Moser

Institute of Computer Science @ UIBK

Summer 2008

- Summary of Last Lecture: Separation Results
- Exercises
- The Immerman-Szelepcsényi Theorem

Separation Results

Theorem

Let $S(n)$ be space-constructible. Then there exists a set in $\text{DSPACE}(S(n))$ that is **not** in $\text{DSPACE}(S'(n))$ for any $S'(n) = o(S(n))$.

Lemma

$$\text{NSPACE}(n^3) \not\subseteq \text{NSPACE}(n^4)$$

A Space-Constructible Function $S(n) \leq O(\log \log n)$

Definition

- 1 let $\ell(n)$ be the least positive number not dividing n
- 2 note that $\ell(n)$ is a prime power
- 3 $\ell(n) = O(\log n)$, because for every ℓ :

$$\prod_{p \leq \ell, p \text{ prime}} p \geq 2^{\Omega(\ell)}$$

Lemma

\exists an unbounded space-constructible function $S(n)$ that is $O(\log \log n)$, namely the space needed to accept $A = \{a^n \mid \ell(n) \text{ is prime}\}$.

Lemma

The function $\lceil \log \log n \rceil$ is **not** space constructible.

Homework

- 1 Miscellaneous Exercises 61 in Kozen, Automata and Computability
- 2 Homework 1.2
- 3 Homework 1.3
- 4 Homework 2.1
- 5 Homework 2.2
- 6 Homework 2.3

Proof (for ①)

- 1 let M be nondeterministic $S(n)$ -space bounded TM; we wish to build a $S(n)$ -space bounded TM N accepting the complement of $L(M)$
- 2 suppose encoding of the configurations of M over Δ $|\Delta| = d$
- 3 configurations on inputs of length n become strings in $\Delta^{S(n)}$
- 4 assume there is a unique accept configuration $\text{accept} \in \Delta^{S(n)}$
- 5 let $\text{start} \in \Delta^{S(n)}$ denote the start configuration on input x $|x| = n$
- 6 if x is accepted, the computation has length $\leq d^{S(n)}$
- 7 define $A_m = \{\alpha \in \Delta^{S(n)} \mid \text{start} \xrightarrow{\leq m} \alpha\}$
- 8 thus $A_0 = \{\text{start}\}$ and M accepts x iff $\text{accept} \in A_{d^{S(n)}}$

Theorem

Immerman-Szelepcsényi Theorem

For $S(n) \geq \log n$, $\text{NSPACE}(S(n)) = \text{co-NSPACE}(S(n))$.

Proof Idea

- 1 assume we have finite set A and a nondeterministic test for membership
- 2 assume we know the size of A
- 3 we can test for non-membership as follows:
 - given y
 - let $n := |A|$
 - guess n elements x , make sure $x \in A$ and $x \neq y$
 - hence $y \notin A$

Two cases:

- ① Assume $S(n)$ is space-constructible
- ② Assume $S(n) \geq \log n$

Construction of N

- lay off $S(n)$ space on the worktape $S(n)$ is space-constructible
- compute sizes $|A_0|, |A_1|, \dots$ inductively
- first $|A_0| = 1$
- suppose $|A_m|$ is computed and written on the worktape:
 - 1 write down $\beta \in \Delta^{S(n)}$
 - 2 determine whether $\beta \in A_{m+1}$
 - 3 increment counter
- suppose $|A_{d^{S(n)}}|$ is known
- guess $|A_{d^{S(n)}}|$ α 's to verify that $\text{start} \xrightarrow{\leq d^{S(n)}} \alpha$ holds
- test $\alpha \neq \text{accept}$
- if successful: $\text{accept} \notin A_{d^{S(n)}}$ and N can accept

the above steps can be implemented such that N uses at most $S(n)$ space

Determine $\beta \in A_{m+1}$

- guess $|A_m|$ elements α of A_m in lexicographic order
- verify for each $\alpha \in A_m$ by guessing a path to test $\text{start} \xrightarrow{\leq m} \alpha$
- test $\alpha \xrightarrow{\leq 1} \beta$
- if $\exists \alpha$ with $\alpha \xrightarrow{=1} \beta$: $\beta \in A_{m+1}$
- if $\forall \alpha$ with $\alpha \not\xrightarrow{=1} \beta$: $\beta \notin A_{m+1}$

In sum: the machine N accepts complement of $L(M)$ is space $S(n)$

Proof (for ②)

To remove the space-constructible assumption for $S(n)$ do the construction on the fly, increasing the available space if necessary

