

- [2] 1 Find two different minimal unsatisfiable cores and the MUC of the following formula:

$$(x \vee y \vee z) \wedge \neg x \wedge (x \vee y) \wedge (x \vee \neg y \vee z) \wedge (x \vee z) \wedge (\neg x \vee \neg y \vee z) \wedge (y \vee \neg z) \wedge \neg z$$


- [3] 2 Use the `minUnsatCore` algorithm to find a minimal unsatisfiable core of the following formula:

$$(x \vee y \vee z) \wedge \neg x \wedge (x \vee y) \wedge (x \vee \neg y) \wedge \neg z$$

- 3 The problem of k -vertex coloring for a given undirected graph assumes that there are k colors given and asks to find a color for every node such that no adjacent nodes have the same color.

More precisely, given a graph $G = (V, E)$ a k -coloring for G exists if there is a function $c : V \rightarrow \{1, \dots, k\}$ such that for all $(u, v) \in E$ we have $c(u) \neq c(v)$. The smallest k which admits a k -coloring of a graph G is called the *chromatic number* of G .

One application is coloring maps. For example, the following graphs are a valid (a) and an invalid (b) 4-coloring of the state map of Austria:



Being in NP, k -vertex coloring can be reduced to SAT, e.g. by using propositional variables n_1, \dots, n_k such that n_i becomes true if and only if node n has color i .

- [3] (a) Use a SAT encoding to determine whether there is a 2-coloring of the state map of Austria. If there is no such coloring, get an unsatisfiable core to determine a minimal subgraph for which no 2-coloring exists.
- [2] (b) Use a SAT encoding to find a 3-coloring of the state map of Austria.
- [3] \star (c) Write a function which takes an undirected graph G and a number k and uses a SAT encoding to determine whether a k -coloring of G exists. (The graph can e.g. be given as an adjacency list, i.e., a list of edges.)