

Computational Logic

Vincent van Oostrom
Course/slides by Aart Middeldorp

Department of Computer Science
University of Innsbruck

SS 2020



Overviews

Overview of previous lecture

The previous lecture was concerned with two things, namely (meta-theoretic) consequences of Model Existence and the presentation of Hilbert systems.

- The rough idea of Model Existence is based on maximally deriving consequences of a given set of formulas, without arriving at a contradiction. Maximality is captured by the notion of a **Hintikka** set, which one can think of as expressing that the set is **closed** under taking consequences, i.e. closed under the tableau expansion rules, and is **consistent** in that it does not contain a formula and its negation, no contradiction. Hintikka's lemma expresses that such sets are **satisfiable** (think of satisfying a branch of a tableau where we have exhausted all possible consequences without it being closed), and Model Existence that every set in a **Propositional Consistency Property**, a collection of sets of formulas having certain closure properties, is satisfiable **because** it can be extended to a Hintikka set (in the collection). From this **completeness** of **tableaux** follows, since if never a closed tableau is obtained (from the **negation** of a formula), a Propositional Consistency Property is obtained, so the negated formula is satisfiable, hence the original formula a **tautology**.

Outline

- Overviews
- First-Order Logic
- Herbrand Models
- Uniform Notation
- Hintikka's Lemma
- Exercises
- Further Reading

Overviews

Overview of previous lecture (ctd)

- Compactness will be commented on below, for first-order logic.
- Interpolation is a meta-theoretic result having uses in model checking and database theory. One can think of it as expressing a restriction on the **converse** of transitivity: Whereas, transitivity expresses that given X implies Y and Y implies Z , it follows that X implies Z , Craig's interpolation theorem expresses that given X implies Z , one can find an **interpolant** Y such that X implies Y and Y implies Z , and that for finding Y we may restrict ourselves to formulas speaking about the variables X and Z ; i.e. an interpolant/way-point Y can always be found using only variables that are in the common language of X and Z . Versions of interpolation hold e.g. for first-order logic (we will see) and equational logic. There are also logics for which interpolation does not hold, e.g. rewriting logic.

Overview of previous lecture (ctd)

- There is usually an exchange in inference systems, between having many axioms and few inference rules, and having few axioms and many inference rules. Hilbert systems are the extreme case of the former, just the inference rule of **modus ponens**, whereas natural deduction is an instance of the latter, having introduction and elimination rules for each connective. The **deduction theorem** connects both by showing that Y can be inferred from X iff there is a Hilbert proof of $X \supset Y$. The proof is constructive, so that to prove the latter, one can first prove the former, and then transform that into a Hilbert proof using the deduction theorem. The Hilbert system presented has only **two** axioms, but was shown complete using Model Existence, showing the versatility of the latter result.

Overview of this lecture

This lecture is concerned with generalising our set-up for to first-order logic, the default logic in automated reasoning, by allowing to express properties of and relations between individuals (predicates instead of just properties), and all (\forall) or some (\exists) of these. The items below, on this page, should be known.

- **syntax**: **terms** to represent individuals (e.g. natural numbers) and operations on them (e.g. addition), **predicates** to describe properties of individuals (unary predicates) and relations between them (binary, ternary, ... predicates), and **quantifiers** to express properties holding for some/all individuals.
- **semantics**: meaning of terms is given by means of a **domain** from which individuals are taken and over which quantifiers range; operations are **interpreted** as functions on the domain, and predicates as relation on it. The meaning of a concrete formula depends on an **assignment** giving meaning to the variables in a term, as elements of the domain. This dependence is there to allow interpreting formulas having quantifiers by **recursion** on the formula. E.g. $(\forall x)\Phi$ is true if **for all** assignments to x , the **subformula** Φ is true. The semantics is then used to define the generalised notions of **validity** and **satisfiability**.

Overview of this lecture (ctd)

- Herbrand models are models where terms are interpreted as **themselves**. That is, Herbrand models are a kind of **syntactic** models; instead of taking as domain say the natural numbers, or people or ... we take the terms themselves as individuals. For the Herbrand model, assignments **are** substitutions for the variables. Herbrand models are counterintuitive (aren't interpretations meant to give **meaning/semantics** to the **language/syntax**? how can syntax fulfill the role of semantics?) but (cf. **free groups**) they work: Herbrand models are typically used for meta-theoretic results connecting semantics to syntax, with the reasoning going roughly as follows: if a formula is valid, then it is true in all models, so in particular in the Herbrand model; thus (Herbrand's theorem), if the formula is existential it suffices to find appropriate **terms** for the existentially quantified variables. This enables automation. Proof search by enumerating terms is the basis for Prolog (how would one check all interpretations in all domains?).
- Uniform notation is generalised by noting that \forall is a generalised conjunction and \exists a generalised disjunction. For example, one can think of $(\exists n)n \geq 5$ in the natural numbers as $(0 \geq 5) \vee (1 \geq 5) \vee (2 \geq 5) \dots$

Overview of this lecture (ctd)

- The meta-theoretic results follow by generalising the propositional case, or rather the converse: Hintikka and Model Existence were set-up for the propositional case such that they would allow easy generalisation to the first-order case. In the propositional case one could often do with **finite** sets of propositions (e.g. for showing completeness results), but that would not do for the first-order case. An intuition for this insufficiency is provided by the above correspondence between quantified formulas and **infinite** con/disjunctions.
- **Compactness** is a meta-theoretic result in that it can be used to show the **limitations** of first-order logic. In particular, it implies that there is no first-order formula that can express that the domain is **finite**. From the **Löwenheim–Skolem** theorem follows a similar limitation of first-order logic: whatever **first-order** axiomatisation one gives of the real numbers, there will always be a **countable** model. This means that first-order formulas cannot capture the uncountability of the real numbers.

Part I: Propositional Logic

compactness, completeness, Hilbert systems, Hintikka's lemma, interpolation, logical consequence, model existence theorem, propositional semantic tableaux, soundness

Part II: First-Order Logic

compactness, completeness, Craig's interpolation theorem, cut elimination, first-order semantic tableaux, Herbrand models, Herbrand's theorem, Hilbert systems, Hintikka's lemma, Löwenheim-Skolem, logical consequence, model existence theorem, prenex form, skolemization, soundness

Part III: Limitations and Extensions of First-Order Logic

Curry-Howard isomorphism, intuitionistic logic, Kripke models, second-order logic, simply-typed λ -calculus

Outline

- Overviews
- First-Order Logic
 - Syntax
 - Substitutions
 - Semantics
- Herbrand Models
- Uniform Notation
- Hintikka's Lemma
- Exercises
- Further Reading

First-Order Languages – Common Items

- **propositional connectives**: primary connectives are basic, secondary connectives are defined, propositional constants \top and \perp
- **quantifiers**: \forall and \exists
- **variables**: v_1, v_2, \dots

Definition

first-order language is determined by specifying

- 1 countable set **R** of **relation** or **predicate symbols**, each of which has positive integer associated with it
- 2 countable set **F** of **function symbols**, each of which has positive integer associated with it
- 3 countable set **C** of **constant symbols**

notation: $L(\mathbf{R}, \mathbf{F}, \mathbf{C})$ (or simply L)

Definition

family of **terms** of $L(\mathbf{R}, \mathbf{F}, \mathbf{C})$ is smallest set such that

- 1 any variable is term of $L(\mathbf{R}, \mathbf{F}, \mathbf{C})$
- 2 any constant symbol (member of **C**) is term of $L(\mathbf{R}, \mathbf{F}, \mathbf{C})$
- 3 if f is n -place function symbol (member of **F**) and t_1, \dots, t_n are terms of $L(\mathbf{R}, \mathbf{F}, \mathbf{C})$ then $f(t_1, \dots, t_n)$ is term of $L(\mathbf{R}, \mathbf{F}, \mathbf{C})$

term is **closed** if it contains no variables

Example

one-place function symbol f , two-place function symbol g , constants a and b , variables x and y

terms: $f(g(a, x))$ $g(f(x), g(x, y))$ $g(a, g(a, g(a, b)))$

Definition

atomic formula of $L(\mathbf{R}, \mathbf{F}, \mathbf{C})$ is any string of form $R(t_1, \dots, t_n)$ where R is n -place relation symbol (member of \mathbf{R}) and t_1, \dots, t_n are terms of $L(\mathbf{R}, \mathbf{F}, \mathbf{C})$
also \top and \perp are atomic formulas of $L(\mathbf{R}, \mathbf{F}, \mathbf{C})$

Definition

family of **formulas** of $L(\mathbf{R}, \mathbf{F}, \mathbf{C})$ is smallest set such that

- 1 any atomic formula of $L(\mathbf{R}, \mathbf{F}, \mathbf{C})$ is formula of $L(\mathbf{R}, \mathbf{F}, \mathbf{C})$
- 2 if A is formula of $L(\mathbf{R}, \mathbf{F}, \mathbf{C})$ so is $\neg A$
- 3 if A and B are formulas of $L(\mathbf{R}, \mathbf{F}, \mathbf{C})$ and \circ is binary connective then $A \circ B$ is formula of $L(\mathbf{R}, \mathbf{F}, \mathbf{C})$
- 4 if A is formula of $L(\mathbf{R}, \mathbf{F}, \mathbf{C})$ and x is variable then $(\forall x)A$ and $(\exists x)A$ are formulas of $L(\mathbf{R}, \mathbf{F}, \mathbf{C})$

Example

two-place relation symbol R , two-place function symbol f

formulas: $(\forall x)(\forall y)(R(x, y) \supset (\exists z)(R(x, z) \wedge R(z, y)))$ $(\forall x)(\exists y)R(f(x, y), z)$

Definition

free-variable occurrences in formula are defined as follows:

- 1 free-variable occurrences in atomic formula are all variable occurrences in that formula
- 2 free-variable occurrences in $\neg A$ are free-variable occurrences in A
- 3 free-variable occurrences in $(A \circ B)$ are free-variable occurrences in A together with free-variable occurrences in B
- 4 free-variable occurrences in $(\forall x)A$ and $(\exists x)A$ are free-variable occurrences in A , except for occurrences of x

variable occurrence is **bound** if it is not free

Example

$(\forall x)[(\exists y)R(f(x, y), c) \supset (\exists z)S(y, z)]$ free-variable occurrences

$(\forall x)[(\exists y)R(f(x, y), c) \supset (\exists z)S(y, z)]$ bound-variable occurrences

Definition

sentence (or **closed formula**) of $L(\mathbf{R}, \mathbf{F}, \mathbf{C})$ is formula of $L(\mathbf{R}, \mathbf{F}, \mathbf{C})$ with no free-variable occurrences

Outline

- Overviews
- First-Order Logic
 - Syntax
 - Substitutions
 - Semantics
- Herbrand Models
- Uniform Notation
- Hintikka's Lemma
- Exercises
- Further Reading

Definition

substitution is mapping $\sigma: \mathbf{V} \rightarrow \mathbf{T}$ from set of variables \mathbf{V} to set of terms \mathbf{T}

Definition

given substitution σ

- 1 $c\sigma = c$ for constant symbol c
- 2 $f(t_1, \dots, t_n)\sigma = f(t_1\sigma, \dots, t_n\sigma)$ for n -place function symbol f

Example

$x\sigma = f(x, y)$, $y\sigma = h(a)$, $z\sigma = g(c, h(x))$
 $j(k(x), y)\sigma = j(k(f(x, y)), h(a))$

Definition

composition of substitutions σ and τ is substitution $\sigma\tau$ such that $x(\sigma\tau) = (x\sigma)\tau$ for each variable x

Lemma

$t(\sigma\tau) = (t\sigma)\tau$ for every term t

Lemma

composition of substitutions is **associative**

Definition

- **support** of substitution σ is set of variables x for which $x\sigma \neq x$
- substitution has **finite support** if support set is finite

Lemma

composition of substitutions having finite support is substitution having finite support

Notation

$\{x_1/t_1, \dots, x_n/t_n\}$ for substitution σ having finite support $\{x_1, \dots, x_n\}$ and $x_i\sigma = t_i$ for $1 \leq i \leq n$

Definition

given substitution σ and variable x , substitution σ_x is defined as follows:

$y\sigma_x = x$ if $y = x$ and $y\sigma_x = y\sigma$ if $y \neq x$

Definition

- 1 $[A(t_1, \dots, t_n)]\sigma = A(t_1\sigma, \dots, t_n\sigma)$, $\top\sigma = \top$, $\perp\sigma = \perp$
- 2 $[\neg X]\sigma = \neg[X\sigma]$
- 3 $(X \circ Y)\sigma = (X\sigma \circ Y\sigma)$
- 4 $[(\forall x)\Phi]\sigma = (\forall x)[\Phi\sigma_x]$
- 5 $[(\exists x)\Phi]\sigma = (\exists x)[\Phi\sigma_x]$

Example

if $\sigma = \{x/a, y/b\}$ then

$$\begin{aligned} [(\forall x)R(x, y) \supset (\exists y)R(x, y)]\sigma &= [(\forall x)R(x, y)]\sigma \supset [(\exists y)R(x, y)]\sigma \\ &= (\forall x)[R(x, y)]\sigma_x \supset (\exists y)[R(x, y)]\sigma_y \\ &= (\forall x)R(x, b) \supset (\exists y)R(a, y) \end{aligned}$$

Remark

$(\Phi\sigma)\tau = \Phi(\sigma\tau)$ need not hold

Example

- $\Phi = (\forall y)R(x, y)$, $\sigma = \{x/y\}$, $\tau = \{y/c\}$
- $(\Phi\sigma)\tau = (\forall y)R(y, y)\tau = (\forall y)R(y, y)$
 - $\sigma\tau = \{x/c, y/c\}$
 - $\Phi(\sigma\tau) = (\forall y)R(c, y)$

Definition

substitution σ being **free for formula** is defined as follows:

- 1 if A is atomic then σ free for A
- 2 σ is free for $\neg X$ if σ is free for X
- 3 σ is free for $(X \circ Y)$ if σ is free for X and σ is free for Y
- 4 σ is free for $(\forall x)\Phi$ and $(\exists x)\Phi$ provided
 - σ_x is free for Φ
 - if y is free variable of Φ other than x , $y\sigma$ does not contain x

Theorem

if substitution σ is free for formula X and substitution τ is free for $X\sigma$ then
 $(X\sigma)\tau = X(\sigma\tau)$

Proof

structural induction on X

- atomic case is obvious
- $X = \neg Y$ σ is free for Y $X\sigma = \neg(Y\sigma)$
 τ is free for $Y\sigma$
 $(Y\sigma)\tau = Y(\sigma\tau)$ follows from induction hypothesis
 $(X\sigma)\tau = [\neg(Y\sigma)]\tau = \neg((Y\sigma)\tau) = \neg(Y(\sigma\tau)) = (\neg Y)(\sigma\tau) = X(\sigma\tau)$
- $X = (Y \circ Z)$ σ is free for Y and Z $X\sigma = (Y\sigma \circ Z\sigma)$
 τ is free for $Y\sigma$ and $Z\sigma$
 $(Y\sigma)\tau = Y(\sigma\tau)$ and $(Z\sigma)\tau = Z(\sigma\tau)$ follow from induction hypothesis
 $(X\sigma)\tau = ((Y\sigma)\tau \circ (Z\sigma)\tau) = (Y(\sigma\tau) \circ Z(\sigma\tau)) = (Y \circ Z)(\sigma\tau) = X(\sigma\tau)$

Proof (cont'd)

structural induction on X

- $X = (\forall x)\Phi$ σ_x is free for Φ τ is free for $[(\forall x)\Phi]\sigma = (\forall x)[\Phi\sigma_x]$
 τ_x is free for $\Phi\sigma_x$
 $(\Phi\sigma_x)\tau_x = \Phi(\sigma_x\tau_x)$ follows from induction hypothesis
 claim: $\Phi(\sigma_x\tau_x) = \Phi(\sigma\tau)_x$
 let y be free variable of Φ
 - if $y = x$ then $y(\sigma_x\tau_x) = (y\sigma_x)\tau_x = x\tau_x = x = y(\sigma\tau)_x$
 - if $y \neq x$ then $y(\sigma_x\tau_x) = (y\sigma_x)\tau_x = (y\sigma)\tau_x = (y\sigma)\tau = y(\sigma\tau) = y(\sigma\tau)_x$
$$(X\sigma)\tau = ((\forall x)[\Phi\sigma_x])\tau = (\forall x)[(\Phi\sigma_x)\tau_x] = (\forall x)[\Phi(\sigma_x\tau_x)]$$

$$= (\forall x)[\Phi(\sigma\tau)_x] = X(\sigma\tau)$$
- $X = (\exists x)\Phi$ similar

Outline

- Overviews
- First-Order Logic
 - Syntax
 - Substitutions
 - Semantics
- Herbrand Models
- Uniform Notation
- Hintikka's Lemma
- Exercises
- Further Reading

Definition

model for first-order language $L(\mathbf{R}, \mathbf{F}, \mathbf{C})$ is pair $\mathbf{M} = \langle \mathbf{D}, \mathbf{I} \rangle$ where

- \mathbf{D} is nonempty set, called **domain** of \mathbf{M}
- \mathbf{I} is mapping, called **interpretation**, that associates
 - to every $c \in \mathbf{C}$ some member $c^{\mathbf{I}} \in \mathbf{D}$
 - to every n -place $f \in \mathbf{F}$ some n -ary function $f^{\mathbf{I}}: \mathbf{D}^n \rightarrow \mathbf{D}$
 - to every n -place $P \in \mathbf{R}$ some n -ary relation $P^{\mathbf{I}} \subseteq \mathbf{D}^n$

Definition

assignment in model $\mathbf{M} = \langle \mathbf{D}, \mathbf{I} \rangle$ is mapping \mathbf{A} from set of variables to set \mathbf{D}
image of variable v under assignment \mathbf{A} is denoted by $v^{\mathbf{A}}$

Definition

given model $\mathbf{M} = \langle \mathbf{D}, \mathbf{I} \rangle$ for language $L(\mathbf{R}, \mathbf{F}, \mathbf{C})$ and assignment \mathbf{A} in \mathbf{M} , value $t^{\mathbf{I}, \mathbf{A}}$ in \mathbf{D} is defined inductively:

- 1 $c^{\mathbf{I}, \mathbf{A}} = c^{\mathbf{I}}$ for constant symbol c
- 2 $v^{\mathbf{I}, \mathbf{A}} = v^{\mathbf{A}}$ for variable v
- 3 $[f(t_1, \dots, t_n)]^{\mathbf{I}, \mathbf{A}} = f^{\mathbf{I}}(t_1^{\mathbf{I}, \mathbf{A}}, \dots, t_n^{\mathbf{I}, \mathbf{A}})$ for n -place function symbol f

Example

L with constant 0, one-place function symbol s , two-place function symbol $+$
terms $t_1 = s(s(0) + s(x))$ and $t_2 = s(x + s(x + s(0)))$

model $\mathbf{M}_2 = \langle \mathbf{D}, \mathbf{I} \rangle$ with $\mathbf{D} = \{a, b\}^*$, $0^{\mathbf{I}} = a$, $s^{\mathbf{I}}(w) = wa$, $+^{\mathbf{I}}(v, w) = vw$

assignment \mathbf{A} with $x^{\mathbf{A}} = aba$

$t_1^{\mathbf{I}, \mathbf{A}} = aaabaaa$ and $t_2^{\mathbf{I}, \mathbf{A}} = abaabaaaa$

Definition

assignment \mathbf{B} in model \mathbf{M} is **x -variant** of assignment \mathbf{A} provided \mathbf{A} and \mathbf{B} assign same values to every variable except possibly x

Definition

given model $\mathbf{M} = \langle \mathbf{D}, \mathbf{I} \rangle$ for language $L(\mathbf{R}, \mathbf{F}, \mathbf{C})$ and assignment \mathbf{A} in \mathbf{M} , truth value $\Phi^{\mathbf{I}, \mathbf{A}}$ for formula Φ of $L(\mathbf{R}, \mathbf{F}, \mathbf{C})$ is defined inductively:

- 1 $[P(t_1, \dots, t_n)]^{\mathbf{I}, \mathbf{A}} = t \iff \langle t_1^{\mathbf{I}, \mathbf{A}}, \dots, t_n^{\mathbf{I}, \mathbf{A}} \rangle \in P^{\mathbf{I}} \quad \top^{\mathbf{I}, \mathbf{A}} = t \quad \perp^{\mathbf{I}, \mathbf{A}} = f$
- 2 $[\neg X]^{\mathbf{I}, \mathbf{A}} = \neg[X]^{\mathbf{I}, \mathbf{A}}$
- 3 $[X \circ Y]^{\mathbf{I}, \mathbf{A}} = X^{\mathbf{I}, \mathbf{A}} \circ Y^{\mathbf{I}, \mathbf{A}}$
- 4 $[(\forall x)\Phi]^{\mathbf{I}, \mathbf{A}} = t \iff \Phi^{\mathbf{I}, \mathbf{B}} = t$ for every assignment \mathbf{B} in \mathbf{M} that is x -variant of \mathbf{A}
- 5 $[(\exists x)\Phi]^{\mathbf{I}, \mathbf{A}} = t \iff \Phi^{\mathbf{I}, \mathbf{B}} = t$ for some assignment \mathbf{B} in \mathbf{M} that is x -variant of \mathbf{A}

Notation

$\Phi^{\mathbf{I}}$ instead of $\Phi^{\mathbf{I}, \mathbf{A}}$ for formulas Φ without free variables

Definitions

- formula Φ of $L(\mathbf{R}, \mathbf{F}, \mathbf{C})$ is **true in model** $\mathbf{M} = \langle \mathbf{D}, \mathbf{I} \rangle$ for $L(\mathbf{R}, \mathbf{F}, \mathbf{C})$ provided $\Phi^{\mathbf{I}, \mathbf{A}} = t$ for all assignments \mathbf{A}
- formula Φ is **valid** if Φ is true in all models for $L(\mathbf{R}, \mathbf{F}, \mathbf{C})$
- set S of formulas is **satisfiable in** $\mathbf{M} = \langle \mathbf{D}, \mathbf{I} \rangle$ provided there exists assignment \mathbf{A} (called **satisfying assignment**) such that $\Phi^{\mathbf{I}, \mathbf{A}} = t$ for all $\Phi \in S$
- set S of formulas is **satisfiable** if S is satisfiable in some model

Example

L with two-place function symbol \oplus and two-place relation symbol R

- model $\mathbf{M} = \langle \mathbf{D}, \mathbf{I} \rangle$ with $\mathbf{D} = \mathbb{N}$, $\oplus^{\mathbf{I}}(x, y) = x + y$, $R^{\mathbf{I}}$ is equality relation
formula $(\exists y)R(x, y \oplus y)$
 $(\exists y)R(x, y \oplus y)^{\mathbf{I}, \mathbf{A}}$ is true if and only if $x^{\mathbf{A}}$ is even number
- model $\mathbf{M} = \langle \mathbf{D}, \mathbf{I} \rangle$ with $\mathbf{D} = \mathbb{N}_+$, $\oplus^{\mathbf{I}}(x, y) = x + y$, $R^{\mathbf{I}}$ is greater-than relation
sentence $(\forall x)(\forall y)(\exists z)R(x \oplus y, z)$ is true in \mathbf{M}

Lemma

given closed term t , formula Φ of first-order language L , model $\mathbf{M} = \langle \mathbf{D}, \mathbf{I} \rangle$ for L
if x is variable and \mathbf{A} assignment such that $x^{\mathbf{A}} = t^{\mathbf{I}}$ then $[\Phi\{x/t\}]^{\mathbf{I}, \mathbf{B}} = \Phi^{\mathbf{I}, \mathbf{A}}$ for any x -variant \mathbf{B} of \mathbf{A}

Lemma

given model $\mathbf{M} = \langle \mathbf{D}, \mathbf{I} \rangle$ for language L , formula Φ in L , assignment \mathbf{A} in \mathbf{M} ,
substitution σ that is free for Φ

if assignment \mathbf{B} is defined by $v^{\mathbf{B}} = (v\sigma)^{\mathbf{I}, \mathbf{A}}$ for each variable v then $\Phi^{\mathbf{I}, \mathbf{B}} = (\Phi\sigma)^{\mathbf{I}, \mathbf{A}}$

Proof

structural induction on Φ

- atomic** and propositional cases are straightforward
 $t^{\mathbf{I}, \mathbf{B}} = (t\sigma)^{\mathbf{I}, \mathbf{A}}$ for all terms t is obtained by induction on t
- $\Phi = (\exists x)\varphi$ suppose $(\Phi\sigma)^{\mathbf{I}, \mathbf{A}} = [(\exists x)(\varphi\sigma_x)]^{\mathbf{I}, \mathbf{A}} = t$
 $(\varphi\sigma_x)^{\mathbf{I}, \mathbf{A}'} = t$ for some x -variant \mathbf{A}' of \mathbf{A}
define assignment \mathbf{B}' by $v^{\mathbf{B}'} = (v\sigma_x)^{\mathbf{I}, \mathbf{A}'}$ for each variable v
 σ_x is free for φ
 $\varphi^{\mathbf{I}, \mathbf{B}'} = (\varphi\sigma_x)^{\mathbf{I}, \mathbf{A}'} = t$ by induction hypothesis

Proof (cont'd)

- $\Phi = (\exists x)\varphi$ $(\Phi\sigma)^{\mathbf{I}, \mathbf{A}} = [(\exists x)(\varphi\sigma_x)]^{\mathbf{I}, \mathbf{A}} = t$ σ is free for Φ
assignment \mathbf{B}' with $v^{\mathbf{B}'} = (v\sigma_x)^{\mathbf{I}, \mathbf{A}'}$ for each variable v
 $\varphi^{\mathbf{I}, \mathbf{B}'} = (\varphi\sigma_x)^{\mathbf{I}, \mathbf{A}'} = t$
claim: \mathbf{B}' is x -variant \mathbf{B}
if $v \neq x$ then $v^{\mathbf{B}'} = (v\sigma_x)^{\mathbf{I}, \mathbf{A}'} = (v\sigma)^{\mathbf{I}, \mathbf{A}} = (v\sigma)^{\mathbf{I}, \mathbf{A}} = v^{\mathbf{B}}$
 $\Phi^{\mathbf{I}, \mathbf{B}} = [(\exists x)\varphi]^{\mathbf{I}, \mathbf{B}} = \varphi^{\mathbf{I}, \mathbf{B}'} = t$
proof of converse direction is similar
- $\Phi = (\forall x)\varphi$ similar

Outline

- Overviews
- First-Order Logic
- Herbrand Models
- Uniform Notation
- Hintikka's Lemma
- Exercises
- Further Reading

Definition

model $\mathbf{M} = \langle \mathbf{D}, \mathbf{I} \rangle$ for language L is **Herbrand model** if

- 1 \mathbf{D} is set of closed terms of L (which is assumed to be nonempty)
- 2 $t^{\mathbf{I}} = t$ for each closed term t

Remark

assignments in Herbrand model are substitutions

Lemma

if $\mathbf{M} = \langle \mathbf{D}, \mathbf{I} \rangle$ is Herbrand model for L then $t^{\mathbf{I}, \mathbf{A}} = (t\mathbf{A})^{\mathbf{I}}$ for each term t of L

Proof

structural induction on t

- if t is variable v then $t^{\mathbf{I}, \mathbf{A}} = v^{\mathbf{I}, \mathbf{A}} = v^{\mathbf{A}} = v\mathbf{A} = (v\mathbf{A})^{\mathbf{I}} = (t\mathbf{A})^{\mathbf{I}}$

Proof (cont'd)

structural induction on t

- if t is constant symbol c of L then $t^{\mathbf{I}, \mathbf{A}} = c^{\mathbf{I}, \mathbf{A}} = c^{\mathbf{I}} = (c\mathbf{A})^{\mathbf{I}} = (t\mathbf{A})^{\mathbf{I}}$
- if $t = f(t_1, \dots, t_n)$ then

$$t^{\mathbf{I}, \mathbf{A}} = f^{\mathbf{I}}(t_1^{\mathbf{I}, \mathbf{A}}, \dots, t_n^{\mathbf{I}, \mathbf{A}}) = f^{\mathbf{I}}((t_1\mathbf{A})^{\mathbf{I}}, \dots, (t_n\mathbf{A})^{\mathbf{I}}) = [f(t_1\mathbf{A}, \dots, t_n\mathbf{A})]^{\mathbf{I}} = (t\mathbf{A})^{\mathbf{I}}$$

Lemma

if $\mathbf{M} = \langle \mathbf{D}, \mathbf{I} \rangle$ is Herbrand model for L then $\Phi^{\mathbf{I}, \mathbf{A}} = (\Phi\mathbf{A})^{\mathbf{I}}$ for each formula Φ of L

Proof

... exercise ...

Lemma

if $\mathbf{M} = \langle \mathbf{D}, \mathbf{I} \rangle$ is Herbrand model for L then

- 1 $(\forall x)\Phi$ is true in $\mathbf{M} \iff \Phi\{x/d\}$ is true in \mathbf{M} for every $d \in \mathbf{D}$
- 2 $(\exists x)\Phi$ is true in $\mathbf{M} \iff \Phi\{x/d\}$ is true in \mathbf{M} for some $d \in \mathbf{D}$

Proof

... exercise ...



William Craig
(1918–2016)



Jacques Herbrand
(1908–1931)



David Hilbert
(1862–1943)



Jaakko Hintikka
(1929–2015)



Outline

- Overviews
- First-Order Logic
- Herbrand Models
- Uniform Notation
- Hintikka's Lemma
- Exercises
- Further Reading

Definition

universal		existential	
γ	$\gamma(t)$	δ	$\delta(t)$
$(\forall x)\Phi$	$\Phi\{x/t\}$	$(\exists x)\Phi$	$\Phi\{x/t\}$
$\neg(\exists x)\Phi$	$\neg\Phi\{x/t\}$	$\neg(\forall x)\Phi$	$\neg\Phi\{x/t\}$

Lemma

$\gamma \equiv (\forall y)\gamma(y)$ and $\delta \equiv (\exists y)\delta(y)$ are valid, provided y is variable new to γ and δ

Lemma

set S of sentences, sentences γ and δ

- 1 if $S \cup \{\gamma\}$ is satisfiable then $S \cup \{\gamma, \gamma(t)\}$ is satisfiable for any closed term t
- 2 if $S \cup \{\delta\}$ is satisfiable then $S \cup \{\delta, \delta(p)\}$ is satisfiable for any constant symbol p that is new to S and δ

Proof

- 1 suppose $S \cup \{\gamma\}$ is satisfiable in model $\mathbf{M} = \langle \mathbf{D}, \mathbf{I} \rangle$
 $(\forall x)\gamma(x)$ is true in \mathbf{M} (with x new to γ)
 $[\gamma(x)]^{\mathbf{I}, \mathbf{A}}$ is true for every assignment \mathbf{A}
let \mathbf{A} be any assignment such that $x^{\mathbf{A}} = t^{\mathbf{I}}$
 $[\gamma(t)]^{\mathbf{I}, \mathbf{A}} = [\gamma\{x/t\}]^{\mathbf{I}, \mathbf{A}} = [\gamma(x)]^{\mathbf{I}, \mathbf{A}} = t$ (using Lemma on slide 43)
- 2 suppose $S \cup \{\delta\}$ is satisfiable in model $\mathbf{M} = \langle \mathbf{D}, \mathbf{I} \rangle$
 $(\exists x)\delta(x)$ is true in \mathbf{M} (with x new to δ)
 $[\delta(x)]^{\mathbf{I}, \mathbf{A}}$ is true for some assignment \mathbf{A}
construct new model $\mathbf{M}^* = \langle \mathbf{D}, \mathbf{J} \rangle$ with \mathbf{J} identical to \mathbf{I} except $p^{\mathbf{J}} = x^{\mathbf{A}}$
 $S \cup \{\delta\}$ is satisfiable in \mathbf{M}^* and $[\delta(x)]^{\mathbf{J}, \mathbf{A}}$ is true
 $[\delta(p)]^{\mathbf{J}, \mathbf{A}} = [\delta\{x/p\}]^{\mathbf{J}, \mathbf{A}} = [\delta(x)]^{\mathbf{J}, \mathbf{A}} = t$ (using Lemma on slide 43)
 $S \cup \{\delta, \delta(p)\}$ is satisfiable (in \mathbf{M}^*)

Definition

rank $r(X)$ of first-order formula: $r(A) = r(\neg A) = r(\top) = r(\perp) = 0$
 $r(\neg \top) = r(\neg \perp) = 1$ $r(\neg \neg Z) = r(Z) + 1$ $r(\alpha) = r(\alpha_1) + r(\alpha_2) + 1$
 $r(\beta) = r(\beta_1) + r(\beta_2) + 1$ $r(\gamma) = r(\gamma(x)) + 1$ $r(\delta) = r(\delta(x)) + 1$

Theorem (First-Order Structural Induction)

every formula of first-order language L has property Q provided

- *basis step*
every atomic formula and its negation has property Q
- *induction steps*
if X has property Q then $\neg \neg X$ has property Q
if α_1 and α_2 have property Q then α has property Q
if β_1 and β_2 have property Q then β has property Q
if $\gamma(t)$ has property Q for each term t then γ has property Q
if $\delta(t)$ has property Q for each term t then δ has property Q

Lemma

if $\mathbf{M} = \langle \mathbf{D}, \mathbf{I} \rangle$ is Herbrand model for L then

- formula γ of L is true in $\mathbf{M} \iff \gamma(d)$ is true in \mathbf{M} for every $d \in \mathbf{D}$
- formula δ of L is true in $\mathbf{M} \iff \gamma(d)$ is true in \mathbf{M} for some $d \in \mathbf{D}$

Proof

... exercise ...

Outline

- Overviews
- First-Order Logic
- Herbrand Models
- Uniform Notation
- Hintikka's Lemma
 - Model Existence Theorem
 - Compactness
 - Löwenheim-Skolem
 - Logical Consequence
- Exercises
- Further Reading

Definition

set \mathbf{H} of sentences of first-order language L is **first-order Hintikka set**, provided

- 1 for any propositional letter A , not both $A \in \mathbf{H}$ and $\neg A \in \mathbf{H}$
- 2 $\perp \notin \mathbf{H}$, $\neg \top \notin \mathbf{H}$
- 3 if $\neg \neg Z \in \mathbf{H}$ then $Z \in \mathbf{H}$
- 4 if $\alpha \in \mathbf{H}$ then $\alpha_1 \in \mathbf{H}$ and $\alpha_2 \in \mathbf{H}$
- 5 if $\beta \in \mathbf{H}$ then $\beta_1 \in \mathbf{H}$ or $\beta_2 \in \mathbf{H}$
- 6 if $\gamma \in \mathbf{H}$ then $\gamma(t) \in \mathbf{H}$ for every closed term t of L
- 7 if $\delta \in \mathbf{H}$ then $\delta(t) \in \mathbf{H}$ for some closed term t of L

Lemma (Hintikka's Lemma)

if \mathbf{H} is first-order Hintikka set with respect to language L with nonempty set of closed terms then \mathbf{H} is satisfiable in Herbrand model

Proof

- \mathbf{H} is first-order Hintikka set with respect to L
- construct model $\mathbf{M} = \langle \mathbf{D}, \mathbf{I} \rangle$
 - \mathbf{D} is collection of closed terms of L
 - $c^{\mathbf{I}} = c$ for constant symbols c of L
 - $f^{\mathbf{I}}(t_1, \dots, t_n) = f(t_1, \dots, t_n)$ for n -place function symbols f of L and $t_1, \dots, t_n \in \mathbf{D}$
 - $\langle t_1, \dots, t_n \rangle$ belongs to $R^{\mathbf{I}}$ for n -place relation symbols R of L if $R(t_1, \dots, t_n) \in \mathbf{H}$
- $t^{\mathbf{I}} = t$ for each closed term t
- claim: if $X \in \mathbf{H}$ then X is true in \mathbf{M} , for every sentence X of L

Proof (cont'd)

claim: if $X \in \mathbf{H}$ then X is true in \mathbf{M} , for each sentence X of L

induction on X

- \top, \perp : trivial
- suppose $R(t_1, \dots, t_n) \in \mathbf{H}$
 $[R(t_1, \dots, t_n)]^{\mathbf{I}, \mathbf{A}} = \mathbf{t}$ because $\langle t_1^{\mathbf{I}, \mathbf{A}}, \dots, t_n^{\mathbf{I}, \mathbf{A}} \rangle = \langle t_1, \dots, t_n \rangle \in R^{\mathbf{I}}$
- negation of atomic formula is straightforward
- α and β are treated as in propositional case
- suppose $\gamma \in \mathbf{H}$
 $\gamma(t) \in \mathbf{H}$ for every closed term t
 $\gamma(t)$ is true in \mathbf{M} for every $t \in \mathbf{D}$ according to induction hypothesis
 γ is true in \mathbf{M} using Lemma on slide 55
- $\delta \dots$ exercise \dots

Outline

- Overviews
- First-Order Logic
- Herbrand Models
- Uniform Notation
- Hintikka's Lemma
 - Model Existence Theorem
 - Compactness
 - Löwenheim-Skolem
 - Logical Consequence
- Exercises
- Further Reading

Definition

given first-order language $L = L(\mathbf{R}, \mathbf{F}, \mathbf{C})$

- **par** is countably infinite set of constant symbols disjoint from \mathbf{C}
- elements of **par** are called **parameters**
- $L^{\mathbf{par}} = L(\mathbf{R}, \mathbf{F}, \mathbf{C} \cup \mathbf{par})$

Definition

collection \mathcal{C} of sets of sentences of $L^{\mathbf{par}}$ is **first-order consistency property** if \mathcal{C} is propositional consistency property and for each $S \in \mathcal{C}$:

- 6 if $\gamma \in S$ then $S \cup \{\gamma(t)\} \in \mathcal{C}$ for every closed term t of $L^{\mathbf{par}}$
- 7 if $\delta \in S$ then $S \cup \{\delta(p)\} \in \mathcal{C}$ for some parameter p of $L^{\mathbf{par}}$

Theorem (First-Order Model Existence)

if \mathcal{C} is first-order consistency property with respect to L and $S \in \mathcal{C}$ is set of sentences over L then S is satisfiable in Herbrand model with respect to $L^{\mathbf{par}}$

Lemma

every first-order consistency property can be extended to one that is subset closed

Definition

alternate first-order consistency property is collection \mathcal{C} meeting conditions for first-order consistency property, except that condition 7 is replaced by

- 7' if $\delta \in S$ then $S \cup \{\delta(p)\} \in \mathcal{C}$ for every parameter p that is new to S

Definition

- **parameter substitution** is mapping π from set of parameters to itself
- π is extended to (sets of) formulas of L^{par} in the obvious way

Lemma

let \mathcal{C} be first-order consistency property that is closed under subsets and let $\mathcal{C}^+ = \{S \mid S\pi \in \mathcal{C} \text{ for some parameter substitution } \pi\}$

- \mathcal{C}^+ extends \mathcal{C}
- \mathcal{C}^+ is closed under subsets
- \mathcal{C}^+ is alternate first-order consistency property

Lemma

every subset closed alternate first-order consistency property can be extended to one of finite character

Outline

- Overviews
- First-Order Logic
- Herbrand Models
- Uniform Notation
- Hintikka's Lemma
 - Model Existence Theorem
 - Compactness
 - Löwenheim-Skolem
 - Logical Consequence
- Exercises
- Further Reading

Proof of First-Order Model Existence

- extend \mathcal{C} to alternate first-order consistency property \mathcal{C}^* of finite character
- let X_1, X_2, X_3, \dots be enumeration of all sentences of L^{par}
- define sequence S_1, S_2, S_3, \dots of members of \mathcal{C}^* :

$$S_1 = S \quad S_{n+1} = \begin{cases} S_n \cup \{X_n\} & \text{if } S_n \cup \{X_n\} \in \mathcal{C}^* \text{ and } X_n \neq \delta \\ S_n \cup \{X_n, \delta(p)\} & \text{if } S_n \cup \{X_n\} \in \mathcal{C}^* \text{ and } X_n = \delta \\ S_n & \text{otherwise} \end{cases}$$

with fixed parameter p which is new to $S_n \cup \{X_n\}$

- $S_1 \subseteq S_2 \subseteq S_3 \subseteq \dots$ and hence $\mathbf{H} = \bigcup_i S_i$ extends S
- $\mathbf{H} \in \mathcal{C}^*$ (because \mathcal{C}^* is of finite character) and \mathbf{H} is maximal in \mathcal{C}^*
- \mathbf{H} is first-order Hintikka set with respect to L^{par}
- \mathbf{H} is satisfiable by Hintikka's Lemma in Herbrand model with respect to L^{par}
- $S \subseteq \mathbf{H}$ is satisfiable in Herbrand model with respect to L^{par}

Theorem (First-Order Compactness)

if every finite subset of a set S of sentences of first-order language L is satisfiable then S is satisfiable

Proof

- define collection \mathcal{C} of sets of sentences of L^{par} as follows: $W \in \mathcal{C}$ if
 - 1 infinitely many parameters are new to W
 - 2 every finite subset of W is satisfiable
- $S \in \mathcal{C}$
- \mathcal{C} is first-order consistency property
- S is satisfiable according to First-Order Model Existence Theorem

Corollary

any set S of sentences of first-order language L that is satisfiable in arbitrarily large finite models is satisfiable in some infinite model

Proof

- suppose S is satisfiable in arbitrary large finite models
- let R be two-place relation symbol not in L and let L' be L extended with R
- there exist sentences A_2, A_3, \dots involving R such that A_i is not true in any model with less than i elements but can be made true in any domain with at least i elements
- consider $S^* = S \cup \{A_2, A_3, \dots\}$
- every finite subset of S^* is satisfiable
- S^* is satisfiable by First-Order Compactness Theorem
- any model in which S^* is satisfiable must be infinite

Example

sentence

$$A_n = (\exists x_1)(\exists x_2) \cdots (\exists x_n) \left[\bigwedge_{i=1}^n R(x_i, x_i) \wedge \bigwedge_{1 \leq i < j \leq n} \neg R(x_i, x_j) \right]$$

- is not true in any model with less than n elements
- can be made true in any domain with at least n elements

Remark

notion of finiteness cannot be captured using machinery of classical first-order logic

Outline

- Overviews
- First-Order Logic
- Herbrand Models
- Uniform Notation
- Hintikka's Lemma
 - Model Existence Theorem
 - Compactness
 - Löwenheim-Skolem
 - Logical Consequence
- Exercises
- Further Reading

Theorem (Löwenheim-Skolem)

if set S of sentences of first-order language L is satisfiable then S is satisfiable in countable model

Proof

- suppose S is satisfiable
- define collection \mathcal{C} of sets of sentences of L^{par} as follows: $W \in \mathcal{C}$ if
 - 1 infinitely many parameters are new to W
 - 2 W is satisfiable
- $S \in \mathcal{C}$
- \mathcal{C} is first-order consistency property
- Model Existence Theorem: S is satisfiable in Herbrand model with respect to L^{par}
- L^{par} has countable alphabet and hence countably many closed terms



William Craig
(1918–2016)



Jacques Herbrand
(1908–1931)



David Hilbert
(1862–1943)



Jaakko Hintikka
(1929–2015)



Leopold Löwenheim
(1878–1957)



Thoralf Skolem
(1887–1963)



Definition

sentence X is **logical consequence** of set S of **sentences**, $S \models_f X$, if X is true in every model in which all members of S are true

Theorem

$S \models_f X$ if and only if $S_0 \models_f X$ for some finite subset S_0 of S

Proof

\Rightarrow if $S \models_f X$ then $S \cup \{\neg X\}$ is not satisfiable
 some finite subset S' of $S \cup \{\neg X\}$ is not satisfiable by compactness
 we may assume that $S' = S_0 \cup \{\neg X\}$ for finite subset S_0 of S
 $S_0 \models_f X$

\Leftarrow easy

Outline

- Overviews
- First-Order Logic
- Herbrand Models
- Uniform Notation
- Hintikka's Lemma
 - Model Existence Theorem
 - Compactness
 - Löwenheim-Skolem
 - Logical Consequence
- Exercises
- Further Reading

Outline

- Overviews
- First-Order Logic
- Herbrand Models
- Uniform Notation
- Hintikka's Lemma
- Exercises
- Further Reading

Fitting

- Give a translation from propositional logic into first-order logic that is natural in the sense that properties carry over, e.g. that a formula is satisfiable/valid/a contradiction iff so is its translation.
- Exercise 5.2.2
- Bonus Exercise 5.2.4
- Exercise 5.3.2
- Exercise 5.3.9 !
- Exercise 5.4.1 or Exercise 5.4.2
- Bonus Exercise 5.5.2 or Exercise 5.6.3
- Exercise 5.9.2
- Bonus Exercise 5.9.3
- Exercise 5.10.1 or 5.10.3(1,2)

Outline

- Overviews
- First-Order Logic
- Herbrand Models
- Uniform Notation
- Hintikka's Lemma
- Exercises
- Further Reading

Fitting

- Section 5.7
- Section 5.8 !
- Section 5.9 !
- Section 5.10
- Section 6.1 !
- Section 6.3 !
- Section 6.4 !
- Section 6.5 !
- Section 8.2