



Logic

Diana Gründlinger

Aart Middeldorp

Fabian Mitterwallner

Alexander Montag

Johannes Niederhauser

Daniel Rainer

Outline

- 1. Summary of Previous Lecture**
- 2. Symbolic Model Checking**
- 3. Intermezzo**
- 4. Linear-Time Temporal Logic (LTL)**
- 5. Further Reading**

Definitions

boolean function f is

- ▶ **monotone** if $f(x_1, \dots, x_n) \leq f(y_1, \dots, y_n)$ for all $x_1 \leq y_1, \dots, x_n \leq y_n$
- ▶ **self-dual** if $f(x_1, \dots, x_n) = \overline{f(\overline{x}_1, \dots, \overline{x}_n)}$
- ▶ **affine** if $f(x_1, \dots, x_n) = c_0 \oplus c_1 x_1 \oplus \dots \oplus c_n x_n$ for some $c_0, \dots, c_n \in \{0, 1\}$

Theorem (Post's Adequacy Theorem)

set X of boolean functions is adequate if and only if following conditions hold:

- 1 $\exists f_1 \in X$ such that $f_1(0, \dots, 0) \neq 0$
- 2 $\exists f_2 \in X$ such that $f_2(1, \dots, 1) \neq 1$
- 3 $\exists f_3 \in X$ which is not monotone
- 4 $\exists f_4 \in X$ which is not self-dual
- 5 $\exists f_5 \in X$ which is not affine

Definitions

- ▶ **CTL (computation tree logic)** formulas are built from atoms, logical connectives, and temporal connectives **AX**, **EX**, **AF**, **EF**, **AG**, **EG**, **AU**, **EU** according to BNF grammar

$$\varphi ::= \perp \mid \top \mid p \mid (\neg\varphi) \mid (\varphi \wedge \varphi) \mid (\varphi \vee \varphi) \mid (\varphi \rightarrow \varphi) \mid (\text{AX}\varphi) \mid (\text{EX}\varphi) \mid (\text{AF}\varphi) \mid (\text{EF}\varphi) \mid (\text{AG}\varphi) \mid (\text{EG}\varphi) \mid \text{A}[\varphi \text{U}\varphi] \mid \text{E}[\varphi \text{U}\varphi]$$

- ▶ **transition system (model)** is triple $\mathcal{M} = (S, \rightarrow, L)$ with
 - ▶ set of states S
 - ▶ transition relation $\rightarrow \subseteq S \times S$ such that $\forall s \in S \exists t \in S$ with $s \rightarrow t$ ("no deadlock")
 - ▶ labelling function $L: S \rightarrow \mathcal{P}(\text{atoms})$
- ▶ **satisfaction** $\mathcal{M}, s \models \varphi$ of CTL formula φ in state $s \in S$ of model $\mathcal{M} = (S, \rightarrow, L)$ is defined by induction on φ

Definition

CTL formulas φ and ψ are **semantically equivalent** ($\varphi \equiv \psi$) if

$$\mathcal{M}, s \models \varphi \iff \mathcal{M}, s \models \psi$$

for all models $\mathcal{M} = (S, \rightarrow, L)$ and states $s \in S$

Theorem

$$\neg \text{AF } \varphi \equiv \text{EG } \neg \varphi$$

$$\text{AF } \varphi \equiv \text{A}[\top \text{ U } \varphi]$$

$$\neg \text{EF } \varphi \equiv \text{AG } \neg \varphi$$

$$\text{EF } \varphi \equiv \text{E}[\top \text{ U } \varphi]$$

$$\neg \text{AX } \varphi \equiv \text{EX } \neg \varphi$$

$$\text{A}[\varphi \text{ U } \psi] \equiv \neg (\text{E}[\neg \psi \text{ U } (\neg \varphi \wedge \neg \psi)]) \vee \text{EG } \neg \psi$$

Theorem

satisfaction of CTL formulas in finite models is **decidable**

CTL Model Checking Algorithm

input: • model $\mathcal{M} = (S, \rightarrow, L)$ and CTL formula φ

output: • $\{s \in S \mid \mathcal{M}, s \models \varphi\}$

label each state $s \in S$ by those subformulas of φ that are satisfied in s

p label $s \iff p \in L(s)$ $\neg\varphi$ label $s \iff s$ is not labelled with φ

$\varphi \wedge \psi$ label $s \iff s$ is labelled with both φ and ψ

$EX\varphi$ label $s \iff t$ is labelled with φ for some t with $s \rightarrow t$

$EG\varphi$ ① label every s that is labelled with φ

② remove label from $s \iff t$ is not labelled with $EG\varphi$ for all t with $s \rightarrow t$

③ repeat ② until no change

$E[\varphi U \psi]$ label $s \iff$ ① s is labelled with ψ

② s is labelled with φ and t with $E[\varphi U \psi]$ for some t with $s \rightarrow t$

③ repeat ② until no change

Part I: Propositional Logic

algebraic normal forms, binary decision diagrams, conjunctive normal forms, DPLL, Horn formulas, natural deduction, Post's adequacy theorem, resolution, SAT, semantics, sorting networks, soundness and completeness, syntax, Tseitin's transformation

Part II: Predicate Logic

natural deduction, quantifier equivalences, resolution, semantics, Skolemization, syntax, undecidability, unification

Part III: Model Checking

adequacy, branching-time temporal logic, CTL*, fairness, linear-time temporal logic, model checking algorithms, symbolic model checking

Part I: Propositional Logic

algebraic normal forms, binary decision diagrams, conjunctive normal forms, DPLL, Horn formulas, natural deduction, Post's adequacy theorem, resolution, SAT, semantics, sorting networks, soundness and completeness, syntax, Tseitin's transformation

Part II: Predicate Logic

natural deduction, quantifier equivalences, resolution, semantics, Skolemization, syntax, undecidability, unification

Part III: Model Checking

adequacy, branching-time temporal logic, CTL*, fairness, **linear-time temporal logic**, model checking algorithms, **symbolic model checking**

Outline

1. Summary of Previous Lecture
- 2. Symbolic Model Checking**
3. Intermezzo
4. Linear-Time Temporal Logic (LTL)
5. Further Reading


symbolic model checking = (CTL) model checking with **BDDs**

symbolic model checking = (CTL) model checking with **BDDs**

Questions

- ▶ how to represent sets of states?
- ▶ how to represent transition relation?
- ▶ how to implement model checking algorithm?

Example



model $\mathcal{M} = (S, \rightarrow, L)$

$S = \{1, 2, 3, 4, 5, 6, 7, 8\}$


$L(1) = \{I_A, I_B\}$ $L(5) = \{I_A, P_B\}$

$L(2) = \{P_A, I_B\}$ $L(6) = \{R_A, P_B\}$

$L(3) = \{R_A, I_B\}$ $L(7) = \{R_A, R_B\}$

$L(4) = \{I_A, R_B\}$ $L(8) = \{P_A, R_B\}$

Example



model $\mathcal{M} = (S, \rightarrow, L)$

$S = \{1, 2, 3, 4, 5, 6, 7, 8\}$

$L(1) = \{I_A, I_B\}$ $L(5) = \{I_A, P_B\}$


$L(2) = \{P_A, I_B\}$ $L(6) = \{R_A, P_B\}$

$L(3) = \{R_A, I_B\}$ $L(7) = \{R_A, R_B\}$

$L(4) = \{I_A, R_B\}$ $L(8) = \{P_A, R_B\}$

- ▶ 8 states require 3 boolean variables

Example



model $\mathcal{M} = (S, \rightarrow, L)$

$S = \{1, 2, 3, 4, 5, 6, 7, 8\}$

$L(1) = \{I_A, I_B\}$ $L(5) = \{I_A, P_B\}$

$L(2) = \{P_A, I_B\}$ $L(6) = \{R_A, P_B\}$

$L(3) = \{R_A, I_B\}$ $L(7) = \{R_A, R_B\}$

$L(4) = \{I_A, R_B\}$ $L(8) = \{P_A, R_B\}$

- ▶ 8 states require 3 boolean variables

state	x	y	z		state	x	y	z	
1	0	0	0	$\bar{x}\bar{y}\bar{z}$	5	1	0	0	$x\bar{y}\bar{z}$
2	0	0	1	$\bar{x}\bar{y}z$	6	1	0	1	$x\bar{y}z$
3	0	1	0	$\bar{x}y\bar{z}$	7	1	1	0	$xy\bar{z}$
4	0	1	1	$\bar{x}yz$	8	1	1	1	xyz

Example (cont'd)

state	state	state	state
1 $\bar{x}\bar{y}\bar{z}$	3 $\bar{x}y\bar{z}$	5 $x\bar{y}\bar{z}$	7 $xy\bar{z}$
2 $\bar{x}y\bar{z}$	4 $\bar{x}yz$	6 $x\bar{y}z$	8 xyz

variable ordering

$[x, y, z]$

set of states $\{1\}$

boolean function $\bar{x}\bar{y}\bar{z}$

reduced OBDD

Example (cont'd)


state	state	state	state
1 $\bar{x}\bar{y}\bar{z}$	3 $\bar{x}y\bar{z}$	5 $x\bar{y}\bar{z}$	7 $xy\bar{z}$
2 $\bar{x}y\bar{z}$	4 $\bar{x}yz$	6 $x\bar{y}z$	8 xyz

variable ordering
[x, y, z]

set of states {1}

boolean function $\bar{x}\bar{y}\bar{z}$

reduced OBDD



Example (cont'd)

state	state	state	state
1 $\bar{x}\bar{y}\bar{z}$	3 $\bar{x}y\bar{z}$	5 $x\bar{y}\bar{z}$	7 $xy\bar{z}$
2 $\bar{x}y\bar{z}$	4 $\bar{x}yz$	6 $x\bar{y}z$	8 xyz

variable ordering
[x, y, z]

set of states


{1}

{1, 2}

boolean function

$\bar{x}\bar{y}\bar{z}$

reduced OBDD



Example (cont'd)

state	state	state	state
1 $\bar{x}\bar{y}\bar{z}$	3 $\bar{x}y\bar{z}$	5 $x\bar{y}\bar{z}$	7 $xy\bar{z}$
2 $\bar{x}y\bar{z}$	4 $\bar{x}yz$	6 $x\bar{y}z$	8 xyz

variable ordering
[x, y, z]

set of states

{1}


{1, 2}

boolean function

$\bar{x}y\bar{z}$

$\bar{x}y$

reduced OBDD



Example (cont'd)

state	state	state	state
1 $\bar{x}\bar{y}\bar{z}$	3 $\bar{x}y\bar{z}$	5 $x\bar{y}\bar{z}$	7 $xy\bar{z}$
2 $\bar{x}y\bar{z}$	4 $\bar{x}yz$	6 $x\bar{y}z$	8 xyz

variable ordering

$[x, y, z]$

set of states

$\{1\}$


$\{1, 2\}$

boolean function

$\bar{x}\bar{y}\bar{z}$

$\bar{x}\bar{y}$

reduced OBDD




Example (cont'd)

state	state	state	state
1 $\bar{x}\bar{y}\bar{z}$	3 $\bar{x}y\bar{z}$	5 $x\bar{y}\bar{z}$	7 $xy\bar{z}$
2 $\bar{x}y\bar{z}$	4 $\bar{x}yz$	6 $x\bar{y}z$	8 xyz

variable ordering
 $[x, y, z]$




set of states	$\{1\}$	$\{1, 2\}$	\emptyset
boolean function	$\bar{x}\bar{y}\bar{z}$	$\bar{x}\bar{y}$	
reduced OBDD			




Example (cont'd)

state	state	state	state
1 $\bar{x}\bar{y}\bar{z}$	3 $\bar{x}y\bar{z}$	5 $x\bar{y}\bar{z}$	7 $xy\bar{z}$
2 $\bar{x}y\bar{z}$	4 $\bar{x}yz$	6 $x\bar{y}z$	8 xyz

variable ordering
[x, y, z]

set of states	{1}	{1, 2}	\emptyset
boolean function	$\bar{x}\bar{y}\bar{z}$	$\bar{x}\bar{y}$	0
reduced OBDD			




Example (cont'd)

state	state	state	state
1 $\bar{x}\bar{y}\bar{z}$	3 $\bar{x}y\bar{z}$	5 $x\bar{y}\bar{z}$	7 $xy\bar{z}$
2 $\bar{x}y\bar{z}$	4 $\bar{x}yz$	6 $x\bar{y}z$	8 xyz

variable ordering
[x, y, z]

set of states	{1}	{1, 2}	\emptyset	{2, 3, 6}
boolean function	$\bar{x}\bar{y}\bar{z}$	$\bar{x}\bar{y}$	0	
reduced OBDD			0	




Example (cont'd)

state	state	state	state
1 $\bar{x}\bar{y}\bar{z}$	3 $\bar{x}y\bar{z}$	5 $x\bar{y}\bar{z}$	7 $xy\bar{z}$
2 $\bar{x}y\bar{z}$	4 $\bar{x}yz$	6 $x\bar{y}z$	8 xyz

variable ordering
[x, y, z]

set of states	{1}	{1, 2}	\emptyset	{2, 3, 6}
boolean function	$\bar{x}\bar{y}\bar{z}$	$\bar{x}\bar{y}$	0	$\bar{y}z + \bar{x}y\bar{z}$
reduced OBDD			0	



0

Example (cont'd)

state	state	state	state
1 $\bar{x}\bar{y}\bar{z}$	3 $\bar{x}y\bar{z}$	5 $xy\bar{z}$	7 xyz
2 $\bar{x}y\bar{z}$	4 $\bar{x}yz$	6 $xy\bar{z}$	8 xyz

variable ordering
 $[x, y, z]$

set of states

$\{1\}$

$\{1, 2\}$

\emptyset

$\{2, 3, 6\}$

boolean function



$\bar{x}\bar{y}\bar{z}$

$\bar{x}\bar{y}$

0


$\bar{y}z + \bar{x}y\bar{z}$

reduced OBDD



Example (cont'd)

state	state	state	state
1 $\bar{x}\bar{y}\bar{z}$	3 $\bar{x}y\bar{z}$	5 $xy\bar{z}$	7 $xy\bar{z}$
2 $\bar{x}y\bar{z}$	4 $\bar{x}yz$	6 $x\bar{y}\bar{z}$	8 xyz




Example (cont'd)

state	state	state	state
1 $\bar{x}\bar{y}\bar{z}$	3 $\bar{x}y\bar{z}$	5 $xy\bar{z}$	7 $xy\bar{z}$
2 $\bar{x}\bar{y}z$	4 $\bar{x}yz$	6 $x\bar{y}z$	8 xyz

transition

1 \rightarrow 3 $\bar{x}\bar{y}\bar{z}\bar{x}'y'\bar{z}'$




Example (cont'd)

state	state	state	state
1 $\bar{x}\bar{y}\bar{z}$	3 $\bar{x}y\bar{z}$	5 $xy\bar{z}$	7 $xy\bar{z}$
2 $\bar{x}\bar{y}z$	4 $\bar{x}yz$	6 $x\bar{y}z$	8 xyz

transition

$1 \rightarrow 3$	$\bar{x}\bar{y}\bar{z}\bar{x}'y'\bar{z}'$
$1 \rightarrow 4$	$\bar{x}\bar{y}\bar{z}\bar{x}'y'z'$




Example (cont'd)

state	state	state	state
1 $\bar{x}\bar{y}\bar{z}$	3 $\bar{x}y\bar{z}$	5 $xy\bar{z}$	7 $xy\bar{z}$
2 $\bar{x}\bar{y}z$	4 $\bar{x}yz$	6 $x\bar{y}z$	8 xyz

transition


$1 \rightarrow 3$	$\bar{x}\bar{y}\bar{z}\bar{x}'y'\bar{z}'$
$1 \rightarrow 4$	$\bar{x}\bar{y}\bar{z}\bar{x}'y'z'$
$2 \rightarrow 1$	$\bar{x}\bar{y}\bar{z}\bar{x}'\bar{y}'\bar{z}'$



Example (cont'd)

state	state	state	state
1 $\bar{x}\bar{y}\bar{z}$	3 $\bar{x}y\bar{z}$	5 $x\bar{y}\bar{z}$	7 $xy\bar{z}$
2 $\bar{x}\bar{y}z$	4 $\bar{x}yz$	6 $x\bar{y}z$	8 xyz

transition	transition	transition	transition
1 \rightarrow 3 $\bar{x}\bar{y}\bar{z}\bar{x}'y'\bar{z}'$	4 \rightarrow 7 $\bar{x}yzx'y'\bar{z}'$		
1 \rightarrow 4 $\bar{x}\bar{y}\bar{z}\bar{x}'y'z'$	5 \rightarrow 1 $x\bar{y}\bar{z}\bar{x}'y'\bar{z}'$		
2 \rightarrow 1 $\bar{x}\bar{y}\bar{z}\bar{x}'\bar{y}'\bar{z}'$	5 \rightarrow 6 $x\bar{y}\bar{z}\bar{x}'y'z'$		
2 \rightarrow 8 $\bar{x}\bar{y}\bar{z}x'y'z'$	6 \rightarrow 3 $x\bar{y}\bar{z}\bar{x}'y'\bar{z}'$		
3 \rightarrow 2 $\bar{x}y\bar{z}\bar{x}'\bar{y}'z'$	7 \rightarrow 6 $xy\bar{z}\bar{x}'y'z'$		
3 \rightarrow 7 $\bar{x}y\bar{z}x'y'\bar{z}'$	7 \rightarrow 8 $xy\bar{z}x'y'z'$		
4 \rightarrow 5 $\bar{x}yzx'\bar{y}'\bar{z}'$	8 \rightarrow 4 $xyz\bar{x}'y'z'$		




Example (cont'd)

state	state	state	state
1 $\bar{x}\bar{y}\bar{z}$	3 $\bar{x}y\bar{z}$	5 $xy\bar{z}$	7 $xy\bar{z}$
2 $\bar{x}\bar{y}z$	4 $\bar{x}yz$	6 $x\bar{y}z$	8 xyz

transition relation

$$\begin{aligned}
 & \bar{x}\bar{y}(\bar{z}\bar{x}'y' + z(\bar{x}'\bar{y}'\bar{z}' + x'y'z')) \\
 & + \bar{x}\bar{y}(\bar{z}(\bar{x}'\bar{y}'z' + x'y'\bar{z}') + zx'\bar{z}') \\
 & + x\bar{y}(\bar{z}(\bar{x}'\bar{y}'\bar{z}' + x'\bar{y}'z') + z\bar{x}'y'\bar{z}') \\
 & + xy(\bar{z}x'z' + z\bar{x}'y'z')
 \end{aligned}$$




Example (cont'd)

state	state	state	state
1 $\bar{x}\bar{y}\bar{z}$	3 $\bar{x}y\bar{z}$	5 $x\bar{y}\bar{z}$	7 $xy\bar{z}$
2 $\bar{x}\bar{y}z$	4 $\bar{x}yz$	6 $x\bar{y}z$	8 xyz

transition relation

$$\begin{aligned}
 & \bar{x}\bar{y}(\bar{z}\bar{x}'y' + z(\bar{x}'\bar{y}'\bar{z}' + x'y'z')) \\
 & + \bar{x}\bar{y}(\bar{z}(\bar{x}'\bar{y}'z' + x'y'\bar{z}') + zx'\bar{z}') \\
 & + x\bar{y}(\bar{z}(\bar{x}'\bar{y}'\bar{z}' + x'\bar{y}'z') + z\bar{x}'y'\bar{z}') \\
 & + xy(\bar{z}x'z' + z\bar{x}'y'z')
 \end{aligned}$$

reduced OBDD with variable ordering $[x, y, z, x', y', z']$ has 24 nodes




Example (cont'd)

state	state	state	state
1 $\bar{x}\bar{y}\bar{z}$	3 $\bar{x}y\bar{z}$	5 $x\bar{y}\bar{z}$	7 $xy\bar{z}$
2 $\bar{x}\bar{y}z$	4 $\bar{x}yz$	6 $x\bar{y}z$	8 xyz

transition relation

$$\begin{aligned}
 & \bar{x}\bar{y}(\bar{z}\bar{x}'y' + z(\bar{x}'\bar{y}'\bar{z}' + x'y'z')) \\
 & + \bar{x}\bar{y}(\bar{z}(\bar{x}'\bar{y}'z' + x'y'\bar{z}') + zx'\bar{z}') \\
 & + x\bar{y}(\bar{z}(\bar{x}'\bar{y}'\bar{z}' + x'\bar{y}'z') + z\bar{x}'y'\bar{z}') \\
 & + xy(\bar{z}x'z' + z\bar{x}'y'z')
 \end{aligned}$$

reduced OBDD with variable ordering $[x, y, z, x', y', z']$ has 24 nodes (B_{\rightarrow})



Definition

model $\mathcal{M} = (S, \rightarrow, L)$

► $[[\varphi]] = \{s \in S \mid \mathcal{M}, s \models \varphi\}$

Definition

model $\mathcal{M} = (S, \rightarrow, L)$

► $[[\varphi]] = \{s \in S \mid \mathcal{M}, s \models \varphi\}$

Lemma

$$[[\top]] = S$$

Definition

model $\mathcal{M} = (S, \rightarrow, L)$

► $[[\varphi]] = \{s \in S \mid \mathcal{M}, s \models \varphi\}$

Lemma

$$[[\top]] = S$$

$$[[\perp]] = \emptyset$$

Definition

model $\mathcal{M} = (S, \rightarrow, L)$

► $[[\varphi]] = \{s \in S \mid \mathcal{M}, s \models \varphi\}$

Lemma

$$[[\top]] = S$$

$$[[\perp]] = \emptyset$$

$$[[p]] = \{s \in S \mid p \in L(s)\}$$

Definition

model $\mathcal{M} = (S, \rightarrow, L)$

► $\llbracket \varphi \rrbracket = \{s \in S \mid \mathcal{M}, s \models \varphi\}$

Lemma

$$\llbracket \top \rrbracket = S$$

$$\llbracket \perp \rrbracket = \emptyset$$

$$\llbracket \neg \varphi \rrbracket = S - \llbracket \varphi \rrbracket$$

$$\llbracket p \rrbracket = \{s \in S \mid p \in L(s)\}$$

Definition

model $\mathcal{M} = (S, \rightarrow, L)$

► $\llbracket \varphi \rrbracket = \{s \in S \mid \mathcal{M}, s \models \varphi\}$

Lemma

$$\llbracket \top \rrbracket = S$$

$$\llbracket \perp \rrbracket = \emptyset$$

$$\llbracket \neg \varphi \rrbracket = S - \llbracket \varphi \rrbracket$$

$$\llbracket \varphi \wedge \psi \rrbracket = \llbracket \varphi \rrbracket \cap \llbracket \psi \rrbracket$$

$$\llbracket p \rrbracket = \{s \in S \mid p \in L(s)\}$$

Definition

model $\mathcal{M} = (S, \rightarrow, L)$

► $\llbracket \varphi \rrbracket = \{s \in S \mid \mathcal{M}, s \models \varphi\}$

Lemma

$$\llbracket \top \rrbracket = S$$

$$\llbracket \perp \rrbracket = \emptyset$$

$$\llbracket \neg \varphi \rrbracket = S - \llbracket \varphi \rrbracket$$

$$\llbracket \varphi \wedge \psi \rrbracket = \llbracket \varphi \rrbracket \cap \llbracket \psi \rrbracket$$

$$\llbracket \varphi \vee \psi \rrbracket = \llbracket \varphi \rrbracket \cup \llbracket \psi \rrbracket$$

$$\llbracket p \rrbracket = \{s \in S \mid p \in L(s)\}$$

Definition

model $\mathcal{M} = (S, \rightarrow, L)$

► $\llbracket \varphi \rrbracket = \{s \in S \mid \mathcal{M}, s \models \varphi\}$

Lemma

$$\llbracket \top \rrbracket = S$$

$$\llbracket \perp \rrbracket = \emptyset$$

$$\llbracket \neg \varphi \rrbracket = S - \llbracket \varphi \rrbracket$$

$$\llbracket \varphi \wedge \psi \rrbracket = \llbracket \varphi \rrbracket \cap \llbracket \psi \rrbracket$$

$$\llbracket \varphi \vee \psi \rrbracket = \llbracket \varphi \rrbracket \cup \llbracket \psi \rrbracket$$

$$\llbracket \varphi \rightarrow \psi \rrbracket = (S - \llbracket \varphi \rrbracket) \cup \llbracket \psi \rrbracket$$

$$\llbracket p \rrbracket = \{s \in S \mid p \in L(s)\}$$

Definition

model $\mathcal{M} = (S, \rightarrow, L)$ $X \subseteq S$

- ▶ $\llbracket \varphi \rrbracket = \{s \in S \mid \mathcal{M}, s \models \varphi\}$
- ▶ $\text{pre}_V(X) = \{s \in S \mid t \in X \text{ for all } t \text{ with } s \rightarrow t\}$

Lemma

$$\llbracket \top \rrbracket = S$$

$$\llbracket \perp \rrbracket = \emptyset$$

$$\llbracket \neg \varphi \rrbracket = S - \llbracket \varphi \rrbracket$$

$$\llbracket \varphi \wedge \psi \rrbracket = \llbracket \varphi \rrbracket \cap \llbracket \psi \rrbracket$$

$$\llbracket \varphi \vee \psi \rrbracket = \llbracket \varphi \rrbracket \cup \llbracket \psi \rrbracket$$

$$\llbracket \varphi \rightarrow \psi \rrbracket = (S - \llbracket \varphi \rrbracket) \cup \llbracket \psi \rrbracket$$

$$\llbracket p \rrbracket = \{s \in S \mid p \in L(s)\}$$

$$\llbracket \text{AX} \varphi \rrbracket = \text{pre}_V(\llbracket \varphi \rrbracket)$$

Definition

model $\mathcal{M} = (S, \rightarrow, L)$ $X \subseteq S$

- ▶ $\llbracket \varphi \rrbracket = \{s \in S \mid \mathcal{M}, s \models \varphi\}$
- ▶ $\text{pre}_\forall(X) = \{s \in S \mid t \in X \text{ for all } t \text{ with } s \rightarrow t\}$
- ▶ $\text{pre}_\exists(X) = \{s \in S \mid s \rightarrow t \text{ for some } t \in X\}$

Lemma

$$\llbracket \top \rrbracket = S$$

$$\llbracket \perp \rrbracket = \emptyset$$

$$\llbracket \neg \varphi \rrbracket = S - \llbracket \varphi \rrbracket$$

$$\llbracket \varphi \wedge \psi \rrbracket = \llbracket \varphi \rrbracket \cap \llbracket \psi \rrbracket$$

$$\llbracket \varphi \vee \psi \rrbracket = \llbracket \varphi \rrbracket \cup \llbracket \psi \rrbracket$$

$$\llbracket \varphi \rightarrow \psi \rrbracket = (S - \llbracket \varphi \rrbracket) \cup \llbracket \psi \rrbracket$$

$$\llbracket p \rrbracket = \{s \in S \mid p \in L(s)\}$$

$$\llbracket \text{AX } \varphi \rrbracket = \text{pre}_\forall(\llbracket \varphi \rrbracket)$$

$$\llbracket \text{EX } \varphi \rrbracket = \text{pre}_\exists(\llbracket \varphi \rrbracket)$$

Definition

model $\mathcal{M} = (S, \rightarrow, L)$ $X \subseteq S$

- ▶ $\llbracket \varphi \rrbracket = \{s \in S \mid \mathcal{M}, s \models \varphi\}$
- ▶ $\text{pre}_{\forall}(X) = \{s \in S \mid t \in X \text{ for all } t \text{ with } s \rightarrow t\}$
- ▶ $\text{pre}_{\exists}(X) = \{s \in S \mid s \rightarrow t \text{ for some } t \in X\}$

Lemma

$$\llbracket \top \rrbracket = S$$

$$\llbracket \perp \rrbracket = \emptyset$$

$$\llbracket \neg \varphi \rrbracket = S - \llbracket \varphi \rrbracket$$

$$\llbracket \varphi \wedge \psi \rrbracket = \llbracket \varphi \rrbracket \cap \llbracket \psi \rrbracket$$

$$\llbracket \varphi \vee \psi \rrbracket = \llbracket \varphi \rrbracket \cup \llbracket \psi \rrbracket$$

$$\llbracket \varphi \rightarrow \psi \rrbracket = (S - \llbracket \varphi \rrbracket) \cup \llbracket \psi \rrbracket$$

$$\llbracket p \rrbracket = \{s \in S \mid p \in L(s)\}$$

$$\llbracket \text{AX } \varphi \rrbracket = \text{pre}_{\forall}(\llbracket \varphi \rrbracket)$$

$$\llbracket \text{EX } \varphi \rrbracket = \text{pre}_{\exists}(\llbracket \varphi \rrbracket)$$

$$\text{pre}_{\forall}(X) = S - \text{pre}_{\exists}(S - X)$$

required operations

complement $S - X$

union $X \cup Y$

intersection $X \cap Y$

$\text{pre}_{\exists}(X)$

Symbolic Model Checking Operations

	required operations	BDD representation
complement	$S - X$	$\text{apply}(\oplus, B_S, B_X)$
union	$X \cup Y$	
intersection	$X \cap Y$	
	$\text{pre}_{\exists}(X)$	

Symbolic Model Checking Operations

	required operations	BDD representation
complement	$S - X$	$\text{apply}(\oplus, B_S, B_X)$
union	$X \cup Y$	$\text{apply}(+, B_X, B_Y)$
intersection	$X \cap Y$	
	$\text{pre}_{\exists}(X)$	

Symbolic Model Checking Operations

	required operations	BDD representation
complement	$S - X$	$\text{apply}(\oplus, B_S, B_X)$
union	$X \cup Y$	$\text{apply}(+, B_X, B_Y)$
intersection	$X \cap Y$	$\text{apply}(\cdot, B_X, B_Y)$
	$\text{pre}_{\exists}(X)$	

	required operations	BDD representation
complement	$S - X$	$\text{apply}(\oplus, B_S, B_X)$
union	$X \cup Y$	$\text{apply}(+, B_X, B_Y)$
intersection	$X \cap Y$	$\text{apply}(\cdot, B_X, B_Y)$
	$\text{pre}_{\exists}(X)$	$\text{apply}(\cdot, B_{\rightarrow}, \underbrace{B_{X'}})$ replace x_1, \dots, x_n by x'_1, \dots, x'_n in B_X

	required operations	BDD representation
complement	$S - X$	$\text{apply}(\oplus, B_S, B_X)$
union	$X \cup Y$	$\text{apply}(+, B_X, B_Y)$
intersection	$X \cap Y$	$\text{apply}(\cdot, B_X, B_Y)$
	$\text{pre}_{\exists}(X)$	$\text{exists}(x'_1, \dots (\text{exists}(x'_n, \text{apply}(\cdot, B_{\rightarrow}, \underbrace{B_{X'}})) \dots))$ replace x_1, \dots, x_n by x'_1, \dots, x'_n in B_X

	required operations	BDD representation
complement	$S - X$	$\text{apply}(\oplus, B_S, B_X)$
union	$X \cup Y$	$\text{apply}(+, B_X, B_Y)$
intersection	$X \cap Y$	$\text{apply}(\cdot, B_X, B_Y)$
	$\text{pre}_{\exists}(X)$	$\text{exists}(x'_1, \dots (\text{exists}(x'_n, \text{apply}(\cdot, B_{\rightarrow}, \underbrace{B_{X'}})) \dots))$ replace x_1, \dots, x_n by x'_1, \dots, x'_n in B_X
		$\text{exists}(x', B) = \text{apply}(+, \text{restrict}(0, x', B), \text{restrict}(1, x', B))$

Lemma

$$\llbracket \text{AF } \varphi \rrbracket = \llbracket \varphi \rrbracket \cup \text{pre}_\forall(\llbracket \text{AF } \varphi \rrbracket)$$

Lemma

$$\llbracket \text{AF } \varphi \rrbracket = \llbracket \varphi \rrbracket \cup \text{pre}_{\forall}(\llbracket \text{AF } \varphi \rrbracket)$$

$$\llbracket \text{EF } \varphi \rrbracket = \llbracket \varphi \rrbracket \cup \text{pre}_{\exists}(\llbracket \text{EF } \varphi \rrbracket)$$

Lemma

$$\llbracket \text{AF } \varphi \rrbracket = \llbracket \varphi \rrbracket \cup \text{pre}_{\forall}(\llbracket \text{AF } \varphi \rrbracket)$$

$$\llbracket \text{EF } \varphi \rrbracket = \llbracket \varphi \rrbracket \cup \text{pre}_{\exists}(\llbracket \text{EF } \varphi \rrbracket)$$

$$\llbracket \text{AG } \varphi \rrbracket = \llbracket \varphi \rrbracket \cap \text{pre}_{\forall}(\llbracket \text{AG } \varphi \rrbracket)$$

Lemma

$$\llbracket \text{AF } \varphi \rrbracket = \llbracket \varphi \rrbracket \cup \text{pre}_{\forall}(\llbracket \text{AF } \varphi \rrbracket)$$

$$\llbracket \text{EF } \varphi \rrbracket = \llbracket \varphi \rrbracket \cup \text{pre}_{\exists}(\llbracket \text{EF } \varphi \rrbracket)$$

$$\llbracket \text{AG } \varphi \rrbracket = \llbracket \varphi \rrbracket \cap \text{pre}_{\forall}(\llbracket \text{AG } \varphi \rrbracket)$$

$$\llbracket \text{EG } \varphi \rrbracket = \llbracket \varphi \rrbracket \cap \text{pre}_{\exists}(\llbracket \text{EG } \varphi \rrbracket)$$

Lemma

$$\llbracket \text{AF } \varphi \rrbracket = \llbracket \varphi \rrbracket \cup \text{pre}_{\forall}(\llbracket \text{AF } \varphi \rrbracket)$$

$$\llbracket \text{EF } \varphi \rrbracket = \llbracket \varphi \rrbracket \cup \text{pre}_{\exists}(\llbracket \text{EF } \varphi \rrbracket)$$

$$\llbracket \text{AG } \varphi \rrbracket = \llbracket \varphi \rrbracket \cap \text{pre}_{\forall}(\llbracket \text{AG } \varphi \rrbracket)$$

$$\llbracket \text{EG } \varphi \rrbracket = \llbracket \varphi \rrbracket \cap \text{pre}_{\exists}(\llbracket \text{EG } \varphi \rrbracket)$$

$$\llbracket \text{A}[\varphi \text{ U } \psi] \rrbracket = \llbracket \psi \rrbracket \cup (\llbracket \varphi \rrbracket \cap \text{pre}_{\forall}(\llbracket \text{A}[\varphi \text{ U } \psi] \rrbracket))$$

$$\llbracket \text{E}[\varphi \text{ U } \psi] \rrbracket = \llbracket \psi \rrbracket \cup (\llbracket \varphi \rrbracket \cap \text{pre}_{\exists}(\llbracket \text{E}[\varphi \text{ U } \psi] \rrbracket))$$

Lemma

$$\llbracket \text{AF } \varphi \rrbracket = \llbracket \varphi \rrbracket \cup \text{pre}_{\forall}(\llbracket \text{AF } \varphi \rrbracket)$$

$$\llbracket \text{EF } \varphi \rrbracket = \llbracket \varphi \rrbracket \cup \text{pre}_{\exists}(\llbracket \text{EF } \varphi \rrbracket)$$

$$\llbracket \text{AG } \varphi \rrbracket = \llbracket \varphi \rrbracket \cap \text{pre}_{\forall}(\llbracket \text{AG } \varphi \rrbracket)$$

$$\llbracket \text{EG } \varphi \rrbracket = \llbracket \varphi \rrbracket \cap \text{pre}_{\exists}(\llbracket \text{EG } \varphi \rrbracket)$$

$$\llbracket \text{A}[\varphi \text{ U } \psi] \rrbracket = \llbracket \psi \rrbracket \cup (\llbracket \varphi \rrbracket \cap \text{pre}_{\forall}(\llbracket \text{A}[\varphi \text{ U } \psi] \rrbracket)) \quad \llbracket \text{E}[\varphi \text{ U } \psi] \rrbracket = \llbracket \psi \rrbracket \cup (\llbracket \varphi \rrbracket \cap \text{pre}_{\exists}(\llbracket \text{E}[\varphi \text{ U } \psi] \rrbracket))$$

Remark

- ▶ $\llbracket \text{AF } \varphi \rrbracket$ is **least fixed point** of function $F_{\text{AF}}(X) = \llbracket \varphi \rrbracket \cup \text{pre}_{\forall}(X)$

Lemma

$$\llbracket \text{AF } \varphi \rrbracket = \llbracket \varphi \rrbracket \cup \text{pre}_{\forall}(\llbracket \text{AF } \varphi \rrbracket)$$

$$\llbracket \text{EF } \varphi \rrbracket = \llbracket \varphi \rrbracket \cup \text{pre}_{\exists}(\llbracket \text{EF } \varphi \rrbracket)$$

$$\llbracket \text{AG } \varphi \rrbracket = \llbracket \varphi \rrbracket \cap \text{pre}_{\forall}(\llbracket \text{AG } \varphi \rrbracket)$$

$$\llbracket \text{EG } \varphi \rrbracket = \llbracket \varphi \rrbracket \cap \text{pre}_{\exists}(\llbracket \text{EG } \varphi \rrbracket)$$

$$\llbracket \text{A}[\varphi \text{ U } \psi] \rrbracket = \llbracket \psi \rrbracket \cup (\llbracket \varphi \rrbracket \cap \text{pre}_{\forall}(\llbracket \text{A}[\varphi \text{ U } \psi] \rrbracket)) \quad \llbracket \text{E}[\varphi \text{ U } \psi] \rrbracket = \llbracket \psi \rrbracket \cup (\llbracket \varphi \rrbracket \cap \text{pre}_{\exists}(\llbracket \text{E}[\varphi \text{ U } \psi] \rrbracket))$$

Remark

- ▶ $\llbracket \text{AF } \varphi \rrbracket$ is least fixed point of function $F_{\text{AF}}(X) = \llbracket \varphi \rrbracket \cup \text{pre}_{\forall}(X)$
- ▶ $\llbracket \text{EG } \varphi \rrbracket$ is **greatest fixed point** of function $F_{\text{EG}}(X) = \llbracket \varphi \rrbracket \cap \text{pre}_{\exists}(X)$

Lemma

$$\llbracket \text{AF } \varphi \rrbracket = \llbracket \varphi \rrbracket \cup \text{pre}_\forall(\llbracket \text{AF } \varphi \rrbracket)$$

$$\llbracket \text{EF } \varphi \rrbracket = \llbracket \varphi \rrbracket \cup \text{pre}_\exists(\llbracket \text{EF } \varphi \rrbracket)$$

$$\llbracket \text{AG } \varphi \rrbracket = \llbracket \varphi \rrbracket \cap \text{pre}_\forall(\llbracket \text{AG } \varphi \rrbracket)$$

$$\llbracket \text{EG } \varphi \rrbracket = \llbracket \varphi \rrbracket \cap \text{pre}_\exists(\llbracket \text{EG } \varphi \rrbracket)$$

$$\llbracket \text{A}[\varphi \text{ U } \psi] \rrbracket = \llbracket \psi \rrbracket \cup (\llbracket \varphi \rrbracket \cap \text{pre}_\forall(\llbracket \text{A}[\varphi \text{ U } \psi] \rrbracket)) \quad \llbracket \text{E}[\varphi \text{ U } \psi] \rrbracket = \llbracket \psi \rrbracket \cup (\llbracket \varphi \rrbracket \cap \text{pre}_\exists(\llbracket \text{E}[\varphi \text{ U } \psi] \rrbracket))$$

Remark

- ▶ $\llbracket \text{AF } \varphi \rrbracket$ is least fixed point of function $F_{\text{AF}}(X) = \llbracket \varphi \rrbracket \cup \text{pre}_\forall(X)$
- ▶ $\llbracket \text{EG } \varphi \rrbracket$ is greatest fixed point of function $F_{\text{EG}}(X) = \llbracket \varphi \rrbracket \cap \text{pre}_\exists(X)$

Theorem (Knaster-Tarski)

every **monotone** function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ with $|S| = n$ admits

- ▶ **least fixed point** $\mu F = F^n(\emptyset)$

Lemma

$$\llbracket \text{AF } \varphi \rrbracket = \llbracket \varphi \rrbracket \cup \text{pre}_\forall(\llbracket \text{AF } \varphi \rrbracket)$$

$$\llbracket \text{EF } \varphi \rrbracket = \llbracket \varphi \rrbracket \cup \text{pre}_\exists(\llbracket \text{EF } \varphi \rrbracket)$$

$$\llbracket \text{AG } \varphi \rrbracket = \llbracket \varphi \rrbracket \cap \text{pre}_\forall(\llbracket \text{AG } \varphi \rrbracket)$$

$$\llbracket \text{EG } \varphi \rrbracket = \llbracket \varphi \rrbracket \cap \text{pre}_\exists(\llbracket \text{EG } \varphi \rrbracket)$$

$$\llbracket \text{A}[\varphi \text{ U } \psi] \rrbracket = \llbracket \psi \rrbracket \cup (\llbracket \varphi \rrbracket \cap \text{pre}_\forall(\llbracket \text{A}[\varphi \text{ U } \psi] \rrbracket)) \quad \llbracket \text{E}[\varphi \text{ U } \psi] \rrbracket = \llbracket \psi \rrbracket \cup (\llbracket \varphi \rrbracket \cap \text{pre}_\exists(\llbracket \text{E}[\varphi \text{ U } \psi] \rrbracket))$$

Remark

- ▶ $\llbracket \text{AF } \varphi \rrbracket$ is least fixed point of function $F_{\text{AF}}(X) = \llbracket \varphi \rrbracket \cup \text{pre}_\forall(X)$
- ▶ $\llbracket \text{EG } \varphi \rrbracket$ is greatest fixed point of function $F_{\text{EG}}(X) = \llbracket \varphi \rrbracket \cap \text{pre}_\exists(X)$

Theorem (Knaster-Tarski)

every **monotone** function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ with $|S| = n$ admits

- ▶ least fixed point $\mu F = F^n(\emptyset)$
- ▶ **greatest fixed point** $\nu F = F^n(S)$

Theorem (Knaster-Tarski)

every **monotone** function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ with $|S| = n$ admits

- ▶ least fixed point $\mu F = F^n(\emptyset)$
- ▶ greatest fixed point $\nu F = F^n(S)$

function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ is monotone if $F(X) \subseteq F(Y)$ whenever $X \subseteq Y \subseteq S$

Theorem (Knaster-Tarski)

every **monotone** function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ with $|S| = n$ admits

- ▶ **least fixed point** $\mu F = F^n(\emptyset)$
- ▶ **greatest fixed point** $\nu F = F^n(S)$

function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ is monotone if $F(X) \subseteq F(Y)$ whenever $X \subseteq Y \subseteq S$

Proof

- ▶ $\emptyset \subseteq F(\emptyset)$

Theorem (Knaster-Tarski)

every **monotone** function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ with $|S| = n$ admits

- ▶ **least fixed point** $\mu F = F^n(\emptyset)$
- ▶ **greatest fixed point** $\nu F = F^n(S)$

function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ is monotone if $F(X) \subseteq F(Y)$ whenever $X \subseteq Y \subseteq S$

Proof

- ▶ $\emptyset \subseteq F(\emptyset) \subseteq F(F(\emptyset))$

monotonicity

Theorem (Knaster-Tarski)

every **monotone** function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ with $|S| = n$ admits

- ▶ **least fixed point** $\mu F = F^n(\emptyset)$
- ▶ **greatest fixed point** $\nu F = F^n(S)$

function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ is monotone if $F(X) \subseteq F(Y)$ whenever $X \subseteq Y \subseteq S$

Proof

- ▶ $\emptyset \subseteq F(\emptyset) \subseteq F(F(\emptyset)) \subseteq \dots \subseteq F^n(\emptyset) \subseteq F^{n+1}(\emptyset)$ induction

Theorem (Knaster–Tarski)

every **monotone** function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ with $|S| = n$ admits

- ▶ **least fixed point** $\mu F = F^n(\emptyset)$
- ▶ **greatest fixed point** $\nu F = F^n(S)$

function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ is monotone if $F(X) \subseteq F(Y)$ whenever $X \subseteq Y \subseteq S$

Proof

- ▶ $\emptyset \subseteq F(\emptyset) \subseteq F(F(\emptyset)) \subseteq \dots \subseteq F^n(\emptyset) \subseteq F^{n+1}(\emptyset)$ induction
- ▶ $\exists 0 \leq i \leq n$ such that $F^i(\emptyset) = F^{i+1}(\emptyset)$ $|S| = n$

Theorem (Knaster-Tarski)

every **monotone** function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ with $|S| = n$ admits

- ▶ **least fixed point** $\mu F = F^n(\emptyset)$
- ▶ **greatest fixed point** $\nu F = F^n(S)$

function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ is monotone if $F(X) \subseteq F(Y)$ whenever $X \subseteq Y \subseteq S$

Proof

- ▶ $\emptyset \subseteq F(\emptyset) \subseteq F(F(\emptyset)) \subseteq \dots \subseteq F^n(\emptyset) \subseteq F^{n+1}(\emptyset)$ induction
- ▶ $\exists 0 \leq i \leq n$ such that $F^i(\emptyset) = F^{i+1}(\emptyset) = F(F^i(\emptyset))$ $|S| = n$
- ▶ $F^i(\emptyset)$ is fixed point of F

Theorem (Knaster–Tarski)

every **monotone** function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ with $|S| = n$ admits

- ▶ **least fixed point** $\mu F = F^n(\emptyset)$
- ▶ **greatest fixed point** $\nu F = F^n(S)$

function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ is monotone if $F(X) \subseteq F(Y)$ whenever $X \subseteq Y \subseteq S$

Proof

- ▶ $\emptyset \subseteq F(\emptyset) \subseteq F(F(\emptyset)) \subseteq \dots \subseteq F^n(\emptyset) \subseteq F^{n+1}(\emptyset)$
- ▶ $\exists 0 \leq i \leq n$ such that $F^i(\emptyset) = F^{i+1}(\emptyset) = F(F^i(\emptyset))$
- ▶ $F^n(\emptyset)$ is fixed point of F

induction

$|S| = n$

$F^i(\emptyset) = F^n(\emptyset)$

Theorem (Knaster–Tarski)

every **monotone** function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ with $|S| = n$ admits

- ▶ **least fixed point** $\mu F = F^n(\emptyset)$
- ▶ **greatest fixed point** $\nu F = F^n(S)$

function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ is monotone if $F(X) \subseteq F(Y)$ whenever $X \subseteq Y \subseteq S$

Proof

- ▶ $\emptyset \subseteq F(\emptyset) \subseteq F(F(\emptyset)) \subseteq \dots \subseteq F^n(\emptyset) \subseteq F^{n+1}(\emptyset)$ induction
- ▶ $\exists 0 \leq i \leq n$ such that $F^i(\emptyset) = F^{i+1}(\emptyset) = F(F^i(\emptyset))$ $|S| = n$
- ▶ $F^n(\emptyset)$ is fixed point of F $F^i(\emptyset) = F^n(\emptyset)$
- ▶ assume X is fixed point of F

Theorem (Knaster-Tarski)

every **monotone** function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ with $|S| = n$ admits

- ▶ **least fixed point** $\mu F = F^n(\emptyset)$
- ▶ **greatest fixed point** $\nu F = F^n(S)$

function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ is monotone if $F(X) \subseteq F(Y)$ whenever $X \subseteq Y \subseteq S$

Proof

- ▶ $\emptyset \subseteq F(\emptyset) \subseteq F(F(\emptyset)) \subseteq \dots \subseteq F^n(\emptyset) \subseteq F^{n+1}(\emptyset)$ induction
- ▶ $\exists 0 \leq i \leq n$ such that $F^i(\emptyset) = F^{i+1}(\emptyset) = F(F^i(\emptyset))$ $|S| = n$
- ▶ $F^n(\emptyset)$ is fixed point of F $F^i(\emptyset) = F^n(\emptyset)$
- ▶ assume X is fixed point of F
- ▶ $\emptyset \subseteq X$

Theorem (Knaster-Tarski)

every **monotone** function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ with $|S| = n$ admits

- ▶ **least fixed point** $\mu F = F^n(\emptyset)$
- ▶ **greatest fixed point** $\nu F = F^n(S)$

function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ is monotone if $F(X) \subseteq F(Y)$ whenever $X \subseteq Y \subseteq S$

Proof

- ▶ $\emptyset \subseteq F(\emptyset) \subseteq F(F(\emptyset)) \subseteq \dots \subseteq F^n(\emptyset) \subseteq F^{n+1}(\emptyset)$ induction
- ▶ $\exists 0 \leq i \leq n$ such that $F^i(\emptyset) = F^{i+1}(\emptyset) = F(F^i(\emptyset))$ $|S| = n$
- ▶ $F^n(\emptyset)$ is fixed point of F $F^i(\emptyset) = F^n(\emptyset)$
- ▶ assume X is fixed point of F
- ▶ $F(\emptyset) \subseteq F(X) = X$ monotonicity

Theorem (Knaster–Tarski)

every **monotone** function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ with $|S| = n$ admits

- ▶ **least fixed point** $\mu F = F^n(\emptyset)$
- ▶ **greatest fixed point** $\nu F = F^n(S)$

function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ is monotone if $F(X) \subseteq F(Y)$ whenever $X \subseteq Y \subseteq S$

Proof

- ▶ $\emptyset \subseteq F(\emptyset) \subseteq F(F(\emptyset)) \subseteq \dots \subseteq F^n(\emptyset) \subseteq F^{n+1}(\emptyset)$ induction
- ▶ $\exists 0 \leq i \leq n$ such that $F^i(\emptyset) = F^{i+1}(\emptyset) = F(F^i(\emptyset))$ $|S| = n$
- ▶ $F^n(\emptyset)$ is fixed point of F $F^i(\emptyset) = F^n(\emptyset)$
- ▶ assume X is fixed point of F
- ▶ $F^n(\emptyset) \subseteq X$ induction

Theorem (Knaster-Tarski)

every **monotone** function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ with $|S| = n$ admits

- ▶ **least fixed point** $\mu F = F^n(\emptyset)$
- ▶ **greatest fixed point** $\nu F = F^n(S)$

function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ is monotone if $F(X) \subseteq F(Y)$ whenever $X \subseteq Y \subseteq S$

Proof

- ▶ $\emptyset \subseteq F(\emptyset) \subseteq F(F(\emptyset)) \subseteq \dots \subseteq F^n(\emptyset) \subseteq F^{n+1}(\emptyset)$ induction
- ▶ $\exists 0 \leq i \leq n$ such that $F^i(\emptyset) = F^{i+1}(\emptyset) = F(F^i(\emptyset))$ $|S| = n$
- ▶ $F^n(\emptyset)$ is fixed point of F $F^i(\emptyset) = F^n(\emptyset)$
- ▶ assume X is fixed point of F
- ▶ $F^n(\emptyset) \subseteq X$ induction
- ▶ $F^n(\emptyset)$ is least fixed point of F

Theorem (Knaster-Tarski)

every **monotone** function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ with $|S| = n$ admits

- ▶ least fixed point $\mu F = F^n(\emptyset)$
- ▶ **greatest fixed point** $\nu F = F^n(S)$

function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ is monotone if $F(X) \subseteq F(Y)$ whenever $X \subseteq Y \subseteq S$

Proof

- ▶ $S \supseteq F(S)$

Theorem (Knaster-Tarski)

every **monotone** function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ with $|S| = n$ admits

- ▶ least fixed point $\mu F = F^n(\emptyset)$
- ▶ **greatest fixed point** $\nu F = F^n(S)$

function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ is monotone if $F(X) \subseteq F(Y)$ whenever $X \subseteq Y \subseteq S$

Proof

- ▶ $S \supseteq F(S) \supseteq F(F(S))$

monotonicity

Theorem (Knaster-Tarski)

every **monotone** function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ with $|S| = n$ admits

- ▶ least fixed point $\mu F = F^n(\emptyset)$
- ▶ **greatest fixed point** $\nu F = F^n(S)$

function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ is monotone if $F(X) \subseteq F(Y)$ whenever $X \subseteq Y \subseteq S$

Proof

- ▶ $S \supseteq F(S) \supseteq F(F(S)) \supseteq F^n(S) \supseteq F^{n+1}(S)$

induction

Theorem (Knaster-Tarski)

every **monotone** function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ with $|S| = n$ admits

- ▶ least fixed point $\mu F = F^n(\emptyset)$
- ▶ **greatest fixed point** $\nu F = F^n(S)$

function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ is monotone if $F(X) \subseteq F(Y)$ whenever $X \subseteq Y \subseteq S$

Proof

- ▶ $S \supseteq F(S) \supseteq F(F(S)) \supseteq F^n(S) \supseteq F^{n+1}(S)$
- ▶ $\exists 0 \leq i \leq n$ such that $F^i(S) = F^{i+1}(S)$

induction

$|S| = n$

Theorem (Knaster-Tarski)

every **monotone** function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ with $|S| = n$ admits

- ▶ least fixed point $\mu F = F^n(\emptyset)$
- ▶ **greatest fixed point** $\nu F = F^n(S)$

function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ is monotone if $F(X) \subseteq F(Y)$ whenever $X \subseteq Y \subseteq S$

Proof

- ▶ $S \supseteq F(S) \supseteq F(F(S)) \supseteq F^n(S) \supseteq F^{n+1}(S)$ induction
- ▶ $\exists 0 \leq i \leq n$ such that $F^i(S) = F^{i+1}(S)$ $|S| = n$
- ▶ $F^i(S)$ is fixed point of F

Theorem (Knaster-Tarski)

every **monotone** function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ with $|S| = n$ admits

- ▶ least fixed point $\mu F = F^n(\emptyset)$
- ▶ **greatest fixed point** $\nu F = F^n(S)$

function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ is monotone if $F(X) \subseteq F(Y)$ whenever $X \subseteq Y \subseteq S$

Proof

- ▶ $S \supseteq F(S) \supseteq F(F(S)) \supseteq F^n(S) \supseteq F^{n+1}(S)$
- ▶ $\exists 0 \leq i \leq n$ such that $F^i(S) = F^{i+1}(S)$
- ▶ $F^n(S)$ is fixed point of F

induction

$|S| = n$

$F^i(S) = F^n(S)$

Theorem (Knaster–Tarski)

every **monotone** function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ with $|S| = n$ admits

- ▶ least fixed point $\mu F = F^n(\emptyset)$
- ▶ **greatest fixed point** $\nu F = F^n(S)$

function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ is monotone if $F(X) \subseteq F(Y)$ whenever $X \subseteq Y \subseteq S$

Proof

- ▶ $S \supseteq F(S) \supseteq F(F(S)) \supseteq F^n(S) \supseteq F^{n+1}(S)$
- ▶ $\exists 0 \leq i \leq n$ such that $F^i(S) = F^{i+1}(S)$
- ▶ $F^n(S)$ is fixed point of F
- ▶ assume X is fixed point of F

induction

$|S| = n$

$F^i(S) = F^n(S)$

Theorem (Knaster-Tarski)

every **monotone** function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ with $|S| = n$ admits

- ▶ least fixed point $\mu F = F^n(\emptyset)$
- ▶ **greatest fixed point** $\nu F = F^n(S)$

function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ is monotone if $F(X) \subseteq F(Y)$ whenever $X \subseteq Y \subseteq S$

Proof

- ▶ $S \supseteq F(S) \supseteq F(F(S)) \supseteq F^n(S) \supseteq F^{n+1}(S)$
- ▶ $\exists 0 \leq i \leq n$ such that $F^i(S) = F^{i+1}(S)$
- ▶ $F^n(S)$ is fixed point of F
- ▶ assume X is fixed point of F
- ▶ $S \supseteq X$

induction

$|S| = n$

$F^i(S) = F^n(S)$

Theorem (Knaster-Tarski)

every **monotone** function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ with $|S| = n$ admits

- ▶ least fixed point $\mu F = F^n(\emptyset)$
- ▶ **greatest fixed point** $\nu F = F^n(S)$

function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ is monotone if $F(X) \subseteq F(Y)$ whenever $X \subseteq Y \subseteq S$

Proof

- ▶ $S \supseteq F(S) \supseteq F(F(S)) \supseteq F^n(S) \supseteq F^{n+1}(S)$ induction
- ▶ $\exists 0 \leq i \leq n$ such that $F^i(S) = F^{i+1}(S)$ $|S| = n$
- ▶ $F^n(S)$ is fixed point of F $F^i(S) = F^n(S)$
- ▶ assume X is fixed point of F
- ▶ $F(S) \supseteq F(X) = X$ monotonicity

Theorem (Knaster-Tarski)

every **monotone** function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ with $|S| = n$ admits

- ▶ least fixed point $\mu F = F^n(\emptyset)$
- ▶ **greatest fixed point** $\nu F = F^n(S)$

function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ is monotone if $F(X) \subseteq F(Y)$ whenever $X \subseteq Y \subseteq S$

Proof

- ▶ $S \supseteq F(S) \supseteq F(F(S)) \supseteq F^n(S) \supseteq F^{n+1}(S)$ induction
- ▶ $\exists 0 \leq i \leq n$ such that $F^i(S) = F^{i+1}(S)$ $|S| = n$
- ▶ $F^n(S)$ is fixed point of F $F^i(S) = F^n(S)$
- ▶ assume X is fixed point of F
- ▶ $F^n(S) \supseteq X$ induction

Theorem (Knaster-Tarski)

every **monotone** function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ with $|S| = n$ admits

- ▶ least fixed point $\mu F = F^n(\emptyset)$
- ▶ **greatest fixed point** $\nu F = F^n(S)$

function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ is monotone if $F(X) \subseteq F(Y)$ whenever $X \subseteq Y \subseteq S$

Proof

- ▶ $S \supseteq F(S) \supseteq F(F(S)) \supseteq F^n(S) \supseteq F^{n+1}(S)$ induction
- ▶ $\exists 0 \leq i \leq n$ such that $F^i(S) = F^{i+1}(S)$ $|S| = n$
- ▶ $F^n(S)$ is fixed point of F $F^i(S) = F^n(S)$
- ▶ assume X is fixed point of F
- ▶ $F^n(S) \supseteq X$ induction
- ▶ $F^n(S)$ is greatest fixed point of F


Definition

function $F_{AF}: \mathcal{P}(S) \rightarrow \mathcal{P}(S): F_{AF}(X) = \llbracket \varphi \rrbracket \cup \text{pre}_V(X)$

Example

$$\varphi = I_B$$

$$\llbracket AF I_B \rrbracket =$$




Definition

function $F_{AF}: \mathcal{P}(S) \rightarrow \mathcal{P}(S): F_{AF}(X) = \llbracket \varphi \rrbracket \cup \text{pre}_V(X)$

Example

$$\varphi = I_B \quad X = \emptyset$$

$$\llbracket AF I_B \rrbracket =$$



Definition


function $F_{AF}: \mathcal{P}(S) \rightarrow \mathcal{P}(S): F_{AF}(X) = \llbracket \varphi \rrbracket \cup \text{pre}_V(X)$

Example

$$\varphi = I_B \quad X = \emptyset$$

$$F_{AF}(X) = \llbracket \varphi \rrbracket = \{1, 2, 3\}$$

$$\llbracket AF I_B \rrbracket =$$



Definition

function $F_{AF}: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$: $F_{AF}(X) = \llbracket \varphi \rrbracket \cup \text{pre}_V(X)$


Example

$$\varphi = I_B \quad X = \emptyset$$

$$F_{AF}(X) = \{1, 2, 3\}$$

$$F_{AF}^2(X) = F_{AF}(F_{AF}(X)) = \{1, 2, 3\} \cup \{6\}$$

$$\llbracket AF I_B \rrbracket =$$



Definition

function $F_{AF}: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$: $F_{AF}(X) = \llbracket \varphi \rrbracket \cup \text{pre}_V(X)$

Example


$$\varphi = I_B \quad X = \emptyset$$

$$F_{AF}(X) = \{1, 2, 3\}$$

$$F_{AF}^2(X) = \{1, 2, 3, 6\}$$

$$F_{AF}^3(X) = \{1, 2, 3\} \cup \{5, 6\}$$

$$\llbracket AF I_B \rrbracket =$$



Definition

function $F_{AF}: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$: $F_{AF}(X) = \llbracket \varphi \rrbracket \cup \text{pre}_V(X)$

Example

$$\varphi = I_B \quad X = \emptyset$$


$$F_{AF}(X) = \{1, 2, 3\}$$

$$F_{AF}^2(X) = \{1, 2, 3, 6\}$$

$$F_{AF}^3(X) = \{1, 2, 3, 5, 6\}$$

$$F_{AF}^4(X) = \{1, 2, 3\} \cup \{5, 6\}$$

$$\llbracket AF I_B \rrbracket =$$



Definition

function $F_{AF}: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$: $F_{AF}(X) = \llbracket \varphi \rrbracket \cup \text{pre}_V(X)$

Example

$$\varphi = I_B \quad X = \emptyset$$


$$F_{AF}(X) = \{1, 2, 3\}$$

$$F_{AF}^2(X) = \{1, 2, 3, 6\}$$

$$F_{AF}^3(X) = \{1, 2, 3, 5, 6\}$$

$$F_{AF}^4(X) = \{1, 2, 3, 5, 6\}$$

$$\llbracket AF I_B \rrbracket = \{1, 2, 3, 5, 6\}$$




Definition

function $F_{EG}: \mathcal{P}(S) \rightarrow \mathcal{P}(S): F_{EG}(X) = \llbracket \varphi \rrbracket \cap \text{pre}_{\exists}(X)$

Example

$$\varphi = P_A \vee I_B$$

$$\llbracket EG(P_A \vee I_B) \rrbracket =$$



Definition


function $F_{EG}: \mathcal{P}(S) \rightarrow \mathcal{P}(S): F_{EG}(X) = \llbracket \varphi \rrbracket \cap \text{pre}_{\exists}(X)$

Example

$$\varphi = P_A \vee I_B$$

$$X = \{1, 2, 3, 4, 5, 6, 7, 8\} = \text{pre}_{\exists}(X)$$

$$\llbracket EG(P_A \vee I_B) \rrbracket =$$



Definition

function $F_{EG}: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$: $F_{EG}(X) = \llbracket \varphi \rrbracket \cap \text{pre}_{\exists}(X)$


Example

$$\varphi = P_A \vee I_B$$

$$X = \{1, 2, 3, 4, 5, 6, 7, 8\} = \text{pre}_{\exists}(X)$$

$$F_{EG}(X) = \llbracket \varphi \rrbracket = \{1, 2, 3, 8\}$$

$$\llbracket EG(P_A \vee I_B) \rrbracket =$$



Definition

function $F_{EG}: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$: $F_{EG}(X) = \llbracket \varphi \rrbracket \cap \text{pre}_{\exists}(X)$

Example


$$\varphi = P_A \vee I_B$$

$$X = \{1, 2, 3, 4, 5, 6, 7, 8\} = \text{pre}_{\exists}(X)$$

$$F_{EG}(X) = \{1, 2, 3, 8\}$$

$$F_{EG}^2(X) = \{1, 2, 3, 8\} \cap \{1, 2, 3, 5, 6, 7\}$$

$$\llbracket EG(P_A \vee I_B) \rrbracket =$$



Definition

function $F_{EG}: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$: $F_{EG}(X) = \llbracket \varphi \rrbracket \cap \text{pre}_{\exists}(X)$

Example

$$\varphi = P_A \vee I_B$$


$$X = \{1, 2, 3, 4, 5, 6, 7, 8\} = \text{pre}_{\exists}(X)$$

$$F_{EG}(X) = \{1, 2, 3, 8\}$$

$$F_{EG}^2(X) = \{1, 2, 3\}$$

$$F_{EG}^3(X) = \{1, 2, 3, 8\} \cap \{1, 2, 3, 5, 6\}$$

$$\llbracket EG(P_A \vee I_B) \rrbracket =$$



Definition

function $F_{EG}: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$: $F_{EG}(X) = \llbracket \varphi \rrbracket \cap \text{pre}_{\exists}(X)$

Example

$$\varphi = P_A \vee I_B$$


$$X = \{1, 2, 3, 4, 5, 6, 7, 8\} = \text{pre}_{\exists}(X)$$

$$F_{EG}(X) = \{1, 2, 3, 8\}$$

$$F_{EG}^2(X) = \{1, 2, 3\}$$

$$F_{EG}^3(X) = \{1, 2, 3\}$$

$$\llbracket EG(P_A \vee I_B) \rrbracket = \{1, 2, 3\}$$



Definition

function $F_{EU}: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$: $F_{EU}(X) = \llbracket \psi \rrbracket \cup (\llbracket \varphi \rrbracket \cap \text{pre}_{\exists}(X))$

Definition

function $F_{EU}: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$: $F_{EU}(X) = \llbracket \psi \rrbracket \cup (\llbracket \varphi \rrbracket \cap \text{pre}_{\exists}(X))$

Lemma

$\llbracket E[\varphi U \psi] \rrbracket$ is least fixed point of monotone function F_{EU}

Definition

function $F_{\text{EU}}: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$: $F_{\text{EU}}(X) = \llbracket \psi \rrbracket \cup (\llbracket \varphi \rrbracket \cap \text{pre}_{\exists}(X))$

Lemma

$\llbracket E[\varphi \cup \psi] \rrbracket$ is least fixed point of **monotone** function F_{EU}

Algorithm

$W := \llbracket \varphi \rrbracket$;

$X := \emptyset$;

$Y := \llbracket \psi \rrbracket$;

repeat until $X = Y$


$X := Y$;

$Y := Y \cup (W \cap \text{pre}_{\exists}(Y))$

return Y


Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$



Example (Huth and Ryan, Exercise 6.12.2(a))


model $\mathcal{M} = (S, \rightarrow, L)$



state	x	y
0	1	0
1	0	1
2	0	0
-	1	1

Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$




state	x	y
0	1	0
1	0	1
2	0	0
-	1	1

x	x'	y	y'	
0	0	0	0	0
0	0	0	1	1 2 → 1
0	0	1	0	0
0	0	1	1	1 1 → 1
0	1	0	0	1 2 → 0
0	1	0	1	0
0	1	1	0	1 1 → 0
0	1	1	1	0


x	x'	y	y'	
1	0	0	0	1 0 → 2
1	0	0	1	0
1	0	1	0	0
1	0	1	1	0
1	1	0	0	0
1	1	0	1	0
1	1	1	0	0
1	1	1	1	0

Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$




state	x	y
0	1	0
1	0	1
2	0	0
-	1	1



Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$




state	x	y
0	1	0
1	0	1
2	0	0
-	1	1

$$\rightarrow: \bar{x}\bar{x}'y' + \bar{x}x'y' + x\bar{x}'\bar{y}\bar{y}'$$

Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$




state	x	y
0	1	0
1	0	1
2	0	0
-	1	1

$$\rightarrow: \bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + xx'\bar{y}\bar{y}'$$

$$AG(p \vee \neg q)$$

Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$




state	x	y
0	1	0
1	0	1
2	0	0
-	1	1

$$\rightarrow: \bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + xx'\bar{y}\bar{y}'$$

$$AG(p \vee \neg q) \equiv \neg E[\top U \neg p \wedge q]$$

Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$



state	x	y
0	1	0
1	0	1
2	0	0
-	1	1

$$\rightarrow: \bar{x}\bar{x}'y' + \bar{x}x'y' + x\bar{x}'y\bar{y}'$$


$$p: x\bar{y}$$

$$q: \bar{x}y$$

$$AG(p \vee \neg q) \equiv \neg E[\top U \neg p \wedge q]$$

Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$



state	x	y
0	1	0
1	0	1
2	0	0
-	1	1

$$\rightarrow: \bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}\bar{y}'$$

$$p: x\bar{y}$$


$$\top: x\bar{y} + \bar{x}y + \bar{x}\bar{y}$$

$$q: \bar{x}y$$

$$AG(p \vee \neg q) \equiv \neg E[\top U \neg p \wedge q]$$

Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$



state	x	y
0	1	0
1	0	1
2	0	0
-	1	1

$$\rightarrow: \bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}y'$$

$$p: x\bar{y}$$


$$q: \bar{x}y$$

$$\top: x\bar{y} + \bar{x}y + \bar{x}\bar{y} = \bar{x} + \bar{y}$$

$$AG(p \vee \neg q) \equiv \neg E[\top U \neg p \wedge q]$$

Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$



state	x	y
0	1	0
1	0	1
2	0	0
-	1	1

$$\rightarrow: \bar{x}\bar{x}'y' + \bar{x}x'y' + x\bar{x}'\bar{y}\bar{y}'$$


$$p: x\bar{y} \quad S, T: x\bar{y} + \bar{x}y + \bar{x}\bar{y} = \bar{x} + \bar{y}$$

$$q: \bar{x}y$$

$$AG(p \vee \neg q) \equiv \neg E[T U \neg p \wedge q]$$

Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$



state	x	y
0	1	0
1	0	1
2	0	0
-	1	1

$$\rightarrow: \bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}y'$$


$$p: x\bar{y} \quad S, \top: x\bar{y} + \bar{x}y + \bar{x}\bar{y} = \bar{x} + \bar{y}$$

$$q: \bar{x}y \quad \neg p \wedge q: ((\bar{x} + \bar{y}) \oplus x\bar{y}) \cdot \bar{x}y$$

$$AG(p \vee \neg q) \equiv \neg E[\top U \neg p \wedge q]$$

Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$



state	x	y
0	1	0
1	0	1
2	0	0
-	1	1

$$\rightarrow: \bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}\bar{y}'$$


$$p: x\bar{y} \quad S, \top: x\bar{y} + \bar{x}y + \bar{x}\bar{y} = \bar{x} + \bar{y}$$

$$q: \bar{x}y \quad \neg p \wedge q: ((\bar{x} + \bar{y}) \oplus x\bar{y}) \cdot \bar{x}y = \bar{x}y$$

$$AG(p \vee \neg q) \equiv \neg E[\top U \neg p \wedge q]$$

Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$



state	x	y
0	1	0
1	0	1
2	0	0
-	1	1

$$\rightarrow: \bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}y'$$

$$p: x\bar{y} \quad S, \top: x\bar{y} + \bar{x}y + \bar{x}\bar{y} = \bar{x} + \bar{y}$$

$$q: \bar{x}y \quad \neg p \wedge q: ((\bar{x} + \bar{y}) \oplus x\bar{y}) \cdot \bar{x}y = \bar{x}y$$


$$AG(p \vee \neg q) \equiv \neg E[\top U \neg p \wedge q]$$

```

W := [[\top]];
X := \emptyset;
Y := [[\neg p \wedge q]];
repeat until X = Y
  X := Y;
  Y := Y \cup (W \cap pre_{\exists}(Y))
return Y
  
```


Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$



state	x	y
0	1	0
1	0	1
2	0	0
-	1	1

$$\rightarrow: \bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}\bar{y}'$$

$$p: x\bar{y} \quad S, \top: x\bar{y} + \bar{x}y + \bar{x}\bar{y} = \bar{x} + \bar{y}$$

$$q: \bar{x}y \quad \neg p \wedge q: ((\bar{x} + \bar{y}) \oplus x\bar{y}) \cdot \bar{x}y = \bar{x}y$$

$$W: \bar{x} + \bar{y}$$

$$AG(p \vee \neg q) \equiv \neg E[\top U \neg p \wedge q]$$

$W := [\top];$

$X := \emptyset;$

$Y := [\neg p \wedge q];$

repeat until $X = Y$


$X := Y;$

$Y := Y \cup (W \cap \text{pre}_{\exists}(Y))$

return Y

Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$



state	x	y
0	1	0
1	0	1
2	0	0
-	1	1

$$\rightarrow: \bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}\bar{y}'$$

$$p: x\bar{y} \quad S, \top: x\bar{y} + \bar{x}y + \bar{x}\bar{y} = \bar{x} + \bar{y}$$

$$q: \bar{x}y \quad \neg p \wedge q: ((\bar{x} + \bar{y}) \oplus x\bar{y}) \cdot \bar{x}y = \bar{x}y$$

$$W: \bar{x} + \bar{y}$$

$$AG(p \vee \neg q) \equiv \neg E[\top U \neg p \wedge q]$$

$W := \llbracket \top \rrbracket;$

$X := \emptyset;$

$Y := \llbracket \neg p \wedge q \rrbracket;$

repeat until $X = Y$

$X := Y;$

$Y := Y \cup (W \cap \text{pre}_{\exists}(Y))$


return Y

$X_0 \ 0$

$Y_0 \ \bar{x}y$

Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$



state	x	y
0	1	0
1	0	1
2	0	0
-	1	1

$$\rightarrow: \bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}\bar{y}'$$

$$p: x\bar{y} \quad S, \top: x\bar{y} + \bar{x}y + \bar{x}\bar{y} = \bar{x} + \bar{y}$$

$$q: \bar{x}y \quad \neg p \wedge q: ((\bar{x} + \bar{y}) \oplus x\bar{y}) \cdot \bar{x}y = \bar{x}y$$

$$W: \bar{x} + \bar{y}$$

$$AG(p \vee \neg q) \equiv \neg E[\top U \neg p \wedge q]$$

$W := \llbracket \top \rrbracket;$

$X := \emptyset;$

$Y := \llbracket \neg p \wedge q \rrbracket;$

repeat until $X = Y$

$X := Y;$

$Y := Y \cup (W \cap \text{pre}_{\exists}(Y))$


return Y

$X_0 \quad 0 \quad X_1 \quad \bar{x}y$

$Y_0 \quad \bar{x}y \quad Y_1 \quad Y_0 \cup (W \cap \text{pre}_{\exists}(Y_0))$

Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$



state	x	y
0	1	0
1	0	1
2	0	0
-	1	1

$$\rightarrow: \bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}y'$$

$$p: x\bar{y} \quad S, \top: x\bar{y} + \bar{x}y + \bar{x}\bar{y} = \bar{x} + \bar{y}$$

$$q: \bar{x}y \quad \neg p \wedge q: ((\bar{x} + \bar{y}) \oplus x\bar{y}) \cdot \bar{x}y = \bar{x}y$$

$$W: \bar{x} + \bar{y}$$

$$AG(p \vee \neg q) \equiv \neg E[\top U \neg p \wedge q]$$

$W := \{\top\};$

$X := \emptyset;$

$Y := \{\neg p \wedge q\};$

repeat until $X = Y$

$X := Y;$

$Y := Y \cup (W \cap \text{pre}_{\exists}(Y))$

return Y


$$X_0 = \emptyset \quad X_1 = \{\bar{x}y\}$$

$$Y_0 = \{\bar{x}y\} \quad Y_1 = Y_0 \cup (W \cap \text{pre}_{\exists}(Y_0))$$

$$\text{pre}_{\exists}(Y_0) = \{\exists x' \exists y' (\rightarrow \cdot Y_0)\}$$

Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$



state	x	y
0	1	0
1	0	1
2	0	0
-	1	1

$$\rightarrow: \bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}y'$$

$$p: x\bar{y} \quad S, \top: x\bar{y} + \bar{x}y + \bar{x}\bar{y} = \bar{x} + \bar{y}$$

$$q: \bar{x}y \quad \neg p \wedge q: ((\bar{x} + \bar{y}) \oplus x\bar{y}) \cdot \bar{x}y = \bar{x}y$$

$$W: \bar{x} + \bar{y}$$

$$AG(p \vee \neg q) \equiv \neg E[\top U \neg p \wedge q]$$

$W := \llbracket \top \rrbracket;$

$X := \emptyset;$

$Y := \llbracket \neg p \wedge q \rrbracket;$

repeat until $X = Y$

$X := Y;$

$Y := Y \cup (W \cap \text{pre}_{\exists}(Y))$

return Y

$$X_0 \ 0 \quad X_1 \ \bar{x}y$$


$$Y_0 \ \bar{x}y \quad Y_1 \ Y_0 \cup (W \cap \text{pre}_{\exists}(Y_0))$$

$$\text{pre}_{\exists}(Y_0) = \exists x' \exists y' (\rightarrow \cdot Y_0')$$

$$= \exists x' \exists y' (\bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}y') \cdot \bar{x}'y'$$

Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$



state	x	y
0	1	0
1	0	1
2	0	0
-	1	1

$$\rightarrow: \bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}y'$$

$$p: x\bar{y} \quad S, \top: x\bar{y} + \bar{x}y + \bar{x}\bar{y} = \bar{x} + \bar{y}$$

$$q: \bar{x}y \quad \neg p \wedge q: ((\bar{x} + \bar{y}) \oplus x\bar{y}) \cdot \bar{x}y = \bar{x}y$$

$$W: \bar{x} + \bar{y}$$

$$AG(p \vee \neg q) \equiv \neg E[\top U \neg p \wedge q]$$

$W := \{\top\};$

$X := \emptyset;$

$Y := \{\neg p \wedge q\};$

repeat until $X = Y$

$X := Y;$

$Y := Y \cup (W \cap \text{pre}_{\exists}(Y))$

return Y

$$X_0 \quad 0 \quad X_1 \quad \bar{x}y$$

$$Y_0 \quad \bar{x}y \quad Y_1 \quad Y_0 \cup (W \cap \text{pre}_{\exists}(Y_0))$$


$$\text{pre}_{\exists}(Y_0) = \exists x' \exists y' (\rightarrow \cdot Y_0')$$

$$= \exists x' \exists y' (\bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}y') \cdot \bar{x}'y'$$

$$= \exists x' \exists y' \bar{x}\bar{x}'y'$$

Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$



state	x	y
0	1	0
1	0	1
2	0	0
-	1	1

$$\rightarrow: \bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}y'$$

$$p: x\bar{y} \quad S, \top: x\bar{y} + \bar{x}y + \bar{x}\bar{y} = \bar{x} + \bar{y}$$

$$q: \bar{x}y \quad \neg p \wedge q: ((\bar{x} + \bar{y}) \oplus x\bar{y}) \cdot \bar{x}y = \bar{x}y$$

$$W: \bar{x} + \bar{y}$$

$$AG(p \vee \neg q) \equiv \neg E[\top U \neg p \wedge q]$$

$W := \{\top\};$

$X := \emptyset;$

$Y := \{\neg p \wedge q\};$

repeat until $X = Y$

$X := Y;$

$Y := Y \cup (W \cap \text{pre}_{\exists}(Y))$

return Y

$$X_0 = \emptyset \quad X_1 = \{\bar{x}y\}$$

$$Y_0 = \{\bar{x}y\} \quad Y_1 = Y_0 \cup (W \cap \text{pre}_{\exists}(Y_0))$$

$$\text{pre}_{\exists}(Y_0) = \exists x' \exists y' (\rightarrow \cdot Y_0')$$


$$= \exists x' \exists y' (\bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}y') \cdot \bar{x}'y'$$

$$= \exists x' \exists y' \bar{x}\bar{x}'y'$$

$$= \exists x' \bar{x}\bar{x}' = \bar{x}$$

Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$



state	x	y
0	1	0
1	0	1
2	0	0
-	1	1

$$\rightarrow: \bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}y'$$

$$p: x\bar{y} \quad S, \top: x\bar{y} + \bar{x}y + \bar{x}\bar{y} = \bar{x} + \bar{y}$$

$$q: \bar{x}y \quad \neg p \wedge q: ((\bar{x} + \bar{y}) \oplus x\bar{y}) \cdot \bar{x}y = \bar{x}y$$

$$W: \bar{x} + \bar{y}$$

$$AG(p \vee \neg q) \equiv \neg E[\top U \neg p \wedge q]$$

$W := \llbracket \top \rrbracket;$

$X := \emptyset;$

$Y := \llbracket \neg p \wedge q \rrbracket;$

repeat until $X = Y$

$X := Y;$

$Y := Y \cup (W \cap \text{pre}_{\exists}(Y))$

return Y

$$X_0 \quad 0 \quad X_1 \quad \bar{x}y$$

$$Y_0 \quad \bar{x}y \quad Y_1 \quad Y_0 \cup (W \cap \text{pre}_{\exists}(Y_0)) = \bar{x}y + (\bar{x} + \bar{y}) \cdot \bar{x} = \bar{x}$$

$$\text{pre}_{\exists}(Y_0) = \exists x' \exists y' (\rightarrow \cdot Y_0')$$


$$= \exists x' \exists y' (\bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}y') \cdot \bar{x}'y'$$

$$= \exists x' \exists y' \bar{x}\bar{x}'y'$$

$$= \exists x' \bar{x}\bar{x}' = \bar{x}$$

Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$



state	x	y
0	1	0
1	0	1
2	0	0
-	1	1

$$\rightarrow: \bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}y'$$

$$p: x\bar{y} \quad S, \top: x\bar{y} + \bar{x}y + \bar{x}\bar{y} = \bar{x} + \bar{y}$$

$$q: \bar{x}y \quad \neg p \wedge q: ((\bar{x} + \bar{y}) \oplus x\bar{y}) \cdot \bar{x}y = \bar{x}y$$

$$W: \bar{x} + \bar{y}$$

$$AG(p \vee \neg q) \equiv \neg E[\top U \neg p \wedge q]$$

$W := \llbracket \top \rrbracket;$

$X := \emptyset;$

$Y := \llbracket \neg p \wedge q \rrbracket;$

repeat until $X = Y$

$X := Y;$

$Y := Y \cup (W \cap \text{pre}_{\exists}(Y))$


return Y

$$X_0 \ 0 \quad X_1 \ \bar{x}y \quad X_2 \ \bar{x}$$

$$Y_0 \ \bar{x}y \quad Y_1 \ \bar{x} \quad Y_2 \ Y_1 \cup (W \cap \text{pre}_{\exists}(Y_1))$$

Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$



state	x	y
0	1	0
1	0	1
2	0	0
-	1	1

$$\rightarrow: \bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}y'$$

$$p: x\bar{y} \quad S, \top: x\bar{y} + \bar{x}y + \bar{x}\bar{y} = \bar{x} + \bar{y}$$

$$q: \bar{x}y \quad \neg p \wedge q: ((\bar{x} + \bar{y}) \oplus x\bar{y}) \cdot \bar{x}y = \bar{x}y$$

$$W: \bar{x} + \bar{y}$$

$$AG(p \vee \neg q) \equiv \neg E[\top U \neg p \wedge q]$$

$W := \llbracket \top \rrbracket;$

$X := \emptyset;$

$Y := \llbracket \neg p \wedge q \rrbracket;$

repeat until $X = Y$

$X := Y;$

$Y := Y \cup (W \cap \text{pre}_{\exists}(Y))$

return Y


$$X_0 \ 0 \quad X_1 \ \bar{x}y \quad X_2 \ \bar{x}$$

$$Y_0 \ \bar{x}y \quad Y_1 \ \bar{x} \quad Y_2 \ Y_1 \cup (W \cap \text{pre}_{\exists}(Y_1))$$

$$\text{pre}_{\exists}(Y_1) = \exists x' \exists y' (\rightarrow \cdot Y_1')$$

Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$



state	x	y
0	1	0
1	0	1
2	0	0
-	1	1

$$\rightarrow: \bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}y'$$

$$p: x\bar{y} \quad S, \top: x\bar{y} + \bar{x}y + \bar{x}\bar{y} = \bar{x} + \bar{y}$$

$$q: \bar{x}y \quad \neg p \wedge q: ((\bar{x} + \bar{y}) \oplus x\bar{y}) \cdot \bar{x}y = \bar{x}y$$

$$W: \bar{x} + \bar{y}$$

$$AG(p \vee \neg q) \equiv \neg E[\top U \neg p \wedge q]$$

$W := \llbracket \top \rrbracket;$

$X := \emptyset;$

$Y := \llbracket \neg p \wedge q \rrbracket;$

repeat until $X = Y$

$X := Y;$

$Y := Y \cup (W \cap \text{pre}_{\exists}(Y))$

return Y

$$X_0 \ 0 \quad X_1 \ \bar{x}y \quad X_2 \ \bar{x}$$


$$Y_0 \ \bar{x}y \quad Y_1 \ \bar{x} \quad Y_2 \ Y_1 \cup (W \cap \text{pre}_{\exists}(Y_1))$$

$$\text{pre}_{\exists}(Y_1) = \exists x' \exists y' (\rightarrow \cdot Y_1')$$

$$= \exists x' \exists y' (\bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}y') \cdot \bar{x}'$$

Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$



state	x	y
0	1	0
1	0	1
2	0	0
-	1	1

$$\rightarrow: \bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}y'$$

$$p: x\bar{y} \quad S, \top: x\bar{y} + \bar{x}y + \bar{x}\bar{y} = \bar{x} + \bar{y}$$

$$q: \bar{x}y \quad \neg p \wedge q: ((\bar{x} + \bar{y}) \oplus x\bar{y}) \cdot \bar{x}y = \bar{x}y$$

$$W: \bar{x} + \bar{y}$$

$$AG(p \vee \neg q) \equiv \neg E[\top U \neg p \wedge q]$$

$W := \llbracket \top \rrbracket;$

$X := \emptyset;$

$Y := \llbracket \neg p \wedge q \rrbracket;$

repeat until $X = Y$

$X := Y;$

$Y := Y \cup (W \cap \text{pre}_{\exists}(Y))$

return Y

$$X_0 \ 0 \quad X_1 \ \bar{x}y \quad X_2 \ \bar{x}$$

$$Y_0 \ \bar{x}y \quad Y_1 \ \bar{x} \quad Y_2 \ Y_1 \cup (W \cap \text{pre}_{\exists}(Y_1))$$


$$\text{pre}_{\exists}(Y_1) = \exists x' \exists y' (\rightarrow \cdot Y_1')$$

$$= \exists x' \exists y' (\bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}y') \cdot \bar{x}$$

$$= \exists x' \exists y' (\bar{x}\bar{x}'y' + x\bar{x}'\bar{y}y')$$

Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$



state	x	y
0	1	0
1	0	1
2	0	0
-	1	1

$$\rightarrow: \bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}y'$$

$$p: x\bar{y} \quad S, \top: x\bar{y} + \bar{x}y + \bar{x}\bar{y} = \bar{x} + \bar{y}$$

$$q: \bar{x}y \quad \neg p \wedge q: ((\bar{x} + \bar{y}) \oplus x\bar{y}) \cdot \bar{x}y = \bar{x}y$$

$$W: \bar{x} + \bar{y}$$

$$AG(p \vee \neg q) \equiv \neg E[\top U \neg p \wedge q]$$

$W := \llbracket \top \rrbracket;$

$X := \emptyset;$

$Y := \llbracket \neg p \wedge q \rrbracket;$

repeat until $X = Y$

$X := Y;$

$Y := Y \cup (W \cap \text{pre}_{\exists}(Y))$

return Y

$$X_0 \ 0 \quad X_1 \ \bar{x}y \quad X_2 \ \bar{x}$$

$$Y_0 \ \bar{x}y \quad Y_1 \ \bar{x} \quad Y_2 \ Y_1 \cup (W \cap \text{pre}_{\exists}(Y_1))$$

$$\text{pre}_{\exists}(Y_1) = \exists x' \exists y' (\rightarrow \cdot Y_1')$$


$$= \exists x' \exists y' (\bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}y') \cdot \bar{x}$$

$$= \exists x' \exists y' (\bar{x}\bar{x}'y' + x\bar{x}'\bar{y}y')$$

$$= \exists x' (\bar{x}\bar{x}' + x\bar{x}'\bar{y}) = \bar{x} + x\bar{y} = \bar{x} + \bar{y}$$

Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$



state	x	y
0	1	0
1	0	1
2	0	0
-	1	1

$$\rightarrow: \bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}y'$$

$$p: x\bar{y} \quad S, \top: x\bar{y} + \bar{x}y + \bar{x}\bar{y} = \bar{x} + \bar{y}$$

$$q: \bar{x}y \quad \neg p \wedge q: ((\bar{x} + \bar{y}) \oplus x\bar{y}) \cdot \bar{x}y = \bar{x}y$$

$$W: \bar{x} + \bar{y}$$

$$AG(p \vee \neg q) \equiv \neg E[\top U \neg p \wedge q]$$

$W := \llbracket \top \rrbracket;$

$X := \emptyset;$

$Y := \llbracket \neg p \wedge q \rrbracket;$

repeat until $X = Y$

$X := Y;$

$Y := Y \cup (W \cap \text{pre}_{\exists}(Y))$

return Y

$$X_0 \ 0 \quad X_1 \ \bar{x}y \quad X_2 \ \bar{x}$$

$$Y_0 \ \bar{x}y \quad Y_1 \ \bar{x} \quad Y_2 \ Y_1 \cup (W \cap \text{pre}_{\exists}(Y_1)) = \bar{x} + \bar{y}$$

$$\text{pre}_{\exists}(Y_1) = \exists x' \exists y' (\rightarrow \cdot Y_1')$$


$$= \exists x' \exists y' (\bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}y') \cdot \bar{x}$$

$$= \exists x' \exists y' (\bar{x}\bar{x}'y' + x\bar{x}'\bar{y}y')$$

$$= \exists x' (\bar{x}\bar{x}' + x\bar{x}'\bar{y}) = \bar{x} + x\bar{y} = \bar{x} + \bar{y}$$

Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$



state	x	y
0	1	0
1	0	1
2	0	0
-	1	1

$$\rightarrow: \bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}\bar{y}'$$

$$p: x\bar{y} \quad S, \top: x\bar{y} + \bar{x}y + \bar{x}\bar{y} = \bar{x} + \bar{y}$$

$$q: \bar{x}y \quad \neg p \wedge q: ((\bar{x} + \bar{y}) \oplus x\bar{y}) \cdot \bar{x}y = \bar{x}y$$

$$W: \bar{x} + \bar{y}$$

$$AG(p \vee \neg q) \equiv \neg E[\top U \neg p \wedge q]$$

$W := \llbracket \top \rrbracket;$

$X := \emptyset;$

$Y := \llbracket \neg p \wedge q \rrbracket;$

repeat until $X = Y$

$X := Y;$

$Y := Y \cup (W \cap \text{pre}_{\exists}(Y))$


return Y

$$X_0 \quad 0 \quad X_1 \quad \bar{x}y \quad X_2 \quad \bar{x} \quad X_3 \quad \bar{x} + \bar{y}$$

$$Y_0 \quad \bar{x}y \quad Y_1 \quad \bar{x} \quad Y_2 \quad \bar{x} + \bar{y} \quad Y_3 \quad Y_2 \cup (W \cap \text{pre}_{\exists}(Y_2))$$

Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$



state	x	y
0	1	0
1	0	1
2	0	0
-	1	1

$$\rightarrow: \bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}y'$$

$$p: x\bar{y} \quad S, \top: x\bar{y} + \bar{x}y + \bar{x}\bar{y} = \bar{x} + \bar{y}$$

$$q: \bar{x}y \quad \neg p \wedge q: ((\bar{x} + \bar{y}) \oplus x\bar{y}) \cdot \bar{x}y = \bar{x}y$$

$$W: \bar{x} + \bar{y}$$

$$AG(p \vee \neg q) \equiv \neg E[\top U \neg p \wedge q]$$

$W := \llbracket \top \rrbracket;$

$X := \emptyset;$

$Y := \llbracket \neg p \wedge q \rrbracket;$

repeat until $X = Y$

$X := Y;$

$Y := Y \cup (W \cap \text{pre}_{\exists}(Y))$

return Y


$$X_0 \ 0 \quad X_1 \ \bar{x}y \quad X_2 \ \bar{x} \quad X_3 \ \bar{x} + \bar{y}$$

$$Y_0 \ \bar{x}y \quad Y_1 \ \bar{x} \quad Y_2 \ \bar{x} + \bar{y} \quad Y_3 \ Y_2 \cup (W \cap \text{pre}_{\exists}(Y_2))$$

$$\text{pre}_{\exists}(Y_2) = \exists x' \exists y' (\rightarrow \cdot Y_2')$$

Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$



state	x	y
0	1	0
1	0	1
2	0	0
-	1	1

$$\rightarrow: \bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}y'$$

$$p: x\bar{y} \quad S, \top: x\bar{y} + \bar{x}y + \bar{x}\bar{y} = \bar{x} + \bar{y}$$

$$q: \bar{x}y \quad \neg p \wedge q: ((\bar{x} + \bar{y}) \oplus x\bar{y}) \cdot \bar{x}y = \bar{x}y$$

$$W: \bar{x} + \bar{y}$$

$$AG(p \vee \neg q) \equiv \neg E[\top U \neg p \wedge q]$$

$W := \llbracket \top \rrbracket;$

$X := \emptyset;$

$Y := \llbracket \neg p \wedge q \rrbracket;$

repeat until $X = Y$

$X := Y;$

$Y := Y \cup (W \cap \text{pre}_{\exists}(Y))$

return Y

$$X_0 \ 0 \quad X_1 \ \bar{x}y \quad X_2 \ \bar{x} \quad X_3 \ \bar{x} + \bar{y}$$


$$Y_0 \ \bar{x}y \quad Y_1 \ \bar{x} \quad Y_2 \ \bar{x} + \bar{y} \quad Y_3 \ Y_2 \cup (W \cap \text{pre}_{\exists}(Y_2))$$

$$\text{pre}_{\exists}(Y_2) = \exists x' \exists y' (\rightarrow \cdot Y_2')$$

$$= \exists x' \exists y' (\bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}y') \cdot (\bar{x}' + \bar{y}')$$

Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$



state	x	y
0	1	0
1	0	1
2	0	0
-	1	1

$$\rightarrow: \bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}y'$$

$$p: x\bar{y} \quad S, \top: x\bar{y} + \bar{x}y + \bar{x}\bar{y} = \bar{x} + \bar{y}$$

$$q: \bar{x}y \quad \neg p \wedge q: ((\bar{x} + \bar{y}) \oplus x\bar{y}) \cdot \bar{x}y = \bar{x}y$$

$$W: \bar{x} + \bar{y}$$

$$AG(p \vee \neg q) \equiv \neg E[\top U \neg p \wedge q]$$

$W := \llbracket \top \rrbracket;$

$X := \emptyset;$

$Y := \llbracket \neg p \wedge q \rrbracket;$

repeat until $X = Y$

$X := Y;$

$Y := Y \cup (W \cap \text{pre}_{\exists}(Y))$

return Y

$$X_0 \ 0 \quad X_1 \ \bar{x}y \quad X_2 \ \bar{x} \quad X_3 \ \bar{x} + \bar{y}$$

$$Y_0 \ \bar{x}y \quad Y_1 \ \bar{x} \quad Y_2 \ \bar{x} + \bar{y} \quad Y_3 \ Y_2 \cup (W \cap \text{pre}_{\exists}(Y_2))$$


$$\text{pre}_{\exists}(Y_2) = \exists x' \exists y' (\rightarrow \cdot Y_2')$$

$$= \exists x' \exists y' (\bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}y') \cdot (\bar{x}' + \bar{y}')$$

$$= \exists x' \exists y' (\bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}y')$$

Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$



state	x	y
0	1	0
1	0	1
2	0	0
-	1	1

$$\rightarrow: \bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}y'$$

$$p: x\bar{y} \quad S, \top: x\bar{y} + \bar{x}y + \bar{x}\bar{y} = \bar{x} + \bar{y}$$

$$q: \bar{x}y \quad \neg p \wedge q: ((\bar{x} + \bar{y}) \oplus x\bar{y}) \cdot \bar{x}y = \bar{x}y$$

$$W: \bar{x} + \bar{y}$$

$$AG(p \vee \neg q) \equiv \neg E[\top U \neg p \wedge q]$$

$W := \llbracket \top \rrbracket;$

$X := \emptyset;$

$Y := \llbracket \neg p \wedge q \rrbracket;$

repeat until $X = Y$

$X := Y;$

$Y := Y \cup (W \cap \text{pre}_{\exists}(Y))$

return Y

$$X_0 \ 0 \quad X_1 \ \bar{x}y \quad X_2 \ \bar{x} \quad X_3 \ \bar{x} + \bar{y}$$

$$Y_0 \ \bar{x}y \quad Y_1 \ \bar{x} \quad Y_2 \ \bar{x} + \bar{y} \quad Y_3 \ Y_2 \cup (W \cap \text{pre}_{\exists}(Y_2))$$

$$\text{pre}_{\exists}(Y_2) = \exists x' \exists y' (\rightarrow \cdot Y_2')$$


$$= \exists x' \exists y' (\bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}y') \cdot (\bar{x}' + \bar{y}')$$

$$= \exists x' \exists y' (\bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}y')$$

$$= \exists x' (\bar{x}\bar{x}' + \bar{x}x' + x\bar{x}'\bar{y}) = \bar{x} + x\bar{y} = \bar{x} + \bar{y}$$

Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$



state	x	y
0	1	0
1	0	1
2	0	0
-	1	1

$$\rightarrow: \bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}y'$$

$$p: x\bar{y} \quad S, \top: x\bar{y} + \bar{x}y + \bar{x}\bar{y} = \bar{x} + \bar{y}$$

$$q: \bar{x}y \quad \neg p \wedge q: ((\bar{x} + \bar{y}) \oplus x\bar{y}) \cdot \bar{x}y = \bar{x}y$$

$$W: \bar{x} + \bar{y}$$

$$AG(p \vee \neg q) \equiv \neg E[\top U \neg p \wedge q]$$

$W := \llbracket \top \rrbracket;$

$X := \emptyset;$

$Y := \llbracket \neg p \wedge q \rrbracket;$

repeat until $X = Y$

$X := Y;$

$Y := Y \cup (W \cap \text{pre}_{\exists}(Y))$

return Y

$$X_0 \ 0 \quad X_1 \ \bar{x}y \quad X_2 \ \bar{x} \quad X_3 \ \bar{x} + \bar{y}$$

$$Y_0 \ \bar{x}y \quad Y_1 \ \bar{x} \quad Y_2 \ \bar{x} + \bar{y} \quad Y_3 \ \bar{x} + \bar{y}$$

$$\text{pre}_{\exists}(Y_2) = \exists x' \exists y' (\rightarrow \cdot Y_2')$$


$$= \exists x' \exists y' (\bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}y') \cdot (\bar{x}' + \bar{y}')$$

$$= \exists x' \exists y' (\bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}y')$$

$$= \exists x' (\bar{x}\bar{x}' + \bar{x}x' + x\bar{x}'\bar{y}) = \bar{x} + x\bar{y} = \bar{x} + \bar{y}$$

Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$



state	x	y
0	1	0
1	0	1
2	0	0
-	1	1

$$\rightarrow: \bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + xx'\bar{y}\bar{y}'$$

$$p: x\bar{y} \quad S, \top: x\bar{y} + \bar{x}y + \bar{x}\bar{y} = \bar{x} + \bar{y}$$

$$q: \bar{x}y \quad \neg p \wedge q: ((\bar{x} + \bar{y}) \oplus x\bar{y}) \cdot \bar{x}y = \bar{x}y$$

$$W: \bar{x} + \bar{y}$$

$$AG(p \vee \neg q) \equiv \neg E[\top U \neg p \wedge q]$$

$W := \llbracket \top \rrbracket;$

$X := \emptyset;$

$Y := \llbracket \neg p \wedge q \rrbracket;$

repeat until $X = Y$

$X := Y;$

$Y := Y \cup (W \cap \text{pre}_{\exists}(Y))$


return Y

$$X_0 \quad 0 \quad X_1 \quad \bar{x}y \quad X_2 \quad \bar{x} \quad X_3 \quad \bar{x} + \bar{y}$$

$$Y_0 \quad \bar{x}y \quad Y_1 \quad \bar{x} \quad Y_2 \quad \bar{x} + \bar{y} \quad Y_3 \quad \bar{x} + \bar{y} \quad X_3 = Y_3$$

Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$



state	x	y
0	1	0
1	0	1
2	0	0
-	1	1

$$\rightarrow: \bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}\bar{y}'$$

$$p: x\bar{y} \quad S, \top: x\bar{y} + \bar{x}y + \bar{x}\bar{y} = \bar{x} + \bar{y}$$

$$q: \bar{x}y \quad \neg p \wedge q: ((\bar{x} + \bar{y}) \oplus x\bar{y}) \cdot \bar{x}y = \bar{x}y$$

$$W: \bar{x} + \bar{y}$$

$$AG(p \vee \neg q) \equiv \neg E[\top U \neg p \wedge q]$$

$W := \llbracket \top \rrbracket;$

$X := \emptyset;$

$Y := \llbracket \neg p \wedge q \rrbracket;$

repeat until $X = Y$

$X := Y;$

$Y := Y \cup (W \cap \text{pre}_{\exists}(Y))$

return Y


$$X_0 \ 0 \quad X_1 \ \bar{x}y \quad X_2 \ \bar{x} \quad X_3 \ \bar{x} + \bar{y}$$

$$Y_0 \ \bar{x}y \quad Y_1 \ \bar{x} \quad Y_2 \ \bar{x} + \bar{y} \quad Y_3 \ \bar{x} + \bar{y} \quad X_3 = Y_3$$

$$E[\top U \neg p \wedge q]: \bar{x} + \bar{y}$$

Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$



state	x	y
0	1	0
1	0	1
2	0	0
-	1	1

$$\rightarrow: \bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}\bar{y}'$$

$$p: x\bar{y} \quad S, \top: x\bar{y} + \bar{x}y + \bar{x}\bar{y} = \bar{x} + \bar{y}$$

$$q: \bar{x}y \quad \neg p \wedge q: ((\bar{x} + \bar{y}) \oplus x\bar{y}) \cdot \bar{x}y = \bar{x}y$$

$$W: \bar{x} + \bar{y}$$

$$AG(p \vee \neg q) \equiv \neg E[\top U \neg p \wedge q]$$

$W := \llbracket \top \rrbracket;$

$X := \emptyset;$

$Y := \llbracket \neg p \wedge q \rrbracket;$

repeat until $X = Y$

$X := Y;$

$Y := Y \cup (W \cap \text{pre}_{\exists}(Y))$

return Y

$$X_0 \ 0 \quad X_1 \ \bar{x}y \quad X_2 \ \bar{x} \quad X_3 \ \bar{x} + \bar{y}$$


$$Y_0 \ \bar{x}y \quad Y_1 \ \bar{x} \quad Y_2 \ \bar{x} + \bar{y} \quad Y_3 \ \bar{x} + \bar{y} \quad X_3 = Y_3$$

$$E[\top U \neg p \wedge q]: \bar{x} + \bar{y}$$

$$AG(p \vee \neg q): (\bar{x} + \bar{y}) \oplus (\bar{x} + \bar{y})$$

Example (Huth and Ryan, Exercise 6.12.2(a))

model $\mathcal{M} = (S, \rightarrow, L)$



state	x	y
0	1	0
1	0	1
2	0	0
-	1	1

$$\rightarrow: \bar{x}\bar{x}'y' + \bar{x}x'\bar{y}' + x\bar{x}'\bar{y}\bar{y}'$$

$$p: x\bar{y} \quad S, \top: x\bar{y} + \bar{x}y + \bar{x}\bar{y} = \bar{x} + \bar{y}$$

$$q: \bar{x}y \quad \neg p \wedge q: ((\bar{x} + \bar{y}) \oplus x\bar{y}) \cdot \bar{x}y = \bar{x}y$$

$$W: \bar{x} + \bar{y}$$

$$AG(p \vee \neg q) \equiv \neg E[\top U \neg p \wedge q]$$

$W := \llbracket \top \rrbracket;$

$X := \emptyset;$

$Y := \llbracket \neg p \wedge q \rrbracket;$

repeat until $X = Y$

$X := Y;$

$Y := Y \cup (W \cap \text{pre}_{\exists}(Y))$

return Y

$$X_0 \ 0 \quad X_1 \ \bar{x}y \quad X_2 \ \bar{x} \quad X_3 \ \bar{x} + \bar{y}$$

$$Y_0 \ \bar{x}y \quad Y_1 \ \bar{x} \quad Y_2 \ \bar{x} + \bar{y} \quad Y_3 \ \bar{x} + \bar{y} \quad X_3 = Y_3$$

$$E[\top U \neg p \wedge q]: \bar{x} + \bar{y}$$

$$AG(p \vee \neg q): (\bar{x} + \bar{y}) \oplus (\bar{x} + \bar{y}) = 0$$

Outline

1. Summary of Previous Lecture
2. Symbolic Model Checking
- 3. Intermezzo**
4. Linear-Time Temporal Logic (LTL)
5. Further Reading

Question

Which of the following statements about symbolic model checking are true ?

- A** For a model with 2 states the reduced BDD B_{\rightarrow} has at most 5 nodes.
- B** The set $\llbracket p \vee \neg p \rrbracket$ corresponds to the reduced BDD $\boxed{0}$.
- C** Every monotone function $F: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ with $|S| = n$ admits a least fixed point $\mu F = F^n(S)$.
- D** $\llbracket \varphi \rightarrow \perp \rrbracket = (S - \llbracket \varphi \rrbracket)$



Outline

1. Summary of Previous Lecture

2. Symbolic Model Checking

3. Intermezzo

4. Linear-Time Temporal Logic (LTL)

Syntax

Semantics

Example

5. Further Reading

Definitions

- ▶ **LTL (linear-time temporal logic)** formulas are built from
 - ▶ atoms p, q, r, p_1, p_2, \dots
 - ▶ logical connectives $\perp, \top, \neg, \wedge, \vee, \rightarrow$

Definitions

- ▶ **LTL (linear-time temporal logic)** formulas are built from
 - ▶ atoms p, q, r, p_1, p_2, \dots
 - ▶ logical connectives $\perp, \top, \neg, \wedge, \vee, \rightarrow$
 - ▶ **temporal connectives** X, F, G, U, W, R

Definitions

► **LTL (linear-time temporal logic)** formulas are built from

- atoms p, q, r, p_1, p_2, \dots
- logical connectives $\perp, \top, \neg, \wedge, \vee, \rightarrow$
- temporal connectives X, F, G, U, W, R

according to following BNF grammar:

$$\varphi ::= \perp \mid \top \mid p \mid (\neg\varphi) \mid (\varphi \wedge \varphi) \mid (\varphi \vee \varphi) \mid (\varphi \rightarrow \varphi) \mid \\ (X\varphi) \mid (F\varphi) \mid (G\varphi) \mid (\varphi U \varphi) \mid (\varphi W \varphi) \mid (\varphi R \varphi)$$

Definitions

- ▶ LTL (linear-time temporal logic) formulas are built from
 - ▶ atoms p, q, r, p_1, p_2, \dots
 - ▶ logical connectives $\perp, \top, \neg, \wedge, \vee, \rightarrow$
 - ▶ temporal connectives X, F, G, U, W, R

according to following BNF grammar:


$$\varphi ::= \perp \mid \top \mid p \mid (\neg\varphi) \mid (\varphi \wedge \varphi) \mid (\varphi \vee \varphi) \mid (\varphi \rightarrow \varphi) \mid \\ (X\varphi) \mid (F\varphi) \mid (G\varphi) \mid (\varphi U\varphi) \mid (\varphi W\varphi) \mid (\varphi R\varphi)$$

- ▶ notational conventions:
 - ▶ binding precedence $\neg, X, F, G > U, W, R > \wedge, \vee > \rightarrow$
 - ▶ omit outer parentheses
 - ▶ $\rightarrow, \wedge, \vee$ are right-associative

Example

formula $F(p \rightarrow Gr) \vee \neg q U p$

parse tree





Example

formula

$F(p \rightarrow Gr) \vee \neg q Up$

$Fp \rightarrow (Gr \vee \neg q) Up$

parse tree





Example

formula

$F(p \rightarrow Gr) \vee \neg q Up$

$Fp \rightarrow (Gr \vee \neg q) Up$

parse tree



X next state

F \exists future state

W weak until

U until

G \forall states globally

R release

Outline

1. Summary of Previous Lecture

2. Symbolic Model Checking

3. Intermezzo

4. Linear-Time Temporal Logic (LTL)

Syntax

Semantics

Example

5. Further Reading

Definition

- ▶ **path** in model $\mathcal{M} = (S, \rightarrow, L)$ is infinite sequence $s_1 \rightarrow s_2 \rightarrow \dots$

Definition

- ▶ path in model $\mathcal{M} = (S, \rightarrow, L)$ is infinite sequence $s_1 \rightarrow s_2 \rightarrow \dots$
- ▶ \forall paths $\pi = s_1 \rightarrow s_2 \rightarrow \dots \quad \forall i \geq 1 \quad \pi^i = s_i \rightarrow s_{i+1} \rightarrow \dots$

Definition

- ▶ path in model $\mathcal{M} = (S, \rightarrow, L)$ is infinite sequence $s_1 \rightarrow s_2 \rightarrow \dots$
- ▶ \forall paths $\pi = s_1 \rightarrow s_2 \rightarrow \dots \quad \forall i \geq 1 \quad \pi^i = s_i \rightarrow s_{i+1} \rightarrow \dots$

Definition

satisfaction of LTL formula φ with respect to path $\pi = s_1 \rightarrow s_2 \rightarrow \dots$ in model $\mathcal{M} = (S, \rightarrow, L)$

$$\pi \models \varphi$$

Definition

- ▶ path in model $\mathcal{M} = (S, \rightarrow, L)$ is infinite sequence $s_1 \rightarrow s_2 \rightarrow \dots$
- ▶ \forall paths $\pi = s_1 \rightarrow s_2 \rightarrow \dots \quad \forall i \geq 1 \quad \pi^i = s_i \rightarrow s_{i+1} \rightarrow \dots$

Definition

satisfaction of LTL formula φ with respect to path $\pi = s_1 \rightarrow s_2 \rightarrow \dots$ in model $\mathcal{M} = (S, \rightarrow, L)$

$$\pi \models \varphi$$

is defined by induction on φ :

$$\pi \models \top$$

$$\pi \not\models \perp$$

Definition

- ▶ path in model $\mathcal{M} = (S, \rightarrow, L)$ is infinite sequence $s_1 \rightarrow s_2 \rightarrow \dots$
- ▶ \forall paths $\pi = s_1 \rightarrow s_2 \rightarrow \dots \quad \forall i \geq 1 \quad \pi^i = s_i \rightarrow s_{i+1} \rightarrow \dots$

Definition

satisfaction of LTL formula φ with respect to path $\pi = s_1 \rightarrow s_2 \rightarrow \dots$ in model $\mathcal{M} = (S, \rightarrow, L)$

$$\pi \models \varphi$$

is defined by induction on φ :

$$\pi \models \top$$

$$\pi \not\models \perp$$

$$\pi \models p \iff p \in L(s_1)$$

Definition

- ▶ path in model $\mathcal{M} = (S, \rightarrow, L)$ is infinite sequence $s_1 \rightarrow s_2 \rightarrow \dots$
- ▶ \forall paths $\pi = s_1 \rightarrow s_2 \rightarrow \dots \quad \forall i \geq 1 \quad \pi^i = s_i \rightarrow s_{i+1} \rightarrow \dots$

Definition

satisfaction of LTL formula φ with respect to path $\pi = s_1 \rightarrow s_2 \rightarrow \dots$ in model $\mathcal{M} = (S, \rightarrow, L)$

$$\pi \models \varphi$$

is defined by induction on φ :

$$\pi \models \top \qquad \pi \not\models \perp$$

$$\pi \models p \iff p \in L(s_1)$$

$$\pi \models \neg\varphi \iff \pi \not\models \varphi$$

Definition

- ▶ path in model $\mathcal{M} = (S, \rightarrow, L)$ is infinite sequence $s_1 \rightarrow s_2 \rightarrow \dots$
- ▶ \forall paths $\pi = s_1 \rightarrow s_2 \rightarrow \dots \quad \forall i \geq 1 \quad \pi^i = s_i \rightarrow s_{i+1} \rightarrow \dots$

Definition


satisfaction of LTL formula φ with respect to path $\pi = s_1 \rightarrow s_2 \rightarrow \dots$ in model $\mathcal{M} = (S, \rightarrow, L)$

$$\pi \models \varphi$$

is defined by induction on φ :


$$\begin{array}{llll} \pi \models \top & \pi \not\models \perp & \pi \models \varphi \wedge \psi & \iff \pi \models \varphi \text{ and } \pi \models \psi \\ \pi \models p & \iff p \in L(s_1) & \pi \models \varphi \vee \psi & \iff \pi \models \varphi \text{ or } \pi \models \psi \\ \pi \models \neg \varphi & \iff \pi \not\models \varphi & \pi \models \varphi \rightarrow \psi & \iff \pi \not\models \varphi \text{ or } \pi \models \psi \end{array}$$

Example



Example

model \mathcal{M}




$\pi_1 \models I_A$

$\pi_1 = 1 \rightarrow 3 \rightarrow 2 \rightarrow 1 \rightarrow 3 \rightarrow 2 \rightarrow \dots$

Example

model \mathcal{M}




$$\pi_1 = 1 \rightarrow 3 \rightarrow 2 \rightarrow 1 \rightarrow 3 \rightarrow 2 \rightarrow \dots$$

$$\pi_1 \models I_A$$

$$\pi_1 \not\models R_A \wedge I_B$$

Example

model \mathcal{M}



$$\pi_1 = 1 \rightarrow 3 \rightarrow 2 \rightarrow 1 \rightarrow 3 \rightarrow 2 \rightarrow \dots$$


$$\pi_1 \models I_A$$

$$\pi_1 \not\models R_A \wedge I_B$$

$$\pi_1 \not\models I_B \rightarrow P_A \vee R_B$$

Example

model \mathcal{M}



$$\pi_1 = 1 \rightarrow 3 \rightarrow 2 \rightarrow 1 \rightarrow 3 \rightarrow 2 \rightarrow \dots$$

$$\pi_2 = 7 \rightarrow 6 \rightarrow 3 \rightarrow 7 \rightarrow 6 \rightarrow 3 \rightarrow \dots$$

$$\pi_1 \models I_A$$

$$\pi_1 \not\models R_A \wedge I_B$$

$$\pi_1 \not\models I_B \rightarrow P_A \vee R_B$$


$$\pi_2 \models I_A$$

$$\pi_2 \models R_A \wedge I_B$$

$$\pi_2 \models I_B \rightarrow P_A \vee R_B$$

Example

model \mathcal{M}



$$\pi_1 = 1 \rightarrow 3 \rightarrow 2 \rightarrow 1 \rightarrow 3 \rightarrow 2 \rightarrow \dots$$

$$\pi_2 = 7 \rightarrow 6 \rightarrow 3 \rightarrow 7 \rightarrow 6 \rightarrow 3 \rightarrow \dots$$

$$\pi_1 \models I_A$$

$$\pi_2 \not\models I_A$$

$$\pi_1 \not\models R_A \wedge I_B$$


$$\pi_2 \models R_A \wedge I_B$$

$$\pi_1 \not\models I_B \rightarrow P_A \vee R_B$$

$$\pi_2 \models I_B \rightarrow P_A \vee R_B$$

Example

model \mathcal{M}



$$\pi_1 = 1 \rightarrow 3 \rightarrow 2 \rightarrow 1 \rightarrow 3 \rightarrow 2 \rightarrow \dots$$

$$\pi_2 = 7 \rightarrow 6 \rightarrow 3 \rightarrow 7 \rightarrow 6 \rightarrow 3 \rightarrow \dots$$

$$\pi_1 \models I_A$$

$$\pi_2 \not\models I_A$$

$$\pi_1 \not\models R_A \wedge I_B$$


$$\pi_2^6 \models R_A \wedge I_B$$

$$\pi_1 \not\models I_B \rightarrow P_A \vee R_B$$

$$\pi_2 \models I_B \rightarrow P_A \vee R_B$$

Example

model \mathcal{M}



$$\pi_1 = 1 \rightarrow 3 \rightarrow 2 \rightarrow 1 \rightarrow 3 \rightarrow 2 \rightarrow \dots$$

$$\pi_2 = 7 \rightarrow 6 \rightarrow 3 \rightarrow 7 \rightarrow 6 \rightarrow 3 \rightarrow \dots$$

$$\pi_1 \models I_A$$

$$\pi_1 \not\models R_A \wedge I_B$$

$$\pi_1 \not\models I_B \rightarrow P_A \vee R_B$$


$$\pi_2 \not\models I_A$$

$$\pi_2^6 \models R_A \wedge I_B$$

$$\pi_2 \models I_B \rightarrow P_A \vee R_B$$

Example

model \mathcal{M}



$$\pi_1 = 1 \rightarrow 3 \rightarrow 2 \rightarrow 1 \rightarrow 3 \rightarrow 2 \rightarrow \dots$$

$$\pi_2 = 7 \rightarrow 6 \rightarrow 3 \rightarrow 7 \rightarrow 6 \rightarrow 3 \rightarrow \dots$$

special notation for infinite paths:

$$\pi_1 = (132)^\omega \quad \pi_2 = (763)^\omega$$

$$\pi_1 \models I_A$$

$$\pi_1 \not\models R_A \wedge I_B$$

$$\pi_1 \not\models I_B \rightarrow P_A \vee R_B$$

$$\pi_2 \not\models I_A$$

$$\pi_2^6 \models R_A \wedge I_B$$

$$\pi_2 \models I_B \rightarrow P_A \vee R_B$$

Definition

satisfaction of LTL formula φ with respect to path $\pi = s_1 \rightarrow s_2 \rightarrow \dots$ in model $\mathcal{M} = (S, \rightarrow, L)$

$$\pi \models \varphi$$

is defined by induction on φ :

$$\pi \models X\varphi \iff \pi^2 \models \varphi$$

Definition

satisfaction of LTL formula φ with respect to path $\pi = s_1 \rightarrow s_2 \rightarrow \dots$ in model $\mathcal{M} = (S, \rightarrow, L)$

$$\pi \models \varphi$$

is defined by induction on φ :

$$\pi \models X\varphi \iff \pi^2 \models \varphi$$

$$\pi \models F\varphi \iff \exists i \geq 1 \pi^i \models \varphi$$

Definition

satisfaction of LTL formula φ with respect to path $\pi = s_1 \rightarrow s_2 \rightarrow \dots$ in model $\mathcal{M} = (S, \rightarrow, L)$

$$\pi \models \varphi$$

is defined by induction on φ :

$$\pi \models \mathbf{X}\varphi \quad \iff \quad \pi^2 \models \varphi$$

$$\pi \models \mathbf{F}\varphi \quad \iff \quad \exists i \geq 1 \quad \pi^i \models \varphi$$

$$\pi \models \mathbf{G}\varphi \quad \iff \quad \forall i \geq 1 \quad \pi^i \models \varphi$$

Definition

satisfaction of LTL formula φ with respect to path $\pi = s_1 \rightarrow s_2 \rightarrow \dots$ in model $\mathcal{M} = (S, \rightarrow, L)$

$$\pi \models \varphi$$

is defined by induction on φ :

$$\pi \models X\varphi \iff \pi^2 \models \varphi$$

$$\pi \models F\varphi \iff \exists i \geq 1 \pi^i \models \varphi$$

$$\pi \models G\varphi \iff \forall i \geq 1 \pi^i \models \varphi$$

$$\pi \models \varphi U \psi \iff \exists i \geq 1 \pi^i \models \psi \text{ and } \forall j < i \pi^j \models \varphi$$

Definition

satisfaction of LTL formula φ with respect to path $\pi = s_1 \rightarrow s_2 \rightarrow \dots$ in model $\mathcal{M} = (S, \rightarrow, L)$

$$\pi \models \varphi$$

is defined by induction on φ :

$$\pi \models \mathbf{X}\varphi \iff \pi^2 \models \varphi$$

$$\pi \models \mathbf{F}\varphi \iff \exists i \geq 1 \pi^i \models \varphi$$

$$\pi \models \mathbf{G}\varphi \iff \forall i \geq 1 \pi^i \models \varphi$$

$$\pi \models \varphi \mathbf{U} \psi \iff \exists i \geq 1 \pi^i \models \psi \text{ and } \forall j < i \pi^j \models \varphi$$

$$\pi \models \varphi \mathbf{W} \psi \iff (\exists i \geq 1 \pi^i \models \psi \text{ and } \forall j < i \pi^j \models \varphi) \text{ or } \forall i \geq 1 \pi^i \models \varphi$$

Definition

satisfaction of LTL formula φ with respect to path $\pi = s_1 \rightarrow s_2 \rightarrow \dots$ in model $\mathcal{M} = (S, \rightarrow, L)$

$$\pi \models \varphi$$

is defined by induction on φ :

$$\pi \models \mathbf{X}\varphi \iff \pi^2 \models \varphi$$

$$\pi \models \mathbf{F}\varphi \iff \exists i \geq 1 \pi^i \models \varphi$$


$$\pi \models \mathbf{G}\varphi \iff \forall i \geq 1 \pi^i \models \varphi$$

$$\pi \models \varphi \mathbf{U} \psi \iff \exists i \geq 1 \pi^i \models \psi \text{ and } \forall j < i \pi^j \models \varphi$$

$$\pi \models \varphi \mathbf{W} \psi \iff (\exists i \geq 1 \pi^i \models \psi \text{ and } \forall j < i \pi^j \models \varphi) \text{ or } \forall i \geq 1 \pi^i \models \varphi$$

$$\pi \models \varphi \mathbf{R} \psi \iff (\exists i \geq 1 \pi^i \models \varphi \text{ and } \forall j \leq i \pi^j \models \psi) \text{ or } \forall i \geq 1 \pi^i \models \psi$$

Example




$$\pi_1 = (132)^\omega$$

$$\pi_2 = (763)^\omega$$

$$\pi_1 \models X(R_A \vee R_B)$$

Example

model \mathcal{M}



$$\pi_1 = (132)^\omega$$

$$\pi_2 = (763)^\omega$$


$$\pi_1 \models X(R_A \vee R_B)$$

$$\pi_1 \models FP_A$$

$$\pi_1 \models XXP_B$$

Example

model \mathcal{M}



$$\pi_1 = (132)^\omega$$

$$\pi_2 = (763)^\omega$$


$$\pi_1 \models X(R_A \vee R_B)$$

$$\pi_1 \models FP_A$$

$$\pi_1 \not\models XXP_B$$

Example

model \mathcal{M}



$$\pi_1 = (132)^\omega$$

$$\pi_2 = (763)^\omega$$


$$\pi_1 \models X(R_A \vee R_B)$$

$$\pi_1 \models FP_A$$

$$\pi_1 \not\models XXP_B$$

Example

model \mathcal{M}



$$\pi_1 = (132)^\omega$$

$$\pi_2 = (763)^\omega$$

$$\pi_1 \models X(R_A \vee R_B)$$

$$\pi_1 \models F P_A$$

$$\pi_1 \not\models X X P_B$$


$$\pi_2 \models F P_A$$

$$\pi_2 \models G \neg I_A$$

$$\pi_2 \models G F P_B$$

Example

model \mathcal{M}



$$\pi_1 = (132)^\omega$$

$$\pi_2 = (763)^\omega$$

$$\pi_1 \models X(R_A \vee R_B)$$

$$\pi_1 \models F P_A$$

$$\pi_1 \not\models X X P_B$$


$$\pi_2 \not\models F P_A$$

$$\pi_2 \models G \neg I_A$$

$$\pi_2 \models G F P_B$$

Example

model \mathcal{M}



$$\pi_1 = (132)^\omega$$

$$\pi_2 = (763)^\omega$$

$$\pi_1 \models X(R_A \vee R_B)$$

$$\pi_1 \models FP_A$$

$$\pi_1 \not\models XXP_B$$


$$\pi_2 \not\models FP_A$$

$$\pi_2 \models G\neg I_A$$

$$\pi_2 \models GFP_B$$

Example

model \mathcal{M}



$$\pi_1 = (132)^\omega$$

$$\pi_2 = (763)^\omega$$

$$\pi_1 \models X(R_A \vee R_B)$$

$$\pi_1 \models FP_A$$

$$\pi_1 \not\models XXP_B$$


$$\pi_2 \not\models FP_A$$

$$\pi_2 \models G\neg I_A$$

$$\pi_2 \models GFP_B$$

Example

model \mathcal{M}



$$\pi_1 = (132)^\omega$$

$$\pi_2 = (763)^\omega$$

$$\pi_1 \models X(R_A \vee R_B)$$

$$\pi_1 \models FP_A$$

$$\pi_1 \not\models XXP_B$$

$$\pi_2 \not\models FP_A$$

$$\pi_2 \models G\neg I_A$$

$$\pi_2 \models GFP_B$$


$$\pi_1 \models I_A \cup P_A$$

$$\pi_2 \models \neg I_A \text{ W } P_A$$

$$\pi_2 \models P_B \text{ R } R_B$$

Example

model \mathcal{M}



$$\pi_1 = (132)^\omega$$

$$\pi_2 = (763)^\omega$$

$$\pi_1 \models X(R_A \vee R_B)$$

$$\pi_1 \models FP_A$$

$$\pi_1 \not\models XXP_B$$

$$\pi_2 \not\models FP_A$$

$$\pi_2 \models G\neg I_A$$


$$\pi_2 \models GFP_B$$

$$\pi_1 \not\models I_A \cup P_A$$

$$\pi_2 \not\models \neg I_A \text{ W } P_A$$

$$\pi_2 \models P_B \text{ R } R_B$$

Example



$$\pi_1 = (132)^\omega$$

$$\pi_2 = (763)^\omega$$

$$\pi_1 \models X(R_A \vee R_B)$$

$$\pi_1 \models FP_A$$

$$\pi_1 \not\models XXP_B$$

$$\pi_2 \not\models FP_A$$

$$\pi_2 \models G\neg I_A$$

$$\pi_2 \models GFP_B$$


$$\pi_1 \not\models I_A \cup P_A$$

$$\pi_2 \models \neg I_A \text{ W } P_A$$

$$\pi_2 \models P_B \text{ R } R_B$$

Example

model \mathcal{M}



$$\pi_1 = (132)^\omega$$

$$\pi_2 = (763)^\omega$$

$$\pi_1 \models X(R_A \vee R_B)$$

$$\pi_1 \models FP_A$$

$$\pi_1 \not\models XXP_B$$

$$\pi_2 \not\models FP_A$$

$$\pi_2 \models G\neg I_A$$

$$\pi_2 \models GFP_B$$

$$\pi_1 \not\models I_A \cup P_A$$

$$\pi_2 \models \neg I_A \mathcal{W} P_A$$

$$\pi_2 \not\models P_B \mathcal{R} R_B$$

Definition

model $\mathcal{M} = (S, \rightarrow, L)$, state $s \in S$, LTL formula φ


$\mathcal{M}, s \models \varphi \iff \forall \text{ paths } \pi = s \rightarrow \dots \quad \pi \models \varphi$ "formula φ holds in state s of model \mathcal{M} "

Definition

model $\mathcal{M} = (S, \rightarrow, L)$, state $s \in S$, LTL formula φ

$\mathcal{M}, s \models \varphi \iff \forall \text{ paths } \pi = s \rightarrow \dots \quad \pi \models \varphi$ "formula φ holds in state s of model \mathcal{M} "

Example




$\mathcal{M}, 1 \not\models G(R_A \rightarrow F P_A)$

Definition

model $\mathcal{M} = (S, \rightarrow, L)$, state $s \in S$, LTL formula φ

$\mathcal{M}, s \models \varphi \iff \forall \text{ paths } \pi = s \rightarrow \dots \quad \pi \models \varphi$ "formula φ holds in state s of model \mathcal{M} "

Example



$\mathcal{M}, 1 \not\models G (R_A \rightarrow F P_A)$


$\mathcal{M}, 4 \not\models \neg (R_B \cup P_B)$

Definition

model $\mathcal{M} = (S, \rightarrow, L)$, state $s \in S$, LTL formula φ

$\mathcal{M}, s \models \varphi \iff \forall \text{ paths } \pi = s \rightarrow \dots \quad \pi \models \varphi$ "formula φ holds in state s of model \mathcal{M} "

Example



$\mathcal{M}, 1 \not\models G (R_A \rightarrow F P_A)$


$\mathcal{M}, 4 \not\models \neg (R_B \cup P_B)$

Definition

model $\mathcal{M} = (S, \rightarrow, L)$, state $s \in S$, LTL formula φ

$\mathcal{M}, s \models \varphi \iff \forall \text{ paths } \pi = s \rightarrow \dots \quad \pi \models \varphi$ "formula φ holds in state s of model \mathcal{M} "

Example



$\mathcal{M}, 1 \not\models G (R_A \rightarrow F P_A)$

$\mathcal{M}, 4 \not\models \neg (R_B \cup P_B)$


$\mathcal{M}, 4 \models R_B \cup P_B$

Definition

model $\mathcal{M} = (S, \rightarrow, L)$, state $s \in S$, LTL formula φ

$\mathcal{M}, s \models \varphi \iff \forall \text{ paths } \pi = s \rightarrow \dots \quad \pi \models \varphi$ "formula φ holds in state s of model \mathcal{M} "

Example



$\mathcal{M}, 1 \not\models G (R_A \rightarrow F P_A)$

$\mathcal{M}, 4 \not\models \neg (R_B \cup P_B)$


$\mathcal{M}, 4 \not\models R_B \cup P_B$

Definition

model $\mathcal{M} = (S, \rightarrow, L)$, state $s \in S$, LTL formula φ

$\mathcal{M}, s \models \varphi \iff \forall \text{ paths } \pi = s \rightarrow \dots \quad \pi \models \varphi$ "formula φ holds in state s of model \mathcal{M} "

Example



$\mathcal{M}, 1 \not\models G (R_A \rightarrow F P_A)$

$\mathcal{M}, 4 \not\models \neg (R_B \cup P_B)$

$\mathcal{M}, 4 \not\models R_B \cup P_B$


$\mathcal{M}, 6 \not\models X (F I_B \wedge ((X \neg P_B) R R_A))$

Definition

model $\mathcal{M} = (S, \rightarrow, L)$, state $s \in S$, LTL formula φ

$\mathcal{M}, s \models \varphi \iff \forall \text{ paths } \pi = s \rightarrow \dots \quad \pi \models \varphi$ "formula φ holds in state s of model \mathcal{M} "

Example



$\mathcal{M}, 1 \not\models G (R_A \rightarrow F P_A)$

$\mathcal{M}, 4 \not\models \neg (R_B \cup P_B)$

$\mathcal{M}, 4 \not\models R_B \cup P_B$

$\mathcal{M}, 6 \models X (F I_B \wedge ((X \neg P_B) R R_A))$

Definition

LTL formulas φ and ψ are **semantically equivalent** ($\varphi \equiv \psi$) if

\forall models $\mathcal{M} = (S, \rightarrow, L)$

$$\pi \models \varphi \iff \pi \models \psi$$

\forall paths π in \mathcal{M}

Definition

LTL formulas φ and ψ are **semantically equivalent** ($\varphi \equiv \psi$) if

\forall models $\mathcal{M} = (S, \rightarrow, L)$

$$\pi \models \varphi \iff \pi \models \psi$$

\forall paths π in \mathcal{M}

Theorem

$$\neg X\varphi \equiv X\neg\varphi$$

Definition

LTL formulas φ and ψ are **semantically equivalent** ($\varphi \equiv \psi$) if

\forall models $\mathcal{M} = (S, \rightarrow, L)$

$$\pi \models \varphi \iff \pi \models \psi$$

\forall paths π in \mathcal{M}

Theorem

$$\neg X\varphi \equiv X\neg\varphi$$

$$\neg F\varphi \equiv G\neg\varphi$$

$$\neg G\varphi \equiv F\neg\varphi$$

Definition

LTL formulas φ and ψ are **semantically equivalent** ($\varphi \equiv \psi$) if

\forall models $\mathcal{M} = (S, \rightarrow, L)$

$$\pi \models \varphi \iff \pi \models \psi$$

\forall paths π in \mathcal{M}

Theorem

$$\neg X \varphi \equiv X \neg \varphi$$

$$\neg F \varphi \equiv G \neg \varphi$$

$$\neg G \varphi \equiv F \neg \varphi$$

$$\neg(\varphi U \psi) \equiv \neg \varphi R \neg \psi$$

Definition

LTL formulas φ and ψ are **semantically equivalent** ($\varphi \equiv \psi$) if

\forall models $\mathcal{M} = (S, \rightarrow, L)$

$$\pi \models \varphi \iff \pi \models \psi$$

\forall paths π in \mathcal{M}

Theorem

$$\neg X\varphi \equiv X\neg\varphi$$

$$\neg F\varphi \equiv G\neg\varphi$$

$$\neg G\varphi \equiv F\neg\varphi$$

$$\neg(\varphi U \psi) \equiv \neg\varphi R \neg\psi$$

$$\neg(\varphi R \psi) \equiv \neg\varphi U \neg\psi$$

$$F(\varphi \vee \psi) \equiv F\varphi \vee F\psi$$

$$G(\varphi \wedge \psi) \equiv G\varphi \wedge G\psi$$

Definition

LTL formulas φ and ψ are **semantically equivalent** ($\varphi \equiv \psi$) if

\forall models $\mathcal{M} = (S, \rightarrow, L)$

$$\pi \models \varphi \iff \pi \models \psi$$

\forall paths π in \mathcal{M}

Theorem

$$\neg X \varphi \equiv X \neg \varphi$$

$$\neg F \varphi \equiv G \neg \varphi$$

$$\neg G \varphi \equiv F \neg \varphi$$

$$\neg(\varphi U \psi) \equiv \neg \varphi R \neg \psi$$

$$\neg(\varphi R \psi) \equiv \neg \varphi U \neg \psi$$

$$F(\varphi \vee \psi) \equiv F \varphi \vee F \psi$$

$$G(\varphi \wedge \psi) \equiv G \varphi \wedge G \psi$$

$$F \varphi \equiv T U \varphi$$

$$G \varphi \equiv \perp R \varphi$$

Definition

LTL formulas φ and ψ are **semantically equivalent** ($\varphi \equiv \psi$) if

\forall models $\mathcal{M} = (S, \rightarrow, L)$

$$\pi \models \varphi \iff \pi \models \psi$$

\forall paths π in \mathcal{M}

Theorem

$$\neg X \varphi \equiv X \neg \varphi$$

$$\neg F \varphi \equiv G \neg \varphi$$

$$\neg G \varphi \equiv F \neg \varphi$$

$$\neg(\varphi U \psi) \equiv \neg \varphi R \neg \psi$$

$$\neg(\varphi R \psi) \equiv \neg \varphi U \neg \psi$$

$$\varphi U \psi \equiv \varphi W \psi \wedge F \psi$$

$$\varphi W \psi \equiv \varphi U \psi \vee G \varphi$$

$$F(\varphi \vee \psi) \equiv F \varphi \vee F \psi$$

$$G(\varphi \wedge \psi) \equiv G \varphi \wedge G \psi$$

$$F \varphi \equiv T U \varphi$$

$$G \varphi \equiv \perp R \varphi$$

Definition

LTL formulas φ and ψ are **semantically equivalent** ($\varphi \equiv \psi$) if

\forall models $\mathcal{M} = (S, \rightarrow, L)$

$$\pi \models \varphi \iff \pi \models \psi$$

\forall paths π in \mathcal{M}

Theorem

$$\neg X \varphi \equiv X \neg \varphi$$

$$\neg F \varphi \equiv G \neg \varphi$$

$$\neg G \varphi \equiv F \neg \varphi$$

$$\neg(\varphi U \psi) \equiv \neg \varphi R \neg \psi$$

$$\neg(\varphi R \psi) \equiv \neg \varphi U \neg \psi$$

$$\varphi U \psi \equiv \varphi W \psi \wedge F \psi$$

$$\varphi W \psi \equiv \varphi U \psi \vee G \varphi$$

$$F(\varphi \vee \psi) \equiv F \varphi \vee F \psi$$

$$G(\varphi \wedge \psi) \equiv G \varphi \wedge G \psi$$

$$F \varphi \equiv T U \varphi$$

$$G \varphi \equiv \perp R \varphi$$

$$\varphi W \psi \equiv \psi R(\varphi \vee \psi)$$

$$\varphi R \psi \equiv \psi W(\varphi \wedge \psi)$$

Definition

LTL formulas φ and ψ are **semantically equivalent** ($\varphi \equiv \psi$) if

\forall models $\mathcal{M} = (S, \rightarrow, L)$

$$\pi \models \varphi \iff \pi \models \psi$$

\forall paths π in \mathcal{M}

Theorem

$$\neg X \varphi \equiv X \neg \varphi$$

$$\neg F \varphi \equiv G \neg \varphi$$

$$\neg G \varphi \equiv F \neg \varphi$$

$$\neg(\varphi U \psi) \equiv \neg \varphi R \neg \psi$$

$$\neg(\varphi R \psi) \equiv \neg \varphi U \neg \psi$$

$$\varphi U \psi \equiv \varphi W \psi \wedge F \psi$$

$$\varphi W \psi \equiv \varphi U \psi \vee G \varphi$$

$$\varphi U \psi \equiv \neg(\neg \psi U (\neg \varphi \wedge \neg \psi)) \wedge F \psi$$

$$F(\varphi \vee \psi) \equiv F \varphi \vee F \psi$$

$$G(\varphi \wedge \psi) \equiv G \varphi \wedge G \psi$$

$$F \varphi \equiv T U \varphi$$

$$G \varphi \equiv \perp R \varphi$$

$$\varphi W \psi \equiv \psi R(\varphi \vee \psi)$$

$$\varphi R \psi \equiv \psi W(\varphi \wedge \psi)$$

Theorem

$$\varphi \text{ U } \psi \equiv \neg(\neg\psi \text{ U } (\neg\varphi \wedge \neg\psi)) \wedge \text{ F } \psi$$

Theorem

$$\varphi \text{ U } \psi \equiv \neg(\neg\psi \text{ U } (\neg\varphi \wedge \neg\psi)) \wedge \text{ F } \psi$$

$$\pi \models \neg(\neg\psi \text{ U } (\neg\varphi \wedge \neg\psi))$$

Theorem

$$\varphi \text{ U } \psi \equiv \neg(\neg\psi \text{ U } (\neg\varphi \wedge \neg\psi)) \wedge \text{ F } \psi$$

$$\pi \models \neg(\neg\psi \text{ U } (\neg\varphi \wedge \neg\psi)) \iff \pi \not\models \neg\psi \text{ U } (\neg\varphi \wedge \neg\psi)$$

Theorem

$$\varphi \mathbf{U} \psi \equiv \neg(\neg\psi \mathbf{U} (\neg\varphi \wedge \neg\psi)) \wedge \mathbf{F} \psi$$

$$\begin{aligned} \pi \models \neg(\neg\psi \mathbf{U} (\neg\varphi \wedge \neg\psi)) &\iff \pi \not\models \neg\psi \mathbf{U} (\neg\varphi \wedge \neg\psi) \\ &\iff \text{not } \exists i \geq 1 (\pi^i \models \neg\varphi \wedge \neg\psi \text{ and } \forall j < i \pi^j \models \neg\psi) \end{aligned}$$

Theorem

$$\varphi \mathbf{U} \psi \equiv \neg(\neg\psi \mathbf{U} (\neg\varphi \wedge \neg\psi)) \wedge \mathbf{F} \psi$$

$$\begin{aligned} \pi \models \neg(\neg\psi \mathbf{U} (\neg\varphi \wedge \neg\psi)) &\iff \pi \not\models \neg\psi \mathbf{U} (\neg\varphi \wedge \neg\psi) \\ &\iff \text{not } \exists i \geq 1 \left(\pi^i \models \neg\varphi \wedge \neg\psi \text{ and } \forall j < i \pi^j \models \neg\psi \right) \\ &\iff \forall i \geq 1 \text{ not } \left(\pi^i \models \neg\varphi \wedge \neg\psi \text{ and } \forall j < i \pi^j \models \neg\psi \right) \end{aligned}$$

Theorem

$$\varphi \text{ U } \psi \equiv \neg(\neg\psi \text{ U } (\neg\varphi \wedge \neg\psi)) \wedge \text{ F } \psi$$

$$\begin{aligned} \pi \models \neg(\neg\psi \text{ U } (\neg\varphi \wedge \neg\psi)) &\iff \pi \not\models \neg\psi \text{ U } (\neg\varphi \wedge \neg\psi) \\ &\iff \text{not } \exists i \geq 1 (\pi^i \models \neg\varphi \wedge \neg\psi \text{ and } \forall j < i \pi^j \models \neg\psi) \\ &\iff \forall i \geq 1 \text{ not } (\pi^i \models \neg\varphi \wedge \neg\psi \text{ and } \forall j < i \pi^j \models \neg\psi) \\ &\iff \forall i \geq 1 (\pi^i \not\models \neg\varphi \wedge \neg\psi \text{ or not } \forall j < i \pi^j \models \neg\psi) \end{aligned}$$

Theorem

$$\varphi \mathbf{U} \psi \equiv \neg(\neg\psi \mathbf{U} (\neg\varphi \wedge \neg\psi)) \wedge \mathbf{F} \psi$$

$$\begin{aligned} \pi \models \neg(\neg\psi \mathbf{U} (\neg\varphi \wedge \neg\psi)) &\iff \pi \not\models \neg\psi \mathbf{U} (\neg\varphi \wedge \neg\psi) \\ &\iff \text{not } \exists i \geq 1 (\pi^i \models \neg\varphi \wedge \neg\psi \text{ and } \forall j < i \pi^j \models \neg\psi) \\ &\iff \forall i \geq 1 \text{ not } (\pi^i \models \neg\varphi \wedge \neg\psi \text{ and } \forall j < i \pi^j \models \neg\psi) \\ &\iff \forall i \geq 1 (\pi^i \not\models \neg\varphi \wedge \neg\psi \text{ or not } \forall j < i \pi^j \models \neg\psi) \\ &\iff \forall i \geq 1 (\pi^i \models \varphi \vee \psi \text{ or } \exists j < i \pi^j \not\models \neg\psi) \end{aligned}$$

Theorem

$$\varphi \mathbf{U} \psi \equiv \neg(\neg\psi \mathbf{U} (\neg\varphi \wedge \neg\psi)) \wedge \mathbf{F} \psi$$

$$\begin{aligned} \pi \models \neg(\neg\psi \mathbf{U} (\neg\varphi \wedge \neg\psi)) &\iff \pi \not\models \neg\psi \mathbf{U} (\neg\varphi \wedge \neg\psi) \\ &\iff \text{not } \exists i \geq 1 (\pi^i \models \neg\varphi \wedge \neg\psi \text{ and } \forall j < i \pi^j \models \neg\psi) \\ &\iff \forall i \geq 1 \text{ not } (\pi^i \models \neg\varphi \wedge \neg\psi \text{ and } \forall j < i \pi^j \models \neg\psi) \\ &\iff \forall i \geq 1 (\pi^i \not\models \neg\varphi \wedge \neg\psi \text{ or not } \forall j < i \pi^j \models \neg\psi) \\ &\iff \forall i \geq 1 (\pi^i \models \varphi \vee \psi \text{ or } \exists j < i \pi^j \not\models \neg\psi) \\ &\iff \forall i \geq 1 (\pi^i \models \varphi \vee \psi \text{ or } \exists j < i \pi^j \models \psi) \end{aligned}$$

Theorem

$$\varphi \mathbf{U} \psi \equiv \neg(\neg\psi \mathbf{U} (\neg\varphi \wedge \neg\psi)) \wedge \mathbf{F} \psi$$

$$\pi \models \neg(\neg\psi \mathbf{U} (\neg\varphi \wedge \neg\psi)) \iff \forall i \geq 1 (\pi^i \models \varphi \vee \psi \text{ or } \exists j < i \pi^j \models \psi)$$

Theorem

$$\varphi \mathbf{U} \psi \equiv \neg(\neg\psi \mathbf{U}(\neg\varphi \wedge \neg\psi)) \wedge \mathbf{F} \psi$$

$$\pi \models \neg(\neg\psi \mathbf{U}(\neg\varphi \wedge \neg\psi)) \iff \forall i \geq 1 (\pi^i \models \varphi \vee \psi \text{ or } \exists j < i \pi^j \models \psi)$$

Proof

- ▶ consider arbitrary path $\pi = s_1 \rightarrow s_2 \rightarrow s_3 \rightarrow \dots$ in arbitrary model \mathcal{M}

Theorem

$$\varphi \mathbf{U} \psi \equiv \neg(\neg\psi \mathbf{U}(\neg\varphi \wedge \neg\psi)) \wedge \mathbf{F} \psi$$

$$\pi \models \neg(\neg\psi \mathbf{U}(\neg\varphi \wedge \neg\psi)) \iff \forall i \geq 1 (\pi^i \models \varphi \vee \psi \text{ or } \exists j < i \pi^j \models \psi)$$

Proof

- ▶ consider arbitrary path $\pi = s_1 \rightarrow s_2 \rightarrow s_3 \rightarrow \dots$ in arbitrary model \mathcal{M}
- ▶ suppose $\pi \models \varphi \mathbf{U} \psi$

Theorem

$$\varphi \mathbf{U} \psi \equiv \neg(\neg\psi \mathbf{U}(\neg\varphi \wedge \neg\psi)) \wedge \mathbf{F} \psi$$

$$\pi \models \neg(\neg\psi \mathbf{U}(\neg\varphi \wedge \neg\psi)) \iff \forall i \geq 1 (\pi^i \models \varphi \vee \psi \text{ or } \exists j < i \pi^j \models \psi)$$

Proof

- ▶ consider arbitrary path $\pi = s_1 \rightarrow s_2 \rightarrow s_3 \rightarrow \dots$ in arbitrary model \mathcal{M}
- ▶ suppose $\pi \models \varphi \mathbf{U} \psi$ and consider smallest m such that $\pi^m \models \psi$

Theorem

$$\varphi \mathbf{U} \psi \equiv \neg(\neg\psi \mathbf{U}(\neg\varphi \wedge \neg\psi)) \wedge \mathbf{F} \psi$$

$$\pi \models \neg(\neg\psi \mathbf{U}(\neg\varphi \wedge \neg\psi)) \iff \forall i \geq 1 (\pi^i \models \varphi \vee \psi \text{ or } \exists j < i \pi^j \models \psi)$$

Proof

- ▶ consider arbitrary path $\pi = s_1 \rightarrow s_2 \rightarrow s_3 \rightarrow \dots$ in arbitrary model \mathcal{M}
- ▶ suppose $\pi \models \varphi \mathbf{U} \psi$ and consider smallest m such that $\pi^m \models \psi$
- ▶ $\pi \models \mathbf{F} \psi$

Theorem

$$\varphi \mathbf{U} \psi \equiv \neg(\neg\psi \mathbf{U}(\neg\varphi \wedge \neg\psi)) \wedge \mathbf{F}\psi$$

$$\pi \models \neg(\neg\psi \mathbf{U}(\neg\varphi \wedge \neg\psi)) \iff \forall i \geq 1 (\pi^i \models \varphi \vee \psi \text{ or } \exists j < i \pi^j \models \psi)$$

Proof

- ▶ consider arbitrary path $\pi = s_1 \rightarrow s_2 \rightarrow s_3 \rightarrow \dots$ in arbitrary model \mathcal{M}
- ▶ suppose $\pi \models \varphi \mathbf{U} \psi$ and consider smallest m such that $\pi^m \models \psi$
- ▶ $\pi \models \mathbf{F}\psi$ and $\pi^i \models \varphi$ for all $i < m$

Theorem

$$\varphi \mathbf{U} \psi \equiv \neg(\neg\psi \mathbf{U}(\neg\varphi \wedge \neg\psi)) \wedge \mathbf{F}\psi$$

$$\pi \models \neg(\neg\psi \mathbf{U}(\neg\varphi \wedge \neg\psi)) \iff \forall i \geq 1 (\pi^i \models \varphi \vee \psi \text{ or } \exists j < i \pi^j \models \psi)$$

Proof

- ▶ consider arbitrary path $\pi = s_1 \rightarrow s_2 \rightarrow s_3 \rightarrow \dots$ in arbitrary model \mathcal{M}
- ▶ suppose $\pi \models \varphi \mathbf{U} \psi$ and consider smallest m such that $\pi^m \models \psi$
- ▶ $\pi \models \mathbf{F}\psi$ and $\pi^i \models \varphi$ for all $i < m$
- ▶ consider $i \geq 1$

Theorem

$$\varphi \mathbf{U} \psi \equiv \neg(\neg\psi \mathbf{U}(\neg\varphi \wedge \neg\psi)) \wedge \mathbf{F}\psi$$

$$\pi \models \neg(\neg\psi \mathbf{U}(\neg\varphi \wedge \neg\psi)) \iff \forall i \geq 1 (\pi^i \models \varphi \vee \psi \text{ or } \exists j < i \pi^j \models \psi)$$

Proof

- ▶ consider arbitrary path $\pi = s_1 \rightarrow s_2 \rightarrow s_3 \rightarrow \dots$ in arbitrary model \mathcal{M}
- ▶ suppose $\pi \models \varphi \mathbf{U} \psi$ and consider smallest m such that $\pi^m \models \psi$
- ▶ $\pi \models \mathbf{F}\psi$ and $\pi^i \models \varphi$ for all $i < m$
- ▶ consider $i \geq 1$
 - ▶ if $i > m$ then $\exists j < i \pi^j \models \psi$

Theorem

$$\varphi \mathbf{U} \psi \equiv \neg(\neg\psi \mathbf{U}(\neg\varphi \wedge \neg\psi)) \wedge \mathbf{F} \psi$$

$$\pi \models \neg(\neg\psi \mathbf{U}(\neg\varphi \wedge \neg\psi)) \iff \forall i \geq 1 (\pi^i \models \varphi \vee \psi \text{ or } \exists j < i \pi^j \models \psi)$$

Proof

- ▶ consider arbitrary path $\pi = s_1 \rightarrow s_2 \rightarrow s_3 \rightarrow \dots$ in arbitrary model \mathcal{M}
- ▶ suppose $\pi \models \varphi \mathbf{U} \psi$ and consider smallest m such that $\pi^m \models \psi$
- ▶ $\pi \models \mathbf{F} \psi$ and $\pi^i \models \varphi$ for all $i < m$
- ▶ consider $i \geq 1$
 - ▶ if $i > m$ then $\exists j < i \pi^j \models \psi$ and if $i \leq m$ then $\pi^i \models \varphi \vee \psi$

Theorem

$$\varphi \mathbf{U} \psi \equiv \neg(\neg\psi \mathbf{U}(\neg\varphi \wedge \neg\psi)) \wedge \mathbf{F} \psi$$

$$\pi \models \neg(\neg\psi \mathbf{U}(\neg\varphi \wedge \neg\psi)) \iff \forall i \geq 1 (\pi^i \models \varphi \vee \psi \text{ or } \exists j < i \pi^j \models \psi)$$

Proof

- ▶ consider arbitrary path $\pi = s_1 \rightarrow s_2 \rightarrow s_3 \rightarrow \dots$ in arbitrary model \mathcal{M}
- ▶ suppose $\pi \models \varphi \mathbf{U} \psi$ and consider smallest m such that $\pi^m \models \psi$
- ▶ $\pi \models \mathbf{F} \psi$ and $\pi^i \models \varphi$ for all $i < m$
- ▶ consider $i \geq 1$
 - ▶ if $i > m$ then $\exists j < i \pi^j \models \psi$ and if $i \leq m$ then $\pi^i \models \varphi \vee \psi$
- ▶ $\pi \models \neg(\neg\psi \mathbf{U}(\neg\varphi \wedge \neg\psi))$

Theorem

$$\varphi \mathbf{U} \psi \equiv \neg(\neg\psi \mathbf{U}(\neg\varphi \wedge \neg\psi)) \wedge \mathbf{F}\psi$$

$$\pi \models \neg(\neg\psi \mathbf{U}(\neg\varphi \wedge \neg\psi)) \iff \forall i \geq 1 (\pi^i \models \varphi \vee \psi \text{ or } \exists j < i \pi^j \models \psi)$$

Proof

- ▶ consider arbitrary path $\pi = s_1 \rightarrow s_2 \rightarrow s_3 \rightarrow \dots$ in arbitrary model \mathcal{M}
- ▶ suppose $\pi \models \neg(\neg\psi \mathbf{U}(\neg\varphi \wedge \neg\psi)) \wedge \mathbf{F}\psi$

Theorem

$$\varphi \mathbf{U} \psi \equiv \neg(\neg\psi \mathbf{U}(\neg\varphi \wedge \neg\psi)) \wedge \mathbf{F}\psi$$

$$\pi \models \neg(\neg\psi \mathbf{U}(\neg\varphi \wedge \neg\psi)) \iff \forall i \geq 1 (\pi^i \models \varphi \vee \psi \text{ or } \exists j < i \pi^j \models \psi)$$

Proof

- ▶ consider arbitrary path $\pi = s_1 \rightarrow s_2 \rightarrow s_3 \rightarrow \dots$ in arbitrary model \mathcal{M}
- ▶ suppose $\pi \models \neg(\neg\psi \mathbf{U}(\neg\varphi \wedge \neg\psi)) \wedge \mathbf{F}\psi$
- ▶ there exists smallest m such that $\pi^m \models \psi$

Theorem

$$\varphi \mathbf{U} \psi \equiv \neg(\neg\psi \mathbf{U}(\neg\varphi \wedge \neg\psi)) \wedge \mathbf{F} \psi$$

$$\pi \models \neg(\neg\psi \mathbf{U}(\neg\varphi \wedge \neg\psi)) \iff \forall i \geq 1 (\pi^i \models \varphi \vee \psi \text{ or } \exists j < i \pi^j \models \psi)$$

Proof

- ▶ consider arbitrary path $\pi = s_1 \rightarrow s_2 \rightarrow s_3 \rightarrow \dots$ in arbitrary model \mathcal{M}
- ▶ suppose $\pi \models \neg(\neg\psi \mathbf{U}(\neg\varphi \wedge \neg\psi)) \wedge \mathbf{F} \psi$
- ▶ there exists **smallest m such that $\pi^m \models \psi$**
- ▶ if $i < m$ then $\pi^i \models \neg\psi$

Theorem

$$\varphi \mathbf{U} \psi \equiv \neg(\neg\psi \mathbf{U}(\neg\varphi \wedge \neg\psi)) \wedge \mathbf{F}\psi$$

$$\pi \models \neg(\neg\psi \mathbf{U}(\neg\varphi \wedge \neg\psi)) \iff \forall i \geq 1 (\pi^i \models \varphi \vee \psi \text{ or } \exists j < i \pi^j \models \psi)$$

Proof

- ▶ consider arbitrary path $\pi = s_1 \rightarrow s_2 \rightarrow s_3 \rightarrow \dots$ in arbitrary model \mathcal{M}
- ▶ suppose $\pi \models \neg(\neg\psi \mathbf{U}(\neg\varphi \wedge \neg\psi)) \wedge \mathbf{F}\psi$
- ▶ there exists smallest m such that $\pi^m \models \psi$
- ▶ if $i < m$ then $\pi^i \models \neg\psi$ and hence $\pi^i \models \varphi$

Theorem

$$\varphi \mathbf{U} \psi \equiv \neg(\neg\psi \mathbf{U}(\neg\varphi \wedge \neg\psi)) \wedge \mathbf{F}\psi$$

$$\pi \models \neg(\neg\psi \mathbf{U}(\neg\varphi \wedge \neg\psi)) \iff \forall i \geq 1 (\pi^i \models \varphi \vee \psi \text{ or } \exists j < i \pi^j \models \psi)$$

Proof

- ▶ consider arbitrary path $\pi = s_1 \rightarrow s_2 \rightarrow s_3 \rightarrow \dots$ in arbitrary model \mathcal{M}
- ▶ suppose $\pi \models \neg(\neg\psi \mathbf{U}(\neg\varphi \wedge \neg\psi)) \wedge \mathbf{F}\psi$
- ▶ there exists smallest m such that $\pi^m \models \psi$
- ▶ if $i < m$ then $\pi^i \models \neg\psi$ and hence $\pi^i \models \varphi$
- ▶ $\pi \models \varphi \mathbf{U} \psi$

Outline

1. Summary of Previous Lecture

2. Symbolic Model Checking

3. Intermezzo

4. Linear-Time Temporal Logic (LTL)

Syntax

Semantics

Example

5. Further Reading

Mutual Exclusion

- ▶ concurrent processes sharing resource

Mutual Exclusion

- ▶ concurrent processes sharing resource
- ▶ identify **critical sections** (including access to shared resource) in each process' code

Mutual Exclusion

- ▶ concurrent processes sharing resource
- ▶ identify critical sections (including access to shared resource) in each process' code
- ▶ at most one process can be in its critical section at any time

Mutual Exclusion

- ▶ concurrent processes sharing resource
- ▶ identify critical sections (including access to shared resource) in each process' code
- ▶ at most one process can be in its critical section at any time

desired:

protocol for determining which process is allowed to enter its critical section at which time

Mutual Exclusion

- ▶ concurrent processes sharing resource
- ▶ identify critical sections (including access to shared resource) in each process' code
- ▶ at most one process can be in its critical section at any time

desired:

protocol for determining which process is allowed to enter its critical section at which time

expected properties:

safety only one process is in its critical section at any time

Mutual Exclusion

- ▶ concurrent processes sharing resource
- ▶ identify critical sections (including access to shared resource) in each process' code
- ▶ at most one process can be in its critical section at any time

desired:

protocol for determining which process is allowed to enter its critical section at which time

expected properties:

safety only one process is in its critical section at any time

liveness whenever process requests to enter its critical section, it will eventually be permitted to do so

Mutual Exclusion

- ▶ concurrent processes sharing resource
- ▶ identify critical sections (including access to shared resource) in each process' code
- ▶ at most one process can be in its critical section at any time

desired:

protocol for determining which process is allowed to enter its critical section at which time

expected properties:

safety only one process is in its critical section at any time

liveness whenever process requests to enter its critical section, it will eventually be permitted to do so

non-blocking each process can always request to enter its critical section

Mutual Exclusion

- ▶ two processes have three states each:
(n) non-critical state

Mutual Exclusion

- ▶ two processes have three states each:

(n) non-critical state (t) trying to enter critical state

Mutual Exclusion

- ▶ two processes have three states each:

(*n*) non-critical state (*t*) trying to enter critical state (*c*) critical state

Mutual Exclusion

- ▶ two processes have three states each:

(n) non-critical state (t) trying to enter critical state (c) critical state


- ▶ each process undergoes transitions in cycle $n_i \rightarrow t_i \rightarrow c_i \rightarrow n_i \rightarrow \dots$ $(n_i t_i c_i)^\omega$

Mutual Exclusion

- ▶ two processes have three states each:
(*n*) non-critical state (*t*) trying to enter critical state (*c*) critical state
- ▶ each process undergoes transitions in cycle $n_i \rightarrow t_i \rightarrow c_i \rightarrow n_i \rightarrow \dots$ $(n_i t_i c_i)^\omega$
- ▶ asynchronous interleaving


Mutual Exclusion (first modeling attempt)

- ▶ two processes have three states each:
(*n*) non-critical state (*t*) trying to enter critical state (*c*) critical state
- ▶ each process undergoes transitions in cycle $n_i \rightarrow t_i \rightarrow c_i \rightarrow n_i \rightarrow \dots$ $(n_i t_i c_i)^\omega$
- ▶ asynchronous interleaving
- ▶ model (protocol):




Mutual Exclusion (first modeling attempt)

- ▶ two processes have three states each:
(*n*) non-critical state (t) trying to enter critical state (c) critical state
- ▶ each process undergoes transitions in cycle $n_i \rightarrow t_i \rightarrow c_i \rightarrow n_i \rightarrow \dots$ ($n_i t_i c_i$) $^\omega$
- ▶ asynchronous interleaving
- ▶ model (protocol):
- ▶ safety: $G \neg (c_1 \wedge c_2)$




Mutual Exclusion (first modeling attempt)

- ▶ two processes have three states each:
(*n*) non-critical state (t) trying to enter critical state (c) critical state
- ▶ each process undergoes transitions in cycle $n_i \rightarrow t_i \rightarrow c_i \rightarrow n_i \rightarrow \dots$ ($n_i t_i c_i$) $^\omega$
- ▶ asynchronous interleaving
- ▶ model (protocol):
- ▶ safety: $G \neg (c_1 \wedge c_2)$ ✓




Mutual Exclusion (first modeling attempt)

- ▶ two processes have three states each:
(*n*) non-critical state (t) trying to enter critical state (c) critical state
- ▶ each process undergoes transitions in cycle $n_i \rightarrow t_i \rightarrow c_i \rightarrow n_i \rightarrow \dots$ ($n_i t_i c_i$) $^\omega$
- ▶ asynchronous interleaving
- ▶ model (protocol):
- ▶ safety: $G \neg (c_1 \wedge c_2)$ ✓
- ▶ liveness: $G (t_1 \rightarrow F c_1)$




Mutual Exclusion (first modeling attempt)

- ▶ two processes have three states each:
(n) non-critical state (t) trying to enter critical state (c) critical state
- ▶ each process undergoes transitions in cycle $n_i \rightarrow t_i \rightarrow c_i \rightarrow n_i \rightarrow \dots$ ($n_i t_i c_i$) $^\omega$
- ▶ asynchronous interleaving
- ▶ model (protocol):
- ▶ safety: $G \neg (c_1 \wedge c_2)$ ✓
- ▶ liveness: $G (t_1 \rightarrow F c_1)$ ✗




Mutual Exclusion (first modeling attempt)

- ▶ two processes have three states each:
(*n*) non-critical state (t) trying to enter critical state (c) critical state
- ▶ each process undergoes transitions in cycle $n_i \rightarrow t_i \rightarrow c_i \rightarrow n_i \rightarrow \dots$ ($n_i t_i c_i$)^ω
- ▶ asynchronous interleaving
- ▶ model (protocol):
- ▶ safety: $G \neg (c_1 \wedge c_2)$ ✓
- ▶ liveness: $G (t_1 \rightarrow F c_1)$ ✗
- ▶ non-blocking holds but is not expressible in LTL




Mutual Exclusion (first modeling attempt)

- ▶ two processes have three states each:
(n) non-critical state (t) trying to enter critical state (c) critical state
- ▶ each process undergoes transitions in cycle $n_i \rightarrow t_i \rightarrow c_i \rightarrow n_i \rightarrow \dots$ ($n_i t_i c_i$) $^\omega$
- ▶ asynchronous interleaving
- ▶ model (protocol):
- ▶ safety: $G \neg (c_1 \wedge c_2)$ ✓
- ▶ liveness: $G (t_1 \rightarrow F c_1)$ ✗
- ▶ non-blocking holds but is not expressible in LTL
 $AG(n_1 \rightarrow EX t_1)$ in CTL




Mutual Exclusion (second modeling attempt)

- ▶ safety: $G \neg (c_1 \wedge c_2)$
- ▶ liveness: $G (t_1 \rightarrow F c_1)$




Mutual Exclusion (second modeling attempt)

- ▶ safety: $G \neg (c_1 \wedge c_2)$ ✓
- ▶ liveness: $G (t_1 \rightarrow F c_1)$ ✓



Mutual Exclusion (second modeling attempt)

- ▶ safety: $G \neg (c_1 \wedge c_2)$ ✓
- ▶ liveness: $G (t_1 \rightarrow F c_1)$ ✓



NuSMV (New Symbolic Model Verifier)

provides language for describing models and checks satisfaction of LTL and CTL formulas

Mutual Exclusion Protocol in NuSMV

```
MODULE main
VAR
  pr1 : process prc ( pr2.st, turn, FALSE ) ;
  pr2 : process prc ( pr1.st, turn, TRUE ) ;
  turn : boolean ;
ASSIGN
  init ( turn ) := FALSE ;

LTLSPEC G ! (( pr1.st = c ) & ( pr2.st = c )) -- safety
LTLSPEC G (( pr1.st = t ) -> F ( pr1.st = c )) -- liveness
LTLSPEC G (( pr2.st = t ) -> F ( pr2.st = c )) -- liveness

MODULE prc ( other-st, turn, myturn )
VAR st : { n, t, c } ;
ASSIGN
  init ( st ) := n ;
  next ( st ) := case
    ( st = n )           : { n, t } ;
    ( st = t ) & ( other-st = n ) : c ;
    ( st = t ) & ( other-st = t ) : c ;
    ( st = c )           : st ;
  TRUE                   : st ;
  esac ;
  next ( turn ) := case
    turn = myturn & st = c      : ! turn ;
    TRUE                         : turn ;
  esac ;

FAIRNESS running
FAIRNESS ! ( st = c )
```

Outline

1. Summary of Previous Lecture
2. Symbolic Model Checking
3. Intermezzo
4. Linear-Time Temporal Logic (LTL)
- 5. Further Reading**

Huth and Ryan

- ▶ Section 3.1
- ▶ Section 3.2
- ▶ Section 3.3
- ▶ Section 3.7
- ▶ Section 6.3

Huth and Ryan

- ▶ Section 3.1
- ▶ Section 3.2
- ▶ Section 3.3
- ▶ Section 3.7
- ▶ Section 6.3

Model Checking Tools

- ▶ NuSMV
- ▶ Spin

Important Concepts

- ▶ $[[\varphi]]$
- ▶ F
- ▶ fixed point
- ▶ G
- ▶ greatest fixed point
- ▶ Knaster–Tarski
- ▶ least fixed point
- ▶ linear-time temporal logic
- ▶ liveness
- ▶ LTL
- ▶ non-blocking
- ▶ path
- ▶ pre_{\forall}
- ▶ pre_{\exists}
- ▶ R
- ▶ safety
- ▶ symbolic model checking
- ▶ U
- ▶ W
- ▶ X

Important Concepts

- ▶ $[[\varphi]]$
- ▶ F
- ▶ fixed point
- ▶ G
- ▶ greatest fixed point
- ▶ Knaster–Tarski
- ▶ least fixed point
- ▶ linear-time temporal logic
- ▶ liveness
- ▶ LTL
- ▶ non-blocking
- ▶ path
- ▶ pre_{\forall}
- ▶ pre_{\exists}
- ▶ R
- ▶ safety
- ▶ symbolic model checking
- ▶ U
- ▶ W
- ▶ X

homework for June 6

Important Concepts

- ▶ $[[\varphi]]$
- ▶ F
- ▶ fixed point
- ▶ G
- ▶ greatest fixed point
- ▶ Knaster–Tarski
- ▶ least fixed point
- ▶ linear-time temporal logic
- ▶ liveness
- ▶ LTL
- ▶ non-blocking
- ▶ path
- ▶ pre_{\forall}
- ▶ pre_{\exists}
- ▶ R
- ▶ safety
- ▶ symbolic model checking
- ▶ U
- ▶ W
- ▶ X

homework for June 6

next week (June 3): online evaluation in presence

Important Concepts

- ▶ $[[\varphi]]$
- ▶ F
- ▶ fixed point
- ▶ G
- ▶ greatest fixed point
- ▶ Knaster–Tarski
- ▶ least fixed point
- ▶ linear-time temporal logic
- ▶ liveness
- ▶ LTL
- ▶ non-blocking
- ▶ path
- ▶ pre_{\forall}
- ▶ pre_{\exists}
- ▶ R
- ▶ safety
- ▶ symbolic model checking
- ▶ U
- ▶ W
- ▶ X

homework for June 6

next week (June 3): online evaluation in presence \implies bring device