



# Computability Theory

Aart Middeldorp

# Outline

- 1. Summary of Previous Lecture**
- 2. Strong Normalization**
- 3. Type Inference**
- 4. Type Inhabitation**
- 5. Intuitionistic Propositional Logic**
- 6. Summary**
- 7. Test**

## Definitions

- Gödel number of CL-term is defined inductively:

$$g(I) = \langle 0 \rangle \quad g(K) = \langle 1 \rangle \quad g(S) = \langle 2 \rangle \quad g(t u) = \langle 3, g(t), g(u) \rangle \quad g(x_i) = \langle 4, i \rangle$$

- $\text{enc}(n) = g(n)$
- sets  $T$  and  $U$  of CL-terms are recursively separable if  $\{g(t) \mid t \in T\}$  and  $\{g(u) \mid u \in U\}$  are recursively separable
- set  $T$  of CL-terms is conversion-closed if  $u \in T$  whenever  $t \in T$  and  $t \leftrightarrow^* u$

## Theorem

CL-representable functions are partial recursive

## Notation

$\ulcorner t \urcorner = \underline{g(t)}$  is Church numeral of Gödel number of CL-term  $t$

## Theorem

$\forall \text{ CL-term } F \ \exists \text{ CL-term } X \text{ such that } F \sqcap X \sqcap \leftrightarrow^* X$

## Theorem

- ▶ non-empty conversion-closed sets of CL-terms are recursively inseparable
- ▶ non-trivial conversion-closed sets of CL-terms are not recursive

## Corollary

set of normalizing CL-terms is not recursive: decision problem

instance: CL-term  $t$

question: is  $t$  normalizing ?

is undecidable

## Definition (Types)

set  $\mathbb{T}$  of **types** is defined inductively:

- ▶  $V \subseteq \mathbb{T}$  infinite set of **type variables**
- ▶  $C \subseteq \mathbb{T}$  **type constants**
- ▶ if  $\sigma, \tau \in \mathbb{T}$  then  $(\sigma \rightarrow \tau) \in \mathbb{T}$

## Definition (Type Assignment, Curry-style)

- ▶ type assignment formula  $t : \tau$  with CL-term  $t$  and type  $\tau$
- ▶ type assignment system **TA**

$$\overline{I : \sigma \rightarrow \sigma} \quad \overline{K : \sigma \rightarrow \tau \rightarrow \sigma} \quad \overline{S : (\rho \rightarrow \sigma \rightarrow \tau) \rightarrow (\rho \rightarrow \sigma) \rightarrow \rho \rightarrow \tau}$$

$$\frac{t : \sigma \rightarrow \tau \quad u : \sigma}{tu : \tau}$$

for all types  $\sigma, \tau, \rho$  and CL-terms  $s$  and  $t$

## Notation

$\Gamma \vdash t : \tau$  if  $t : \tau$  can be derived in TA from assumptions in  $\Gamma$

## Theorem

if  $\Gamma, x : \sigma \vdash t : \tau$  and  $x \notin \text{Var}(\Gamma)$  then  $\Gamma \vdash [x]t : \sigma \rightarrow \tau$

## Theorem (Subject Reduction)

if  $\Gamma \vdash t : \tau$  and  $t \rightarrow u$  then  $\Gamma \vdash u : \tau$

## Definition

CL-term  $t$  with  $\text{Var}(t) = \{x_1, \dots, x_n\}$  is **typable** if

$$x_1 : \rho_1, \dots, x_n : \rho_n \vdash t : \tau$$

for some types  $\rho_1, \dots, \rho_n, \tau$

## Part I: Recursive Function Theory

Ackermann function, bounded minimization, bounded recursion, course-of-values recursion, diagonalization, diophantine sets, elementary functions, fixed point theorem, Fibonacci numbers, Gödel numbering, Gödel's  $\beta$  function, Grzegorczyk hierarchy, loop programs, minimization, normal form theorem, partial recursive functions, primitive recursion, recursive enumerability, recursive inseparability, s-m-n theorem, total recursive functions, undecidability, while programs, ...

## Part II: Combinatory Logic and Lambda Calculus

$\alpha$ -equivalence, abstraction, arithmetization,  $\beta$ -reduction, CL-representability, combinators, combinatorial completeness, Church numerals, Church–Rosser theorem, Curry–Howard isomorphism, de Bruijn notation,  $\eta$ -reduction, fixed point theorem, intuitionistic propositional logic,  $\lambda$ -definability, normalization theorem, termination, typing, undecidability, Z property, ...

## Part I: Recursive Function Theory

Ackermann function, bounded minimization, bounded recursion, course-of-values recursion, diagonalization, diophantine sets, elementary functions, fixed point theorem, Fibonacci numbers, Gödel numbering, Gödel's  $\beta$  function, Grzegorczyk hierarchy, loop programs, minimization, normal form theorem, partial recursive functions, primitive recursion, recursive enumerability, recursive inseparability, s-m-n theorem, total recursive functions, undecidability, while programs, ...

## Part II: Combinatory Logic and Lambda Calculus

$\alpha$ -equivalence, abstraction, arithmetization,  $\beta$ -reduction, CL-representability, combinators, combinatorial completeness, Church numerals, Church–Rosser theorem, Curry–Howard isomorphism, de Bruijn notation,  $\eta$ -reduction, fixed point theorem, **intuitionistic propositional logic**,  $\lambda$ -definability, normalization theorem, **termination, typing**, undecidability, Z property, ...

# Outline

1. Summary of Previous Lecture
2. **Strong Normalization**
3. Type Inference
4. Type Inhabitation
5. Intuitionistic Propositional Logic
6. Summary
7. Test

## Theorem (Strong Normalization)

typable CL-terms are terminating

## Theorem (Strong Normalization)

typable CL-terms are terminating (**SN**)

## Theorem (Strong Normalization)

typable CL-terms are terminating (SN)

### Definition

typable CL-term  $t$  is **strongly computable** (SC) if

- ▶  $t$  has atomic type  $\tau \in \mathbb{V} \cup \mathbb{C}$  and is SN

## Theorem (Strong Normalization)

typable CL-terms are terminating (SN)

### Definition

typable CL-term  $t$  is **strongly computable** (SC) if

- ▶  $t$  has atomic type  $\tau \in \mathbb{V} \cup \mathbb{C}$  and is SN
- ▶  $t$  has type  $\sigma \rightarrow \tau$  and  $tu$  is SC whenever  $u : \sigma$  is SC

## Lemma ①

for any type  $\tau$

- ① every term  $x u_1 \dots u_n$  of type  $\tau$  with variable  $x$  and SN terms  $u_1, \dots, u_n$  is SC

## Lemma ①

for any type  $\tau$

- ① every term  $x u_1 \dots u_n$  of type  $\tau$  with variable  $x$  and SN terms  $u_1, \dots, u_n$  is SC
- ② every SC term of type  $\tau$  is SN

## Lemma ①

for any type  $\tau$

- ① every term  $x u_1 \dots u_n$  of type  $\tau$  with variable  $x$  and SN terms  $u_1, \dots, u_n$  is SC
- ② every SC term of type  $\tau$  is SN

## Proof

induction on  $\tau$

## Lemma ①

for any type  $\tau$

- ① every term  $x u_1 \dots u_n$  of type  $\tau$  with variable  $x$  and SN terms  $u_1, \dots, u_n$  is SC
- ② every SC term of type  $\tau$  is SN

## Proof

induction on  $\tau$

- ▶ base case:  $\tau$  is atomic

## Lemma ①

for any type  $\tau$

- ① every term  $x u_1 \cdots u_n$  of type  $\tau$  with variable  $x$  and **SN terms  $u_1, \dots, u_n$**  is SC
- ② every SC term of type  $\tau$  is SN

## Proof

induction on  $\tau$

- ▶ base case:  $\tau$  is atomic
  - ①  $t = x u_1 \cdots u_n$  is SN

## Lemma ①

for any type  $\tau$

- ① every term  $x u_1 \cdots u_n$  of type  $\tau$  with variable  $x$  and SN terms  $u_1, \dots, u_n$  is SC
- ② every SC term of type  $\tau$  is SN

## Proof

induction on  $\tau$

► base case:  $\tau$  is atomic

$$\textcircled{1} \quad t = x u_1 \cdots u_n \text{ is SN} \implies t \text{ is SC}$$

## Lemma ①

for any type  $\tau$

- ① every term  $x u_1 \cdots u_n$  of type  $\tau$  with variable  $x$  and SN terms  $u_1, \dots, u_n$  is SC
- ② every SC term of type  $\tau$  is SN

## Proof

induction on  $\tau$

- base case:  $\tau$  is atomic
  - ①  $t = x u_1 \cdots u_n$  is SN  $\implies t$  is SC
  - ②  $t : \tau$  is SC

## Lemma ①

for any type  $\tau$

- ① every term  $x u_1 \cdots u_n$  of type  $\tau$  with variable  $x$  and SN terms  $u_1, \dots, u_n$  is SC
- ② every SC term of type  $\tau$  is SN

## Proof

induction on  $\tau$

► base case:  $\tau$  is atomic

- ①  $t = x u_1 \cdots u_n$  is SN  $\implies t$  is SC
- ②  $t : \tau$  is SC  $\implies t$  is SN

## Lemma ①

for any type  $\tau$

- ① every term  $x u_1 \dots u_n$  of type  $\tau$  with variable  $x$  and SN terms  $u_1, \dots, u_n$  is SC
- ② every SC term of type  $\tau$  is SN

## Proof

induction on  $\tau$

- ▶ base case:  $\tau$  is atomic
  - ①  $t = x u_1 \dots u_n$  is SN  $\implies t$  is SC
  - ②  $t : \tau$  is SC  $\implies t$  is SN
- ▶ step case:  $\tau = \rho \rightarrow \sigma$

## Lemma ①

for any type  $\tau$

- ① every term  $x u_1 \cdots u_n$  of type  $\tau$  with variable  $x$  and SN terms  $u_1, \dots, u_n$  is SC
- ② every SC term of type  $\tau$  is SN

## Proof

induction on  $\tau$

- ▶ base case:  $\tau$  is atomic
  - ①  $t = x u_1 \cdots u_n$  is SN  $\implies t$  is SC
  - ②  $t : \tau$  is SC  $\implies t$  is SN
- ▶ step case:  $\tau = \rho \rightarrow \sigma$ 
  - ①  $v : \rho$  is SC

## Lemma ①

for any type  $\tau$

- ① every term  $x u_1 \dots u_n$  of type  $\tau$  with variable  $x$  and SN terms  $u_1, \dots, u_n$  is SC
- ② every SC term of type  $\tau$  is SN

## Proof

induction on  $\tau$

- ▶ base case:  $\tau$  is atomic
  - ①  $t = x u_1 \dots u_n$  is SN  $\implies t$  is SC
  - ②  $t : \tau$  is SC  $\implies t$  is SN
- ▶ step case:  $\tau = \rho \rightarrow \sigma$ 
  - ①  $v : \rho$  is SC  $\implies v$  is SN

## Lemma ①

for any type  $\tau$

- ① every term  $xu_1 \dots u_n$  of type  $\tau$  with variable  $x$  and SN terms  $u_1, \dots, u_n$  is SC
- ② every SC term of type  $\tau$  is SN

## Proof

induction on  $\tau$

- ▶ base case:  $\tau$  is atomic
  - ①  $t = xu_1 \dots u_n$  is SN  $\implies t$  is SC
  - ②  $t : \tau$  is SC  $\implies t$  is SN
- ▶ step case:  $\tau = \rho \rightarrow \sigma$ 
  - ①  $v : \rho$  is SC  $\implies v$  is SN  $\implies xu_1 \dots u_n v : \sigma$  is SC

## Lemma ①

for any type  $\tau$

- ① every term  $x u_1 \dots u_n$  of type  $\tau$  with variable  $x$  and SN terms  $u_1, \dots, u_n$  is SC
- ② every SC term of type  $\tau$  is SN

## Proof

induction on  $\tau$

- ▶ base case:  $\tau$  is atomic
  - ①  $t = x u_1 \dots u_n$  is SN  $\implies t$  is SC
  - ②  $t : \tau$  is SC  $\implies t$  is SN
- ▶ step case:  $\tau = \rho \rightarrow \sigma$ 
  - ①  $v : \rho$  is SC  $\implies v$  is SN  $\implies \textcolor{red}{x u_1 \dots u_n v : \sigma}$  is SC  $\implies x u_1 \dots u_n : \tau$  is SC

## Lemma ①

for any type  $\tau$

- ① every term  $xu_1 \dots u_n$  of type  $\tau$  with variable  $x$  and SN terms  $u_1, \dots, u_n$  is SC
- ② every SC term of type  $\tau$  is SN

## Proof

induction on  $\tau$

- ▶ base case:  $\tau$  is atomic
  - ①  $t = xu_1 \dots u_n$  is SN  $\implies t$  is SC
  - ②  $t : \tau$  is SC  $\implies t$  is SN
- ▶ step case:  $\tau = \rho \rightarrow \sigma$ 
  - ①  $v : \rho$  is SC  $\implies v$  is SN  $\implies xu_1 \dots u_n v : \sigma$  is SC  $\implies xu_1 \dots u_n : \tau$  is SC
  - ② consider SC term  $t : \tau$

## Lemma ①

for any type  $\tau$

- ① every term  $xu_1 \dots u_n$  of type  $\tau$  with variable  $x$  and SN terms  $u_1, \dots, u_n$  is SC
- ② every SC term of type  $\tau$  is SN

## Proof

induction on  $\tau$

- ▶ base case:  $\tau$  is atomic
  - ①  $t = xu_1 \dots u_n$  is SN  $\implies t$  is SC
  - ②  $t : \tau$  is SC  $\implies t$  is SN
- ▶ step case:  $\tau = \rho \rightarrow \sigma$ 
  - ①  $v : \rho$  is SC  $\implies v$  is SN  $\implies xu_1 \dots u_n v : \sigma$  is SC  $\implies xu_1 \dots u_n : \tau$  is SC
  - ② consider SC term  $t : \tau$  and  $x : \rho$  with  $x \in \mathcal{V} \setminus \text{Var}(t)$

## Lemma ①

for any type  $\tau$

- ① every term  $x u_1 \cdots u_n$  of type  $\tau$  with variable  $x$  and SN terms  $u_1, \dots, u_n$  is SC
- ② every SC term of type  $\tau$  is SN

## Proof

induction on  $\tau$

- ▶ base case:  $\tau$  is atomic
  - ①  $t = x u_1 \cdots u_n$  is SN  $\implies t$  is SC
  - ②  $t : \tau$  is SC  $\implies t$  is SN
- ▶ step case:  $\tau = \rho \rightarrow \sigma$ 
  - ①  $v : \rho$  is SC  $\implies v$  is SN  $\implies x u_1 \cdots u_n v : \sigma$  is SC  $\implies x u_1 \cdots u_n : \tau$  is SC
  - ② consider SC term  $t : \tau$  and  $x : \rho$  with  $x \in \mathcal{V} \setminus \text{Var}(t)$   
 $\implies x$  is SC

## Lemma ①

for any type  $\tau$

- ① every term  $xu_1 \dots u_n$  of type  $\tau$  with variable  $x$  and SN terms  $u_1, \dots, u_n$  is SC
- ② every SC term of type  $\tau$  is SN

## Proof

induction on  $\tau$

- ▶ base case:  $\tau$  is atomic
  - ①  $t = xu_1 \dots u_n$  is SN  $\implies t$  is SC
  - ②  $t : \tau$  is SC  $\implies t$  is SN
- ▶ step case:  $\tau = \rho \rightarrow \sigma$ 
  - ①  $v : \rho$  is SC  $\implies v$  is SN  $\implies xu_1 \dots u_n v : \sigma$  is SC  $\implies xu_1 \dots u_n : \tau$  is SC
  - ② consider SC term  $t : \tau$  and  $x : \rho$  with  $x \in V \setminus \text{Var}(t)$   
 $\implies x$  is SC  $\implies tx$  is SC

## Lemma ①

for any type  $\tau$

- ① every term  $xu_1 \dots u_n$  of type  $\tau$  with variable  $x$  and SN terms  $u_1, \dots, u_n$  is SC
- ② every SC term of type  $\tau$  is SN

## Proof

induction on  $\tau$

- ▶ base case:  $\tau$  is atomic
  - ①  $t = xu_1 \dots u_n$  is SN  $\implies t$  is SC
  - ②  $t : \tau$  is SC  $\implies t$  is SN
- ▶ step case:  $\tau = \rho \rightarrow \sigma$ 
  - ①  $v : \rho$  is SC  $\implies v$  is SN  $\implies xu_1 \dots u_n v : \sigma$  is SC  $\implies xu_1 \dots u_n : \tau$  is SC
  - ② consider SC term  $t : \tau$  and  $x : \rho$  with  $x \in \mathcal{V} \setminus \text{Var}(t)$   
 $\implies x$  is SC  $\implies tx$  is SC  $\implies tx$  is SN

## Lemma ①

for any type  $\tau$

- ① every term  $xu_1 \dots u_n$  of type  $\tau$  with variable  $x$  and SN terms  $u_1, \dots, u_n$  is SC
- ② every SC term of type  $\tau$  is SN

## Proof

induction on  $\tau$

- ▶ base case:  $\tau$  is atomic
  - ①  $t = xu_1 \dots u_n$  is SN  $\implies t$  is SC
  - ②  $t : \tau$  is SC  $\implies t$  is SN
- ▶ step case:  $\tau = \rho \rightarrow \sigma$ 
  - ①  $v : \rho$  is SC  $\implies v$  is SN  $\implies xu_1 \dots u_n v : \sigma$  is SC  $\implies xu_1 \dots u_n : \tau$  is SC
  - ② consider SC term  $t : \tau$  and  $x : \rho$  with  $x \in \mathcal{V} \setminus \text{Var}(t)$   
 $\implies x$  is SC  $\implies tx$  is SC  $\implies \text{tx is SN}$   $\implies t$  is SN

## Lemma ②

S, K and I are SC

## Lemma ②

S, K and I are SC

### Proof

K :  $\sigma \rightarrow \tau \rightarrow \sigma$

## Lemma ②

S, K and I are SC

### Proof

K :  $\sigma \rightarrow \tau \rightarrow \sigma$

►  $\sigma = \sigma_1 \rightarrow \dots \rightarrow \sigma_n \rightarrow \theta$  with atomic type  $\theta$  and  $n \geq 0$

## Lemma ②

S, K and I are SC

### Proof

K :  $\sigma \rightarrow \tau \rightarrow \sigma$

- $\sigma = \sigma_1 \rightarrow \dots \rightarrow \sigma_n \rightarrow \theta$  with atomic type  $\theta$  and  $n \geq 0$
- consider arbitrary SC terms  $s : \sigma$   $t : \tau$   $u_1 : \sigma_1$   $\dots$   $u_n : \sigma_n$

## Lemma ②

S, K and I are SC

### Proof

K :  $\sigma \rightarrow \tau \rightarrow \sigma$

- ▶  $\sigma = \sigma_1 \rightarrow \dots \rightarrow \sigma_n \rightarrow \theta$  with atomic type  $\theta$  and  $n \geq 0$
- ▶ consider arbitrary SC terms  $s : \sigma$     $t : \tau$     $u_1 : \sigma_1$     $\dots$     $u_n : \sigma_n$
- ▶  $s$  is SC    $\implies$     $s u_1 \dots u_n$  is SC

## Lemma ②

S, K and I are SC

### Proof

K :  $\sigma \rightarrow \tau \rightarrow \sigma$

- ▶  $\sigma = \sigma_1 \rightarrow \dots \rightarrow \sigma_n \rightarrow \theta$  with atomic type  $\theta$  and  $n \geq 0$
- ▶ consider arbitrary SC terms  $s : \sigma$     $t : \tau$     $u_1 : \sigma_1$    ...    $u_n : \sigma_n$
- ▶  $s$  is SC    $\Rightarrow$     $s u_1 \dots u_n$  is SC    $\Rightarrow$     $s u_1 \dots u_n$  is SN

## Lemma ②

S, K and I are SC

### Proof

K :  $\sigma \rightarrow \tau \rightarrow \sigma$

- ▶  $\sigma = \sigma_1 \rightarrow \dots \rightarrow \sigma_n \rightarrow \theta$  with atomic type  $\theta$  and  $n \geq 0$
- ▶ consider arbitrary SC terms  $s : \sigma$     $t : \tau$     $u_1 : \sigma_1$     $\dots$     $u_n : \sigma_n$
- ▶  $s$  is SC    $\implies$     $s u_1 \dots u_n$  is SC    $\implies$     $s u_1 \dots u_n$  is SN
- ▶  $t$  is SC    $\implies$     $t$  is SN

## Lemma ②

S, K and I are SC

### Proof

K :  $\sigma \rightarrow \tau \rightarrow \sigma$

- ▶  $\sigma = \sigma_1 \rightarrow \dots \rightarrow \sigma_n \rightarrow \theta$  with atomic type  $\theta$  and  $n \geq 0$
- ▶ consider arbitrary SC terms  $s : \sigma$   $t : \tau$   $u_1 : \sigma_1 \dots u_n : \sigma_n$
- ▶  $s$  is SC  $\implies su_1 \dots u_n$  is SC  $\implies su_1 \dots u_n$  is SN
- ▶  $t$  is SC  $\implies t$  is SN
- ▶  $Kstu_1 \dots u_n$  is SN

## Lemma ②

S, K and I are SC

### Proof

K :  $\sigma \rightarrow \tau \rightarrow \sigma$

- ▶  $\sigma = \sigma_1 \rightarrow \dots \rightarrow \sigma_n \rightarrow \theta$  with atomic type  $\theta$  and  $n \geq 0$
- ▶ consider arbitrary SC terms  $s : \sigma$   $t : \tau$   $u_1 : \sigma_1 \dots u_n : \sigma_n$
- ▶  $s$  is SC  $\implies su_1 \dots u_n$  is SC  $\implies su_1 \dots u_n$  is SN
- ▶  $t$  is SC  $\implies t$  is SN
- ▶  $Kstu_1 \dots u_n$  is SN: any infinite reduction starts with

$$Kstu_1 \dots u_n \xrightarrow{*} Ks't'u'_1 \dots u'_n$$

with  $s \xrightarrow{*} s'$   $t \xrightarrow{*} t'$   $u_1 \xrightarrow{*} u'_1$   $u_n \xrightarrow{*} u'_n$

## Lemma ②

S, K and I are SC

### Proof

K :  $\sigma \rightarrow \tau \rightarrow \sigma$

- $\sigma = \sigma_1 \rightarrow \dots \rightarrow \sigma_n \rightarrow \theta$  with atomic type  $\theta$  and  $n \geq 0$
- consider arbitrary SC terms  $s : \sigma$   $t : \tau$   $u_1 : \sigma_1 \dots u_n : \sigma_n$
- $s$  is SC  $\implies su_1 \dots u_n$  is SC  $\implies su_1 \dots u_n$  is SN
- $t$  is SC  $\implies t$  is SN
- $Kstu_1 \dots u_n$  is SN: any infinite reduction starts with

$$Kstu_1 \dots u_n \xrightarrow{*} Ks't'u'_1 \dots u'_n \xrightarrow{*} s'u'_1 \dots u'_n \xrightarrow{*} \dots$$

with  $s \xrightarrow{*} s'$   $t \xrightarrow{*} t'$   $u_1 \xrightarrow{*} u'_1$   $u_n \xrightarrow{*} u'_n$

## Lemma ②

S, K and I are SC

### Proof

K :  $\sigma \rightarrow \tau \rightarrow \sigma$

- $\sigma = \sigma_1 \rightarrow \dots \rightarrow \sigma_n \rightarrow \theta$  with atomic type  $\theta$  and  $n \geq 0$
- consider arbitrary SC terms  $s : \sigma$   $t : \tau$   $u_1 : \sigma_1 \dots u_n : \sigma_n$
- $s$  is SC  $\implies su_1 \dots u_n$  is SC  $\implies su_1 \dots u_n$  is SN
- $t$  is SC  $\implies t$  is SN
- Kstu<sub>1</sub>...u<sub>n</sub> is SN: any infinite reduction starts with

$$\begin{aligned} \text{Kstu}_1 \dots u_n &\rightarrow^* \text{K}s' t' u'_1 \dots u'_n \rightarrow s' u'_1 \dots u'_n \rightarrow^* \dots \\ &\implies su_1 \dots u_n \rightarrow s' u'_1 \dots u'_n \rightarrow^* \dots \end{aligned}$$

with  $s \rightarrow^* s'$   $t \rightarrow^* t'$   $u_1 \rightarrow^* u'_1$   $u_n \rightarrow^* u'_n$

## Lemma ②

S, K and I are SC

### Proof

K :  $\sigma \rightarrow \tau \rightarrow \sigma$

- $\sigma = \sigma_1 \rightarrow \dots \rightarrow \sigma_n \rightarrow \theta$  with atomic type  $\theta$  and  $n \geq 0$
- consider arbitrary SC terms  $s : \sigma$   $t : \tau$   $u_1 : \sigma_1 \dots u_n : \sigma_n$
- $s$  is SC  $\implies su_1 \dots u_n$  is SC  $\implies su_1 \dots u_n$  is SN
- $t$  is SC  $\implies t$  is SN
- $Kstu_1 \dots u_n$  is SN: any infinite reduction starts with

$$\begin{aligned} Kstu_1 \dots u_n &\rightarrow^* Ks't'u'_1 \dots u'_n \rightarrow s'u'_1 \dots u'_n \rightarrow^* \dots \\ &\implies su_1 \dots u_n \rightarrow s'u'_1 \dots u'_n \rightarrow^* \dots \end{aligned}$$

with  $s \rightarrow^* s'$   $t \rightarrow^* t'$   $u_1 \rightarrow^* u'_1$   $u_n \rightarrow^* u'_n$

## Lemma ②

S, K and I are SC

### Proof

K :  $\sigma \rightarrow \tau \rightarrow \sigma$

- $\sigma = \sigma_1 \rightarrow \dots \rightarrow \sigma_n \rightarrow \theta$  with atomic type  $\theta$  and  $n \geq 0$
- consider arbitrary SC terms  $s : \sigma$   $t : \tau$   $u_1 : \sigma_1 \dots u_n : \sigma_n$
- $s$  is SC  $\implies su_1 \dots u_n$  is SC  $\implies su_1 \dots u_n$  is SN
- $t$  is SC  $\implies t$  is SN
- Kstu<sub>1</sub>...u<sub>n</sub> is SN: any infinite reduction starts with

$$\begin{aligned} \text{Kstu}_1 \dots u_n &\rightarrow^* \text{Ks}'t'u'_1 \dots u'_n \rightarrow s'u'_1 \dots u'_n \rightarrow^* \dots \\ &\implies su_1 \dots u_n \rightarrow s'u'_1 \dots u'_n \rightarrow^* \dots \end{aligned}$$

with  $s \rightarrow^* s'$   $t \rightarrow^* t'$   $u_1 \rightarrow^* u'_1$   $u_n \rightarrow^* u'_n$

S and I: homework exercise

## Lemma

every typable CL-term is SC

## Lemma

every typable CL-term is SC

## Proof

induction on term  $t$

## Lemma

every typable CL-term is SC

## Proof

induction on term  $t$

- ▶  $t$  is variable

## Lemma

every typable CL-term is SC

## Proof

induction on term  $t$

- $t$  is variable  $\implies t$  is SC by lemma 1

## Lemma

every typable CL-term is SC

## Proof

induction on term  $t$

- ▶  $t$  is variable  $\implies t$  is SC by lemma 1
- ▶  $t \in \{I, K, S\}$

## Lemma

every typable CL-term is SC

## Proof

induction on term  $t$

- ▶  $t$  is variable  $\implies t$  is SC by lemma ①
- ▶  $t \in \{I, K, S\}$   $\implies t$  is SC by lemma ②

## Lemma

every typable CL-term is SC

## Proof

induction on term  $t$

- ▶  $t$  is variable  $\implies t$  is SC by lemma ①
- ▶  $t \in \{I, K, S\}$   $\implies t$  is SC by lemma ②
- ▶  $t = t_1 t_2$

## Lemma

every typable CL-term is SC

## Proof

induction on term  $t$

- ▶  $t$  is variable  $\implies t$  is SC by lemma ①
- ▶  $t \in \{I, K, S\}$   $\implies t$  is SC by lemma ②
- ▶  $t = t_1 t_2$   $\implies t_1$  and  $t_2$  are SC by induction hypothesis

## Lemma

every typable CL-term is SC

## Proof

induction on term  $t$

- ▶  $t$  is variable  $\implies t$  is SC by lemma ①
- ▶  $t \in \{I, K, S\}$   $\implies t$  is SC by lemma ②
- ▶  $t = t_1 t_2$   $\implies t_1$  and  $t_2$  are SC by induction hypothesis  $\implies t$  is SC

## Lemma

every typable CL-term is SC

## Proof

induction on term  $t$

- ▶  $t$  is variable  $\implies t$  is SC by lemma ①
- ▶  $t \in \{I, K, S\}$   $\implies t$  is SC by lemma ②
- ▶  $t = t_1 t_2$   $\implies t_1$  and  $t_2$  are SC by induction hypothesis  $\implies t$  is SC

## Corollary

typable CL-terms are SN

# Outline

1. Summary of Previous Lecture
2. Strong Normalization
- 3. Type Inference**
4. Type Inhabitation
5. Intuitionistic Propositional Logic
6. Summary
7. Test

## Theorem

problem

instance: CL-term  $t$

question: is  $t$  typable ?

is decidable

## Theorem

problem

instance: CL-term  $t$

question: is  $t$  typable?

is decidable

## Definition

principal type of combinator  $t$  is any type  $\sigma$  such that

①  $\vdash t : \sigma$

## Theorem

problem

instance: CL-term  $t$

question: is  $t$  typable?

is decidable

## Definition

principal type of combinator  $t$  is any type  $\sigma$  such that

- ①  $\vdash t : \sigma$
- ② if  $\vdash t : \tau$  then  $\tau$  is substitution instance of  $\sigma$

## Theorem

problem

instance: CL-term  $t$

question: is  $t$  typable?

is decidable

## Definition

principal type of combinator  $t$  is any type  $\sigma$  such that

- ①  $\vdash t : \sigma$
- ② if  $\vdash t : \tau$  then  $\tau$  is substitution instance of  $\sigma$

## Example

SKK has principal type  $a \rightarrow a$  (where  $a$  is type variable)

## Theorem

every typable combinator has principal type

## Theorem

every typable combinator has principal type

## Type Inference

principal types can be computed by typing rules of TA (with type variables  $\sigma, \tau, \rho$ )

$$\text{I} : \sigma \rightarrow \sigma \quad \text{K} : \sigma \rightarrow \tau \rightarrow \sigma \quad \text{S} : (\rho \rightarrow \sigma \rightarrow \tau) \rightarrow (\rho \rightarrow \sigma) \rightarrow \rho \rightarrow \tau$$

$$\frac{t : \sigma \rightarrow \tau \quad u : \sigma}{tu : \tau}$$

and unification algorithm

## Example ①

principle type of SKK

## Example ①

principle type of  $\text{SKK} : \beta$

- ▶  $\text{SK} : \alpha \rightarrow \beta$      $\text{K} : \alpha$

## Example ①

principle type of  $\text{SKK} : \beta$

- ▶  $\text{SK} : \alpha \rightarrow \beta$     $\text{K} : \alpha$
- ▶ unification problem

$$\alpha \approx \sigma_1 \rightarrow \tau_1 \rightarrow \sigma_1$$

## Example ①

principle type of  $\text{SKK} : \beta$

- ▶  $\text{SK} : \alpha \rightarrow \beta \quad \text{K} : \alpha$
- ▶  $\text{S} : \gamma \rightarrow \delta \quad \text{K} : \gamma$
- ▶ unification problem

$$\alpha \approx \sigma_1 \rightarrow \tau_1 \rightarrow \sigma_1$$

## Example ①

principle type of  $\text{SKK} : \beta$

- ▶  $\text{SK} : \alpha \rightarrow \beta \quad \text{K} : \alpha$
- ▶  $\text{S} : \gamma \rightarrow \delta \quad \text{K} : \gamma$
- ▶ unification problem

$$\alpha \approx \sigma_1 \rightarrow \tau_1 \rightarrow \sigma_1 \quad \delta \approx \alpha \rightarrow \beta$$

## Example ①

principle type of  $\text{SKK} : \beta$

- ▶  $\text{SK} : \alpha \rightarrow \beta \quad \text{K} : \alpha$
- ▶  $\text{S} : \gamma \rightarrow \delta \quad \text{K} : \gamma$
- ▶ unification problem

$$\alpha \approx \sigma_1 \rightarrow \tau_1 \rightarrow \sigma_1 \quad \delta \approx \alpha \rightarrow \beta \quad \gamma \rightarrow \delta \approx (\rho_2 \rightarrow \sigma_2 \rightarrow \tau_2) \rightarrow (\rho_2 \rightarrow \sigma_2) \rightarrow \rho_2 \rightarrow \tau_2$$

## Example ①

principle type of  $\text{SKK} : \beta$

- ▶  $\text{SK} : \alpha \rightarrow \beta \quad \text{K} : \alpha$
- ▶  $\text{S} : \gamma \rightarrow \delta \quad \text{K} : \gamma$
- ▶ unification problem

$$\begin{aligned}\alpha &\approx \sigma_1 \rightarrow \tau_1 \rightarrow \sigma_1 & \delta &\approx \alpha \rightarrow \beta & \gamma \rightarrow \delta &\approx (\rho_2 \rightarrow \sigma_2 \rightarrow \tau_2) \rightarrow (\rho_2 \rightarrow \sigma_2) \rightarrow \rho_2 \rightarrow \tau_2 \\ \gamma &\approx \sigma_3 \rightarrow \tau_3 \rightarrow \sigma_3\end{aligned}$$

## Example ①

principle type of  $\text{SKK} : \beta$

- ▶  $\text{SK} : \alpha \rightarrow \beta \quad \text{K} : \alpha$
- ▶  $\text{S} : \gamma \rightarrow \delta \quad \text{K} : \gamma$
- ▶ unification problem

$$\begin{array}{lll} \alpha \approx \sigma_1 \rightarrow \tau_1 \rightarrow \sigma_1 & \delta \approx \alpha \rightarrow \beta & \gamma \rightarrow \delta \approx (\rho_2 \rightarrow \sigma_2 \rightarrow \tau_2) \rightarrow (\rho_2 \rightarrow \sigma_2) \rightarrow \rho_2 \rightarrow \tau_2 \\ \gamma \approx \sigma_3 \rightarrow \tau_3 \rightarrow \sigma_3 & \gamma \approx \rho_2 \rightarrow \sigma_2 \rightarrow \tau_2 & \delta \approx (\rho_2 \rightarrow \sigma_2) \rightarrow \rho_2 \rightarrow \tau_2 \end{array}$$

## Example ①

principle type of  $\text{SKK} : \beta$

- ▶  $\text{SK} : \alpha \rightarrow \beta \quad \text{K} : \alpha$
- ▶  $\text{S} : \gamma \rightarrow \delta \quad \text{K} : \gamma$
- ▶ unification problem

$$\begin{array}{lll} \alpha \approx \sigma_1 \rightarrow \tau_1 \rightarrow \sigma_1 & \delta \approx \alpha \rightarrow \beta \\ \gamma \approx \sigma_3 \rightarrow \tau_3 \rightarrow \sigma_3 & \gamma \approx \rho_2 \rightarrow \sigma_2 \rightarrow \tau_2 & \delta \approx (\rho_2 \rightarrow \sigma_2) \rightarrow \rho_2 \rightarrow \tau_2 \\ \alpha \approx \rho_2 \rightarrow \sigma_2 & \beta \approx \rho_2 \rightarrow \tau_2 & \end{array}$$

## Example ①

principle type of  $\text{SKK} : \beta$

- ▶  $\text{SK} : \alpha \rightarrow \beta \quad \text{K} : \alpha$
- ▶  $\text{S} : \gamma \rightarrow \delta \quad \text{K} : \gamma$
- ▶ unification problem

$$\alpha \approx \sigma_1 \rightarrow \tau_1 \rightarrow \sigma_1 \quad \delta \approx \alpha \rightarrow \beta$$

$$\gamma \approx \sigma_3 \rightarrow \tau_3 \rightarrow \sigma_3 \quad \gamma \approx \rho_2 \rightarrow \sigma_2 \rightarrow \tau_2$$

$$\alpha \approx \rho_2 \rightarrow \sigma_2 \quad \beta \approx \rho_2 \rightarrow \tau_2 \quad \rho_2 \approx \sigma_1 \quad \sigma_2 \approx \tau_1 \rightarrow \sigma_1$$

## Example ①

principle type of  $\text{SKK} : \beta$

- ▶  $\text{SK} : \alpha \rightarrow \beta \quad \text{K} : \alpha$
- ▶  $\text{S} : \gamma \rightarrow \delta \quad \text{K} : \gamma$
- ▶ unification problem

$$\begin{array}{lll} \alpha \approx \sigma_1 \rightarrow \tau_1 \rightarrow \sigma_1 & \delta \approx \alpha \rightarrow \beta \\ \gamma \approx \sigma_3 \rightarrow \tau_3 \rightarrow \sigma_3 & \gamma \approx \rho_2 \rightarrow \sigma_2 \rightarrow \tau_2 \\ & \beta \approx \rho_2 \rightarrow \tau_2 & \rho_2 \approx \sigma_1 \quad \sigma_2 \approx \tau_1 \rightarrow \sigma_1 \\ \sigma_3 \approx \rho_2 & \tau_3 \approx \sigma_2 & \sigma_3 \approx \tau_2 \end{array}$$

## Example ①

principle type of  $\text{SKK} : \beta$

- ▶  $\text{SK} : \alpha \rightarrow \beta \quad \text{K} : \alpha$
- ▶  $\text{S} : \gamma \rightarrow \delta \quad \text{K} : \gamma$
- ▶ unification problem

$$\alpha \approx \sigma_1 \rightarrow \tau_1 \rightarrow \sigma_1 \quad \delta \approx \alpha \rightarrow \beta$$

$$\gamma \approx \sigma_3 \rightarrow \tau_3 \rightarrow \sigma_3$$

$$\beta \approx \rho_2 \rightarrow \tau_2 \quad \sigma_2 \approx \tau_1 \rightarrow \sigma_1 \approx \tau_3$$

$$\sigma_3 \approx \rho_2 \approx \sigma_1 \approx \tau_2$$

## Example ①

principle type of **SKK** :  $\beta$

- ▶ **SK** :  $\alpha \rightarrow \beta$     **K** :  $\alpha$
- ▶ **S** :  $\gamma \rightarrow \delta$     **K** :  $\gamma$
- ▶ unification problem

$$\alpha \approx \sigma_1 \rightarrow \tau_1 \rightarrow \sigma_1 \quad \delta \approx \alpha \rightarrow \beta$$

$$\gamma \approx \sigma_3 \rightarrow \tau_3 \rightarrow \sigma_3$$

$$\beta \approx \rho_2 \rightarrow \tau_2 \quad \sigma_2 \approx \tau_1 \rightarrow \sigma_1 \approx \tau_3$$

$$\sigma_3 \approx \rho_2 \approx \sigma_1 \approx \tau_2$$

- ▶ mgu

$$\left\{ \begin{array}{ll} \alpha \mapsto \sigma_1 \rightarrow \tau_1 \rightarrow \sigma_1 & \gamma \mapsto \sigma_1 \rightarrow (\tau_1 \rightarrow \sigma_1) \rightarrow \sigma_1 \\ \beta \mapsto \sigma_1 \rightarrow \sigma_1 & \delta \mapsto (\sigma_1 \rightarrow \tau_1 \rightarrow \sigma_1) \rightarrow \sigma_1 \rightarrow \sigma_1 \end{array} \right\}$$

## Example ①

principle type of  $\text{SKK} : \sigma_1 \rightarrow \sigma_1$

►  $\text{SK} : \alpha \rightarrow \beta \quad \text{K} : \alpha$

►  $\text{S} : \gamma \rightarrow \delta \quad \text{K} : \gamma$

► unification problem

$$\alpha \approx \sigma_1 \rightarrow \tau_1 \rightarrow \sigma_1 \quad \delta \approx \alpha \rightarrow \beta$$

$$\gamma \approx \sigma_3 \rightarrow \tau_3 \rightarrow \sigma_3$$

$$\beta \approx \rho_2 \rightarrow \tau_2 \quad \sigma_2 \approx \tau_1 \rightarrow \sigma_1 \approx \tau_3$$

$$\sigma_3 \approx \rho_2 \approx \sigma_1 \approx \tau_2$$

► mgu

$$\left\{ \begin{array}{ll} \alpha \mapsto \sigma_1 \rightarrow \tau_1 \rightarrow \sigma_1 & \gamma \mapsto \sigma_1 \rightarrow (\tau_1 \rightarrow \sigma_1) \rightarrow \sigma_1 \\ \beta \mapsto \sigma_1 \rightarrow \sigma_1 & \delta \mapsto (\sigma_1 \rightarrow \tau_1 \rightarrow \sigma_1) \rightarrow \sigma_1 \rightarrow \sigma_1 \end{array} \right\}$$

## Example ②

SII

## Example ②

SII

- ▶ SI :  $\alpha \rightarrow \beta$     I :  $\alpha$

## Example ②

SII

- ▶ SI :  $\alpha \rightarrow \beta$     I :  $\alpha$
- ▶ unification problem

$$\alpha \approx \sigma_1 \rightarrow \sigma_1$$

## Example ②

SII

- ▶ **SI** :  $\alpha \rightarrow \beta$     | :  $\alpha$
- ▶ **S** :  $\gamma \rightarrow \delta$     | :  $\gamma$
- ▶ unification problem

$$\alpha \approx \sigma_1 \rightarrow \sigma_1$$

## Example ②

SII

- ▶ SI :  $\alpha \rightarrow \beta$    I :  $\alpha$
- ▶ S :  $\gamma \rightarrow \delta$    I :  $\gamma$
- ▶ unification problem

$$\alpha \approx \sigma_1 \rightarrow \sigma_1 \quad \delta \approx \alpha \rightarrow \beta$$

## Example ②

SII

- ▶ **SI** :  $\alpha \rightarrow \beta$     I :  $\alpha$
- ▶ **S** :  $\gamma \rightarrow \delta$     I :  $\gamma$
- ▶ unification problem

$$\alpha \approx \sigma_1 \rightarrow \sigma_1$$

$$\delta \approx \alpha \rightarrow \beta$$

$$\gamma \rightarrow \delta \approx (\rho_2 \rightarrow \sigma_2 \rightarrow \tau_2) \rightarrow (\rho_2 \rightarrow \sigma_2) \rightarrow \rho_2 \rightarrow \tau_2$$

## Example ②

SII

- ▶ **SI** :  $\alpha \rightarrow \beta$     **I** :  $\alpha$
- ▶ **S** :  $\gamma \rightarrow \delta$     **I** :  $\gamma$
- ▶ unification problem

$$\alpha \approx \sigma_1 \rightarrow \sigma_1$$

$$\gamma \approx \sigma_3 \rightarrow \sigma_3$$

$$\delta \approx \alpha \rightarrow \beta$$

$$\gamma \rightarrow \delta \approx (\rho_2 \rightarrow \sigma_2 \rightarrow \tau_2) \rightarrow (\rho_2 \rightarrow \sigma_2) \rightarrow \rho_2 \rightarrow \tau_2$$

## Example ②

SII

- ▶ **SI** :  $\alpha \rightarrow \beta$      $I : \alpha$
- ▶ **S** :  $\gamma \rightarrow \delta$      $I : \gamma$
- ▶ unification problem

$$\alpha \approx \sigma_1 \rightarrow \sigma_1$$

$$\gamma \approx \sigma_3 \rightarrow \sigma_3$$

$$\delta \approx \alpha \rightarrow \beta$$

$$\gamma \approx \rho_2 \rightarrow \sigma_2 \rightarrow \tau_2$$

$$\gamma \rightarrow \delta \approx (\rho_2 \rightarrow \sigma_2 \rightarrow \tau_2) \rightarrow (\rho_2 \rightarrow \sigma_2) \rightarrow \rho_2 \rightarrow \tau_2$$

$$\delta \approx (\rho_2 \rightarrow \sigma_2) \rightarrow \rho_2 \rightarrow \tau_2$$

## Example ②

SII

- ▶ **SI** :  $\alpha \rightarrow \beta$      $I : \alpha$
- ▶ **S** :  $\gamma \rightarrow \delta$      $I : \gamma$
- ▶ unification problem

$$\alpha \approx \sigma_1 \rightarrow \sigma_1$$

$$\delta \approx \alpha \rightarrow \beta$$

$$\gamma \approx \sigma_3 \rightarrow \sigma_3$$

$$\gamma \approx \rho_2 \rightarrow \sigma_2 \rightarrow \tau_2$$

$$\delta \approx (\rho_2 \rightarrow \sigma_2) \rightarrow \rho_2 \rightarrow \tau_2$$

$$\sigma_3 \approx \rho_2 \approx \sigma_2 \rightarrow \tau_2$$

## Example ②

SII

- ▶ **SI** :  $\alpha \rightarrow \beta$      $I : \alpha$
- ▶ **S** :  $\gamma \rightarrow \delta$      $I : \gamma$
- ▶ unification problem

$$\alpha \approx \sigma_1 \rightarrow \sigma_1$$

$$\delta \approx \alpha \rightarrow \beta$$

$$\gamma \approx \sigma_3 \rightarrow \sigma_3$$

$$\delta \approx (\rho_2 \rightarrow \sigma_2) \rightarrow \rho_2 \rightarrow \tau_2$$

$$\sigma_3 \approx \rho_2 \approx \sigma_2 \rightarrow \tau_2$$

$$\alpha \approx \rho_2 \rightarrow \sigma_2 \quad \beta \approx \rho_2 \rightarrow \tau_2$$

## Example ②

SII

- ▶ **SI** :  $\alpha \rightarrow \beta$     **I** :  $\alpha$
- ▶ **S** :  $\gamma \rightarrow \delta$     **I** :  $\gamma$
- ▶ unification problem

$$\alpha \approx \sigma_1 \rightarrow \sigma_1$$

$$\delta \approx \alpha \rightarrow \beta$$

$$\gamma \approx \sigma_3 \rightarrow \sigma_3$$

$$\sigma_3 \approx \rho_2 \approx \sigma_2 \rightarrow \tau_2$$

$$\alpha \approx \rho_2 \rightarrow \sigma_2$$

$$\beta \approx \rho_2 \rightarrow \tau_2$$

$$\sigma_1 \approx \rho_2 \approx \sigma_2$$

## Example ②

### SII

- ▶ **SI** :  $\alpha \rightarrow \beta$     | :  $\alpha$
- ▶ **S** :  $\gamma \rightarrow \delta$     | :  $\gamma$
- ▶ unification problem

$$\alpha \approx \sigma_1 \rightarrow \sigma_1 \qquad \delta \approx \alpha \rightarrow \beta$$

$$\gamma \approx \sigma_3 \rightarrow \sigma_3$$

$$\sigma_3 \approx \rho_2 \approx \sigma_2 \rightarrow \tau_2 \approx \sigma_1 \approx \sigma_2$$

$$\beta \approx \rho_2 \rightarrow \tau_2$$

## Example ②

SII

- ▶ **SI** :  $\alpha \rightarrow \beta$      $I : \alpha$
- ▶ **S** :  $\gamma \rightarrow \delta$      $I : \gamma$
- ▶ unification problem

$$\alpha \approx \sigma_1 \rightarrow \sigma_1 \qquad \delta \approx \alpha \rightarrow \beta$$

$$\gamma \approx \sigma_3 \rightarrow \sigma_3$$

$$\sigma_3 \approx \rho_2 \approx \sigma_2 \rightarrow \tau_2 \approx \sigma_1 \approx \sigma_2$$

$$\beta \approx \rho_2 \rightarrow \tau_2$$

has no solution

## Example ②

SII cannot be typed

- ▶ SI :  $\alpha \rightarrow \beta$    I :  $\alpha$
- ▶ S :  $\gamma \rightarrow \delta$    I :  $\gamma$
- ▶ unification problem

$$\alpha \approx \sigma_1 \rightarrow \sigma_1 \quad \delta \approx \alpha \rightarrow \beta$$

$$\gamma \approx \sigma_3 \rightarrow \sigma_3$$

$$\sigma_3 \approx \rho_2 \approx \sigma_2 \rightarrow \tau_2 \approx \sigma_1 \approx \sigma_2$$

$$\beta \approx \rho_2 \rightarrow \tau_2$$

has no solution

## Example ③

principle type of  $B = S(KS)K$

## Example ③

principle type of  $B = S(KS)K : \beta$

- ▶  $S(KS) : \alpha \rightarrow \beta$     $K : \alpha$

## Example ③

principle type of  $B = S(KS)K : \beta$

- ▶  $S(KS) : \alpha \rightarrow \beta$     $K : \alpha$

- ▶ unification problem

$$\alpha \approx \sigma_1 \rightarrow \tau_1 \rightarrow \sigma_1$$

## Example ③

principle type of  $B = S(KS)K : \beta$

►  $S(KS) : \alpha \rightarrow \beta \quad K : \alpha$

►  $S : \gamma \rightarrow \delta \quad KS : \gamma$

► unification problem

$$\alpha \approx \sigma_1 \rightarrow \tau_1 \rightarrow \sigma_1$$

## Example ③

principle type of  $B = S(KS)K : \beta$

- ▶  $S(KS) : \alpha \rightarrow \beta \quad K : \alpha$
- ▶  $S : \gamma \rightarrow \delta \quad KS : \gamma$
  
- ▶ unification problem

$$\alpha \approx \sigma_1 \rightarrow \tau_1 \rightarrow \sigma_1 \quad \delta \approx \alpha \rightarrow \beta$$

## Example ③

principle type of  $B = S(KS)K : \beta$

►  $S(KS) : \alpha \rightarrow \beta \quad K : \alpha$

►  $S : \gamma \rightarrow \delta \quad KS : \gamma$

► unification problem

$$\alpha \approx \sigma_1 \rightarrow \tau_1 \rightarrow \sigma_1 \quad \delta \approx \alpha \rightarrow \beta \quad \gamma \rightarrow \delta \approx (\rho_2 \rightarrow \sigma_2 \rightarrow \tau_2) \rightarrow (\rho_2 \rightarrow \sigma_2) \rightarrow \rho_2 \rightarrow \tau_2$$

## Example ③

principle type of  $B = S(KS)K : \beta$

- ▶  $S(KS) : \alpha \rightarrow \beta \quad K : \alpha$
- ▶  $S : \gamma \rightarrow \delta \quad KS : \gamma$
- ▶  $K : \epsilon \rightarrow \eta \quad S : \epsilon$
- ▶ unification problem

$$\alpha \approx \sigma_1 \rightarrow \tau_1 \rightarrow \sigma_1 \quad \delta \approx \alpha \rightarrow \beta \quad \gamma \rightarrow \delta \approx (\rho_2 \rightarrow \sigma_2 \rightarrow \tau_2) \rightarrow (\rho_2 \rightarrow \sigma_2) \rightarrow \rho_2 \rightarrow \tau_2$$

## Example ③

principle type of  $B = S(KS)K : \beta$

- ▶  $S(KS) : \alpha \rightarrow \beta \quad K : \alpha$
- ▶  $S : \gamma \rightarrow \delta \quad KS : \gamma$
- ▶  $K : \epsilon \rightarrow \eta \quad S : \epsilon$
- ▶ unification problem

$$\begin{aligned}\alpha &\approx \sigma_1 \rightarrow \tau_1 \rightarrow \sigma_1 & \delta &\approx \alpha \rightarrow \beta & \gamma \rightarrow \delta &\approx (\rho_2 \rightarrow \sigma_2 \rightarrow \tau_2) \rightarrow (\rho_2 \rightarrow \sigma_2) \rightarrow \rho_2 \rightarrow \tau_2 \\ \gamma &\approx \eta\end{aligned}$$

## Example ③

principle type of  $B = S(KS)K : \beta$

- ▶  $S(KS) : \alpha \rightarrow \beta \quad K : \alpha$
- ▶  $S : \gamma \rightarrow \delta \quad KS : \gamma$
- ▶  $K : \epsilon \rightarrow \eta \quad S : \epsilon$
- ▶ unification problem

$$\begin{aligned}\alpha &\approx \sigma_1 \rightarrow \tau_1 \rightarrow \sigma_1 & \delta &\approx \alpha \rightarrow \beta & \gamma \rightarrow \delta &\approx (\rho_2 \rightarrow \sigma_2 \rightarrow \tau_2) \rightarrow (\rho_2 \rightarrow \sigma_2) \rightarrow \rho_2 \rightarrow \tau_2 \\ \gamma &\approx \eta & \epsilon \rightarrow \eta &\approx \sigma_3 \rightarrow \tau_3 \rightarrow \sigma_3\end{aligned}$$

## Example ③

principle type of  $B = S(KS)K : \beta$

- ▶  $S(KS) : \alpha \rightarrow \beta \quad K : \alpha$
- ▶  $S : \gamma \rightarrow \delta \quad KS : \gamma$
- ▶  $K : \epsilon \rightarrow \eta \quad S : \epsilon$
- ▶ unification problem

$$\begin{array}{lll} \alpha \approx \sigma_1 \rightarrow \tau_1 \rightarrow \sigma_1 & \delta \approx \alpha \rightarrow \beta & \gamma \rightarrow \delta \approx (\rho_2 \rightarrow \sigma_2 \rightarrow \tau_2) \rightarrow (\rho_2 \rightarrow \sigma_2) \rightarrow \rho_2 \rightarrow \tau_2 \\ \gamma \approx \eta & \epsilon \rightarrow \eta \approx \sigma_3 \rightarrow \tau_3 \rightarrow \sigma_3 & \epsilon \approx (\rho_4 \rightarrow \sigma_4 \rightarrow \tau_4) \rightarrow (\rho_4 \rightarrow \sigma_4) \rightarrow \rho_4 \rightarrow \tau_4 \end{array}$$

### Example ③

principle type of  $B = S(KS)K : (\sigma_4 \rightarrow \tau_4) \rightarrow (\rho_4 \rightarrow \sigma_4) \rightarrow \rho_4 \rightarrow \tau_4$

- ▶  $S(KS) : \alpha \rightarrow \beta \quad K : \alpha$
- ▶  $S : \gamma \rightarrow \delta \quad KS : \gamma$
- ▶  $K : \epsilon \rightarrow \eta \quad S : \epsilon$
- ▶ unification problem

$$\begin{array}{lll} \alpha \approx \sigma_1 \rightarrow \tau_1 \rightarrow \sigma_1 & \delta \approx \alpha \rightarrow \beta & \gamma \rightarrow \delta \approx (\rho_2 \rightarrow \sigma_2 \rightarrow \tau_2) \rightarrow (\rho_2 \rightarrow \sigma_2) \rightarrow \rho_2 \rightarrow \tau_2 \\ \gamma \approx \eta & \epsilon \rightarrow \eta \approx \sigma_3 \rightarrow \tau_3 \rightarrow \sigma_3 & \epsilon \approx (\rho_4 \rightarrow \sigma_4 \rightarrow \tau_4) \rightarrow (\rho_4 \rightarrow \sigma_4) \rightarrow \rho_4 \rightarrow \tau_4 \end{array}$$

- ▶ mgu

$$\left\{ \beta \mapsto (\sigma_4 \rightarrow \tau_4) \rightarrow (\rho_4 \rightarrow \sigma_4) \rightarrow \rho_4 \rightarrow \tau_4 \quad \dots \right\}$$

# Outline

1. Summary of Previous Lecture
2. Strong Normalization
3. Type Inference
- 4. Type Inhabitation**
5. Intuitionistic Propositional Logic
6. Summary
7. Test

## Definition

type  $\tau$  is **inhabited** if  $\vdash t : \tau$  for some combinator  $t$

## Definition

type  $\tau$  is inhabited if  $\vdash t : \tau$  for some combinator  $t$

## Remark

not every type is inhabited

## Definition

type  $\tau$  is inhabited if  $\vdash t : \tau$  for some combinator  $t$

## Remark

not every type is inhabited

## Theorem

problem

instance: type  $\tau$

question: is  $\tau$  inhabited ?

is decidable

## Remark

$$I : \sigma \rightarrow \sigma$$

$$K : \sigma \rightarrow \tau \rightarrow \sigma$$

$$S : (\rho \rightarrow \sigma \rightarrow \tau) \rightarrow (\rho \rightarrow \sigma) \rightarrow \rho \rightarrow \tau$$

## Remark

$$\sigma \rightarrow \sigma$$

$$\sigma \rightarrow \tau \rightarrow \sigma$$

$$(\rho \rightarrow \sigma \rightarrow \tau) \rightarrow (\rho \rightarrow \sigma) \rightarrow \rho \rightarrow \tau$$

valid formulas in propositional logic

## Remark

$$\sigma \rightarrow \sigma$$

$$\sigma \rightarrow \tau \rightarrow \sigma$$

$$(\rho \rightarrow \sigma \rightarrow \tau) \rightarrow (\rho \rightarrow \sigma) \rightarrow \rho \rightarrow \tau$$

valid formulas in **intuitionistic** propositional logic

## Theorem (Curry-Howard)

type  $\tau$  (without type constants) is inhabited if and only if  $\tau$  is valid formula in implication fragment of intuitionistic propositional logic

# Outline

1. Summary of Previous Lecture
2. Strong Normalization
3. Type Inference
4. Type Inhabitation
5. Intuitionistic Propositional Logic

Kripke models

6. Summary
7. Test

## Syntax

- ▶ basic connectives  $\rightarrow \wedge \vee \perp$

## Syntax

- ▶ basic connectives  $\rightarrow \wedge \vee \perp$
- ▶ derived connectives
  - ▶  $\neg\varphi$  abbreviates  $\varphi \rightarrow \perp$

## Syntax

► basic connectives  $\rightarrow \wedge \vee \perp$

► derived connectives

►  $\neg\varphi$  abbreviates  $\varphi \rightarrow \perp$

►  $\top$  abbreviates  $\perp \rightarrow \perp$

## Syntax

- ▶ basic connectives  $\rightarrow \wedge \vee \perp$
- ▶ derived connectives
  - ▶  $\neg\varphi$  abbreviates  $\varphi \rightarrow \perp$
  - ▶  $\top$  abbreviates  $\perp \rightarrow \perp$
  - ▶  $\varphi \leftrightarrow \psi$  abbreviates  $(\varphi \rightarrow \psi) \wedge (\psi \rightarrow \varphi)$

## Syntax

- ▶ basic connectives  $\rightarrow \wedge \vee \perp$
- ▶ derived connectives
  - ▶  $\neg\varphi$  abbreviates  $\varphi \rightarrow \perp$
  - ▶  $\top$  abbreviates  $\perp \rightarrow \perp$
  - ▶  $\varphi \leftrightarrow \psi$  abbreviates  $(\varphi \rightarrow \psi) \wedge (\psi \rightarrow \varphi)$
- ▶ implication fragment contains only  $\rightarrow$

## Syntax

- ▶ basic connectives  $\rightarrow \wedge \vee \perp$
- ▶ derived connectives
  - ▶  $\neg\varphi$  abbreviates  $\varphi \rightarrow \perp$
  - ▶  $\top$  abbreviates  $\perp \rightarrow \perp$
  - ▶  $\varphi \leftrightarrow \psi$  abbreviates  $(\varphi \rightarrow \psi) \wedge (\psi \rightarrow \varphi)$
- ▶ implication fragment contains only  $\rightarrow$

## Formal Semantics

- ▶ Heyting algebras
- ▶ Kripke models

## Syntax

- ▶ basic connectives  $\rightarrow \wedge \vee \perp$
- ▶ derived connectives
  - ▶  $\neg\varphi$  abbreviates  $\varphi \rightarrow \perp$
  - ▶  $\top$  abbreviates  $\perp \rightarrow \perp$
  - ▶  $\varphi \leftrightarrow \psi$  abbreviates  $(\varphi \rightarrow \psi) \wedge (\psi \rightarrow \varphi)$
- ▶ implication fragment contains only  $\rightarrow$

## Formal Semantics

- ▶ Heyting algebras
- ▶ Kripke models

## Definition

Kripke model is triple  $\mathcal{C} = \langle C, \leq, \Vdash \rangle$

## Definition

Kripke model is triple  $\mathcal{C} = \langle C, \leq, \Vdash \rangle$  with

- ▶ non-empty set  $C$  of states

## Definition

Kripke model is triple  $\mathcal{C} = \langle C, \leqslant, \Vdash \rangle$  with

- ▶ non-empty set  $C$  of states
- ▶ partial order  $\leqslant$  on  $C$

## Definition

Kripke model is triple  $\mathcal{C} = \langle C, \leq, \Vdash \rangle$  with

- ▶ non-empty set  $C$  of states
- ▶ partial order  $\leq$  on  $C$
- ▶ binary relation  $\Vdash$  between elements of  $C$  and propositional atoms

## Definition

Kripke model is triple  $\mathcal{C} = \langle C, \leqslant, \Vdash \rangle$  with

- ▶ non-empty set  $C$  of states
- ▶ partial order  $\leqslant$  on  $C$
- ▶ binary relation  $\Vdash$  between elements of  $C$  and propositional atoms such that  $d \Vdash p$  whenever  $c \Vdash p$  and  $c \leqslant d$

## Definition

Kripke model is triple  $\mathcal{C} = \langle C, \leqslant, \Vdash \rangle$  with

- ▶ non-empty set  $C$  of states
- ▶ partial order  $\leqslant$  on  $C$
- ▶ binary relation  $\Vdash$  between elements of  $C$  and propositional atoms such that  $d \Vdash p$  whenever  $c \Vdash p$  and  $c \leqslant d$

## Definition

Kripke model  $\mathcal{C} = \langle C, \leqslant, \Vdash \rangle$ ,  $c \in C$

- ▶  $c \Vdash \varphi \wedge \psi$  if and only if  $c \Vdash \varphi$  and  $c \Vdash \psi$

## Definition

Kripke model is triple  $\mathcal{C} = \langle C, \leqslant, \Vdash \rangle$  with

- ▶ non-empty set  $C$  of states
- ▶ partial order  $\leqslant$  on  $C$
- ▶ binary relation  $\Vdash$  between elements of  $C$  and propositional atoms such that  $d \Vdash p$  whenever  $c \Vdash p$  and  $c \leqslant d$

## Definition

Kripke model  $\mathcal{C} = \langle C, \leqslant, \Vdash \rangle$ ,  $c \in C$

- ▶  $c \Vdash \varphi \wedge \psi$  if and only if  $c \Vdash \varphi$  and  $c \Vdash \psi$
- ▶  $c \Vdash \varphi \vee \psi$  if and only if  $c \Vdash \varphi$  or  $c \Vdash \psi$

## Definition

Kripke model is triple  $\mathcal{C} = \langle C, \leqslant, \Vdash \rangle$  with

- ▶ non-empty set  $C$  of states
- ▶ partial order  $\leqslant$  on  $C$
- ▶ binary relation  $\Vdash$  between elements of  $C$  and propositional atoms such that  $d \Vdash p$  whenever  $c \Vdash p$  and  $c \leqslant d$

## Definition

Kripke model  $\mathcal{C} = \langle C, \leqslant, \Vdash \rangle$ ,  $c \in C$

- ▶  $c \Vdash \varphi \wedge \psi$  if and only if  $c \Vdash \varphi$  and  $c \Vdash \psi$
- ▶  $c \Vdash \varphi \vee \psi$  if and only if  $c \Vdash \varphi$  or  $c \Vdash \psi$
- ▶  $c \Vdash \varphi \rightarrow \psi$  if and only if  $d \Vdash \psi$  for all  $d \geqslant c$  with  $d \Vdash \varphi$

## Definition

Kripke model is triple  $\mathcal{C} = \langle C, \leqslant, \Vdash \rangle$  with

- ▶ non-empty set  $C$  of states
- ▶ partial order  $\leqslant$  on  $C$
- ▶ binary relation  $\Vdash$  between elements of  $C$  and propositional atoms such that  $d \Vdash p$  whenever  $c \Vdash p$  and  $c \leqslant d$

## Definition

Kripke model  $\mathcal{C} = \langle C, \leqslant, \Vdash \rangle$ ,  $c \in C$

- ▶  $c \Vdash \varphi \wedge \psi$  if and only if  $c \Vdash \varphi$  and  $c \Vdash \psi$
- ▶  $c \Vdash \varphi \vee \psi$  if and only if  $c \Vdash \varphi$  or  $c \Vdash \psi$
- ▶  $c \Vdash \varphi \rightarrow \psi$  if and only if  $d \Vdash \psi$  for all  $d \geqslant c$  with  $d \Vdash \varphi$
- ▶  $c \not\Vdash \perp$

## Terminology

$c$  forces  $p$  if  $c \Vdash p$

$c$  forces  $p$  if  $c \Vdash p$

### Example

Kripke model  $\mathcal{C} = \langle C, \leqslant, \Vdash \rangle$  with  $C = \{a, b, c\}$ ,  $a \leqslant b$ ,  $a \leqslant c$ ,  $b \Vdash p$ ,  $c \Vdash q$

$c$  forces  $p$  if  $c \Vdash p$

### Example

Kripke model  $\mathcal{C} = \langle C, \leqslant, \Vdash \rangle$  with  $C = \{a, b, c\}$ ,  $a \leqslant b$ ,  $a \leqslant c$ ,  $b \Vdash p$ ,  $c \Vdash q$

- ▶  $a \Vdash (p \rightarrow q) \rightarrow q$

$c$  forces  $p$  if  $c \Vdash p$

### Example

Kripke model  $\mathcal{C} = \langle C, \leqslant, \Vdash \rangle$  with  $C = \{a, b, c\}$ ,  $a \leqslant b$ ,  $a \leqslant c$ ,  $b \Vdash p$ ,  $c \Vdash q$

- ▶  $a \Vdash (p \rightarrow q) \rightarrow q$
- ▶  $a \Vdash \neg\neg(p \vee q)$

$c$  forces  $p$  if  $c \Vdash p$

### Example

Kripke model  $\mathcal{C} = \langle C, \leqslant, \Vdash \rangle$  with  $C = \{a, b, c\}$ ,  $a \leqslant b$ ,  $a \leqslant c$ ,  $b \Vdash p$ ,  $c \Vdash q$

- ▶  $a \Vdash (p \rightarrow q) \rightarrow q$
- ▶  $a \Vdash \neg\neg(p \vee q)$
- ▶  $a \nVdash p \vee \neg p$

## Terminology

$c$  forces  $p$  if  $c \Vdash p$

### Example

Kripke model  $\mathcal{C} = \langle C, \leqslant, \Vdash \rangle$  with  $C = \{a, b, c\}$ ,  $a \leqslant b$ ,  $a \leqslant c$ ,  $b \Vdash p$ ,  $c \Vdash q$

- ▶  $a \Vdash (p \rightarrow q) \rightarrow q$
- ▶  $a \Vdash \neg\neg(p \vee q)$
- ▶  $a \nvDash p \vee \neg p$

### Definition

Kripke model  $\mathcal{C} = \langle C, \leqslant, \Vdash \rangle$ ,  $c \in C$

- ▶  $c \Vdash \Gamma$  if  $c \Vdash \varphi$  for all  $\varphi \in \Gamma$

## Terminology

$c$  forces  $p$  if  $c \Vdash p$

### Example

Kripke model  $\mathcal{C} = \langle C, \leq, \Vdash \rangle$  with  $C = \{a, b, c\}$ ,  $a \leq b$ ,  $a \leq c$ ,  $b \Vdash p$ ,  $c \Vdash q$

- ▶  $a \Vdash (p \rightarrow q) \rightarrow q$
- ▶  $a \Vdash \neg\neg(p \vee q)$
- ▶  $a \nvDash p \vee \neg p$

### Definition

Kripke model  $\mathcal{C} = \langle C, \leq, \Vdash \rangle$ ,  $c \in C$

- ▶  $c \Vdash \Gamma$  if  $c \Vdash \varphi$  for all  $\varphi \in \Gamma$
- ▶  $\mathcal{C} \Vdash \varphi$  if  $c \Vdash \varphi$  for all  $c \in C$

## Definition

$\Gamma \Vdash \varphi$  if  $c \Vdash \varphi$  whenever  $c \Vdash \Gamma$  for all Kripke models  $\mathcal{C} = \langle C, \leq, \Vdash \rangle$  and  $c \in C$

## Definition

$\Gamma \Vdash \varphi$  if  $c \Vdash \varphi$  whenever  $c \Vdash \Gamma$  for all Kripke models  $\mathcal{C} = \langle C, \leqslant, \Vdash \rangle$  and  $c \in C$

## Lemma (Monotonicity)

if  $c \leqslant d$  and  $c \Vdash \varphi$  then  $d \Vdash \varphi$

## Definition

$\Gamma \Vdash \varphi$  if  $c \Vdash \varphi$  whenever  $c \Vdash \Gamma$  for all Kripke models  $\mathcal{C} = \langle C, \leqslant, \Vdash \rangle$  and  $c \in C$

## Lemma (Monotonicity)

if  $c \leqslant d$  and  $c \Vdash \varphi$  then  $d \Vdash \varphi$

## Lemma

if  $\Vdash \varphi \vee \psi$  then  $\Vdash \varphi$  or  $\Vdash \psi$

# Outline

1. Summary of Previous Lecture
2. Strong Normalization
3. Type Inference
4. Type Inhabitation
5. Intuitionistic Propositional Logic
6. Summary
7. Test

## Important Concepts

- ▶  $\Vdash$
- ▶ implication fragment
- ▶ intuitionistic propositional logic
- ▶ Kripke model
- ▶ principal type
- ▶ SC
- ▶ type inference
- ▶ type inhabitation

## Important Concepts

- ▶  $\Vdash$
- ▶ implication fragment
- ▶ intuitionistic propositional logic
- ▶ Kripke model
- ▶ principal type
- ▶ SC
- ▶ type inference
- ▶ type inhabitation

homework for January 8

## Important Concepts

- ▶  $\Vdash$
- ▶ implication fragment
- ▶ intuitionistic propositional logic
- ▶ Kripke model
- ▶ principal type
- ▶ SC
- ▶ type inference
- ▶ type inhabitation

homework for January 8

next lecture (January 8): online evaluation in presence

## Important Concepts

- ▶  $\Vdash$
- ▶ implication fragment
- ▶ intuitionistic propositional logic
- ▶ Kripke model
- ▶ principal type
- ▶ SC
- ▶ type inference
- ▶ type inhabitation

homework for January 8

next lecture (January 8): online evaluation in presence  $\implies$  bring device

# Outline

1. Summary of Previous Lecture
2. Strong Normalization
3. Type Inference
4. Type Inhabitation
5. Intuitionistic Propositional Logic
6. Summary
7. Test

## Test on January 29

- ▶ 15:15–18:00 in HS 10

## Test on January 29

- ▶ 15:15–18:00 in HS 10
- ▶ online registration required before 10 am on January 23

## Test on January 29

- ▶ 15:15–18:00 in HS 10
- ▶ online registration required before 10 am on January 23
- ▶ closed book

## Test on January 29

- ▶ 15:15–18:00 in HS 10
- ▶ online registration required before 10 am on January 23
- ▶ closed book

## Earlier Exams / Tests

- ▶ SS 2022 (test)
- ▶ WS 2014 – 1
- ▶ SS 2007
- ▶ WS 2017 – 2
- ▶ SS 2012
- ▶ SS 2006 – 2
- ▶ WS 2017 – 1
- ▶ SS 2008 – 2
- ▶ SS 2006 – 1
- ▶ WS 2014 – 2
- ▶ SS 2008 – 1
- ▶ WS 2004