



Automata and Logic

Aart Middeldorp and Samuel Frontull

Definitions

► **deterministic finite automaton (DFA)** is quintuple $M = (Q, \Sigma, \delta, s, F)$ with

- ① Q : finite set of **states**
- ② Σ : **input alphabet**
- ③ $\delta: Q \times \Sigma \rightarrow Q$: **transition function**
- ④ $s \in Q$: **start state**
- ⑤ $F \subseteq Q$: **final (accept) states**

► $\hat{\delta}: Q \times \Sigma^* \rightarrow Q$ is inductively defined by

$$\hat{\delta}(q, \epsilon) = q \qquad \hat{\delta}(q, xa) = \delta(\hat{\delta}(q, x), a)$$

- string $x \in \Sigma^*$ is **accepted** by M if $\hat{\delta}(s, x) \in F$
- string $x \in \Sigma^*$ is **rejected** by M if $\hat{\delta}(s, x) \notin F$
- language accepted by M : $L(M) = \{x \in \Sigma^* \mid \hat{\delta}(s, x) \in F\}$

Outline

1. Summary of Previous Lecture
2. Nondeterministic Finite Automata
3. Epsilon Transitions
4. Intermezzo
5. Closure Properties
6. Hamming Distance
7. Further Reading

Definition

set $A \subseteq \Sigma^*$ is **regular** if $A = L(M)$ for some DFA M

Theorem

regular sets are **effectively** closed under **intersection**, **union**, and **complement**

Automata

- ▶ (deterministic, **nondeterministic**, alternating) **finite automata**
- ▶ regular expressions
- ▶ (alternating) Büchi automata

Logic


- ▶ (weak) monadic second-order logic
- ▶ Presburger arithmetic
- ▶ linear-time temporal logic

Outline

1. Summary of Previous Lecture
2. **Nondeterministic Finite Automata**
3. Epsilon Transitions
4. Intermezzo
5. Closure Properties
6. Hamming Distance
7. Further Reading

Example

NFA $M = (Q, \Sigma, \Delta, S, F)$



1 $Q = \{1, 2, 3\}$

2 $\Sigma = \{a, b\}$

3 $\Delta: Q \times \Sigma \rightarrow 2^Q$

4 $S = \{1\}$

5 $F = \{3\}$

Δ	a	b
1	$\{1, 2\}$	$\{1\}$
2	$\{3\}$	$\{1, 3\}$
3	$\{3\}$	$\emptyset$

Definitions

▶ **nondeterministic finite automaton (NFA)** is quintuple $N = (Q, \Sigma, \Delta, S, F)$ with

- ① Q : finite set of states
- ② Σ : input alphabet
- ③ $\Delta: Q \times \Sigma \rightarrow 2^Q$: transition function
- ④ $S \subseteq Q$: **set** of start states
- ⑤ $F \subseteq Q$: final (accept) states

▶ $\hat{\Delta}: 2^Q \times \Sigma^* \rightarrow 2^Q$ is inductively defined by

$$\hat{\Delta}(A, \epsilon) = A$$

$$\hat{\Delta}(A, xa) = \bigcup_{q \in \hat{\Delta}(A, x)} \Delta(q, a)$$

▶ $x \in \Sigma^*$ is accepted by N if $\hat{\Delta}(S, x) \cap F \neq \emptyset$

▶ $L(N) = \{x \in \Sigma^* \mid \hat{\Delta}(S, x) \cap F \neq \emptyset\}$

Theorem

every set accepted by NFA is regular

Proof (subset construction)

► NFA $N = (Q_N, \Sigma, \Delta_N, S_N, F_N)$

► $L(N) = L(M)$ for DFA $M = (Q_M, \Sigma, \delta_M, S_M, F_M)$ with

① $Q_M = 2^{Q_N}$

② $\delta_M(A, a) = \widehat{\Delta}_N(A, a)$ for all $A \subseteq Q_N$ and $a \in \Sigma$


③ $S_M = S_N$

④ $F_M = \{A \subseteq Q_N \mid A \cap F_N \neq \emptyset\}$

► claim: $\widehat{\delta}_M(A, x) = \widehat{\Delta}_N(A, x)$ for all $A \subseteq Q_N$ and $x \in \Sigma^*$

proof of claim: easy induction on $|x|$

Example




$A = \emptyset$ $E = \{1, 2\}$
 $B = \{1\}$ $F = \{1, 3\}$
 $C = \{2\}$ $G = \{2, 3\}$
 $D = \{3\}$ $H = \{1, 2, 3\}$

	a	b		a	b
A	A	A	E	H	F
B	E	B	F	H	B
C	D	F	G	D	F
D	D	A	H	H	F

abbbaababbabbbaababba

remove inaccessible states



Outline

1. Summary of Previous Lecture

2. Nondeterministic Finite Automata

3. Epsilon Transitions


4. Intermezzo

5. Closure Properties

6. Hamming Distance

7. Further Reading

Example



$C_\epsilon(\{1\}) = \{1, 2, 3\}$
 $\widehat{\Delta}(\{1\}, b) = \{2, 3\}$

Definitions

► NFA with ϵ -transitions (NFA_ϵ) is sextuple $N = (Q, \Sigma, \epsilon, \Delta, S, F)$ such that

① $\epsilon \notin \Sigma$

② $N_\epsilon = (Q, \Sigma \cup \{\epsilon\}, \Delta, S, F)$ is NFA over alphabet $\Sigma \cup \{\epsilon\}$

► ϵ -closure of set $A \subseteq Q$ is defined as $C_\epsilon(A) = \bigcup \{\widehat{\Delta}_{N_\epsilon}(A, x) \mid x \in \{\epsilon\}^*\}$

► $\widehat{\Delta}_N: 2^Q \times \Sigma^* \rightarrow 2^Q$ is inductively defined by

$$\widehat{\Delta}_N(A, \epsilon) = C_\epsilon(A)$$

$$\widehat{\Delta}_N(A, xa) = \bigcup \{C_\epsilon(\Delta(q, a)) \mid q \in \widehat{\Delta}_N(A, x)\}$$

Lemma

$C_\epsilon(A)$ is least extension of A that is closed under ϵ -transitions:

$$q \in C_\epsilon(A) \implies \Delta_{N_\epsilon}(q, \epsilon) \subseteq C_\epsilon(A)$$


Theorem

every set accepted by NFA_ϵ is regular

Proof (construction)

- ▶ $NFA_\epsilon N_1 = (Q, \Sigma, \epsilon, \Delta_1, S, F_1)$
- ▶ $L(N_1) = L(N_2)$ for $NFA N_2 = (Q, \Sigma, \Delta_2, S, F_2)$ with
 - ① $\Delta_2(q, a) = \hat{\Delta}_1(\{q\}, a)$ for all $q \in Q$ and $a \in \Sigma$
 - ② $F_2 = \{q \mid C_\epsilon(\{q\}) \cap F_1 \neq \emptyset\}$


Example



Example

$NFA_\epsilon N_1 = (\{1, 2, 3\}, \{a, b, c\}, \epsilon, \Delta_1, \{1\}, \{3\})$ with


Δ_1	a	b	c	ϵ
1	$\{1\}$	$\emptyset$	$\emptyset$	$\{2\}$
2	$\emptyset$	$\{2\}$	$\emptyset$	$\{3\}$
3	$\emptyset$	$\emptyset$	$\{3\}$	$\emptyset$



$NFA N_2 = (\{1, 2, 3\}, \{a, b, c\}, \Delta_2, \{1\}, F_2)$ with

▶ $F_2 = \{q \mid C_\epsilon(\{q\}) \cap \{3\} \neq \emptyset\} = \{q \mid 3 \in C_\epsilon(\{q\})\} = \{1, 2, 3\}$

Δ_2	a	b	c
1	$\{1, 2, 3\}$	$\{2, 3\}$	$\{3\}$
2	$\emptyset$	$\{2, 3\}$	$\{3\}$
3	$\emptyset$	$\emptyset$	$\{3\}$



Outline

1. Summary of Previous Lecture
2. Nondeterministic Finite Automata
3. Epsilon Transitions
4. Intermezzo
5. Closure Properties
6. Hamming Distance
7. Further Reading

Question

What is the language accepted by the NFA_ε given by the following transition table ?

		ε	a	b
→ 1		∅	{2}	∅
2		{3}	∅	∅
3		∅	{4}	{2}
4 F		{1}	∅	∅

- A $\{xyx \mid x \in \{a\} \text{ and } y \in \{b\}^*\}$
- B the set of all strings over $\{a, b\}$ starting and ending with a
- C $\{xyz \mid x, z \in \{a, b\} \text{ and } y \in \{aab\}^*\}$
- D $\{\{a\}\{b\}^*\{a\}\}^+$



Theorem

regular sets are **effectively** closed under **union**, concatenation, and asterate

Proof



- ▶ $A = L(N_1)$ for NFA $N_1 = (Q_1, \Sigma, \Delta_1, S_1, F_1)$
 $B = L(N_2)$ for NFA $N_2 = (Q_2, \Sigma, \Delta_2, S_2, F_2)$
- ▶ without loss of generality $Q_1 \cap Q_2 = \emptyset$
- ▶ $A \cup B = L(N)$ for NFA $N = (Q, \Sigma, \Delta, S, F)$ with
 - ① $Q = Q_1 \cup Q_2$
 - ② $S = S_1 \cup S_2$
 - ③ $F = F_1 \cup F_2$
 - ④ $\Delta(q, a) = \begin{cases} \Delta_1(q, a) & \text{if } q \in Q_1 \\ \Delta_2(q, a) & \text{if } q \in Q_2 \end{cases}$

Outline

1. Summary of Previous Lecture
2. Nondeterministic Finite Automata
3. Epsilon Transitions
4. Intermezzo
5. Closure Properties
6. Hamming Distance
7. Further Reading

Example

$\{x \in \{a\}^* \mid |x| \text{ is divisible by 3}\} \cup \{x \in \{a\}^* \mid |x| \text{ is divisible by 4}\}$



$\{x \in \{a\}^* \mid |x| \text{ is divisible by 3 or 4}\}$

Theorem


regular sets are **effectively** closed under union, **concatenation**, and asterate

Proof

- ▶ $A = L(N_1)$ for NFA $N_1 = (Q_1, \Sigma, \Delta_1, S_1, F_1)$
- $B = L(N_2)$ for NFA $N_2 = (Q_2, \Sigma, \Delta_2, S_2, F_2)$
- ▶ without loss of generality $Q_1 \cap Q_2 = \emptyset$
- ▶ $AB = L(N)$ for NFA $_{\epsilon}$ $N = (Q, \Sigma, \epsilon, \Delta, S_1, F_2)$ with
 - ① $Q = Q_1 \cup Q_2$
 - ② $\Delta(q, a) = \begin{cases} \Delta_1(q, a) & \text{if } q \in Q_1 \text{ and } a \in \Sigma \\ \Delta_2(q, a) & \text{if } q \in Q_2 \text{ and } a \in \Sigma \\ S_2 & \text{if } q \in F_1 \text{ and } a = \epsilon \\ \emptyset & \text{otherwise} \end{cases}$

Example

$\{x \in \{a\}^* \mid |x| \text{ is divisible by } 3\}$ $\{x \in \{a\}^* \mid |x| \text{ is divisible by } 4\}$



$\{x \in \{a\}^* \mid |x| \notin \{1, 2, 5\}\}$

Theorem


regular sets are **effectively** closed under union, concatenation, and **asterate**

Proof


- ▶ $A = L(N_1)$ for NFA $N_1 = (Q_1, \Sigma, \Delta_1, S_1, F_1)$
- ▶ $A^* = L(N)$ for NFA $_{\epsilon}$ $N = (Q, \Sigma, \epsilon, \Delta, S, F)$ with
 - ① $Q = Q_1 \uplus \{s\}$
 - ② $S = \{s\}$
 - ③ $F = \{s\}$
 - ④ $\Delta(q, a) = \begin{cases} \Delta_1(q, a) & \text{if } q \in Q_1 \text{ and } a \in \Sigma \\ S_1 & \text{if } q = s \text{ and } a = \epsilon \\ S & \text{if } q \in F_1 \text{ and } a = \epsilon \\ \emptyset & \text{otherwise} \end{cases}$

Example

$\{a\}^* \{b\}^*$



$(\{a\}^* \{b\}^*)^* = \{a, b\}^*$



Outline

1. Summary of Previous Lecture
2. Nondeterministic Finite Automata
3. Epsilon Transitions
4. Intermezzo
5. Closure Properties
- 6. Hamming Distance**
7. Further Reading

Definitions

- ▶ **Hamming distance** $H(x, y)$ is number of places where bit strings x and y differ
- ▶ if $|x| \neq |y|$ then $H(x, y) = \infty$
- ▶ $N_k(A) = \{x \in \{0, 1\}^* \mid H(x, y) \leq k \text{ for some } y \in A\}$

Lemma

$A \subseteq \{0, 1\}^*$ is regular $\implies N_2(A)$ is regular

Lemma

$A \subseteq \{0, 1\}^*$ is regular $\implies N_2(A)$ is regular

Proof

- ▶ $A = L(M)$ for DFA $M = (Q_M, \{0, 1\}, \delta_M, s_M, F_M)$
- ▶ define NFA $N = (Q_N, \{0, 1\}, \Delta_N, s_N, F_N)$ with
 - ① $Q_N = Q_M \times \{0, 1, 2\}$
 - ② $\Delta_N((p, 0), a) = \{(q, 0) \mid \delta_M(p, a) = q\} \cup \{(q, 1) \mid \delta_M(p, b) = q \text{ for some } b \neq a\}$
 $\Delta_N((p, 1), a) = \{(q, 1) \mid \delta_M(p, a) = q\} \cup \{(q, 2) \mid \delta_M(p, b) = q \text{ for some } b \neq a\}$
 $\Delta_N((p, 2), a) = \{(q, 2) \mid \delta_M(p, a) = q\}$ for all $a \in \Sigma$
 - ③ $s_N = \{(s_M, 0)\}$
 - ④ $F_N = F_M \times \{0, 1, 2\}$

Proof (cont'd)

- ▶ key property:

$$(q, j) \in \widehat{\Delta}_N(\{(p, i)\}, y) \iff \widehat{\delta}_M(p, x) = q \text{ for some } x \in \{0, 1\}^*$$

for all $p, q \in Q_M$, $y \in \{0, 1\}^*$, $i, j \in \{0, 1, 2\}$ such that $|x| = |y|$ and $H(x, y) = j - i$

- ▶ $N_2(A) = \{y \mid H(y, x) \leq 2 \text{ for some } x \in A\}$
 $= \{y \mid H(y, x) = k \text{ for some } x \in A \text{ and } k \in \{0, 1, 2\}\}$
 $= \{y \mid H(y, x) = k \text{ and } \widehat{\delta}_M(s_M, x) = q \text{ for some } x \in A, k \in \{0, 1, 2\} \text{ and } q \in F_M\}$
 $= \{y \mid (q, k) \in \widehat{\Delta}_N(\{(s_M, 0)\}, y) \text{ for some } q \in F_M \text{ and } k \in \{0, 1, 2\}\}$
 $= \{y \mid (q, k) \in \widehat{\Delta}_N(\{(s_M, 0)\}, y) \text{ for some } (q, k) \in F_N\}$
 $= \{y \mid \widehat{\Delta}_N(\{(s_M, 0)\}, y) \cap F_N \neq \emptyset\}$
 $= L(N)$

Outline

- 1. Summary of Previous Lecture
- 2. Nondeterministic Finite Automata
- 3. Epsilon Transitions
- 4. Intermezzo
- 5. Closure Properties
- 6. Hamming Distance
- 7. Further Reading

Kozen

- ▶ Lecture 5 and 6

Important Concepts

- | | | |
|--------------------------|--------------------|-----------------------|
| ▶ ϵ -transition | ▶ Hamming distance | ▶ NFA_{ϵ} |
| ▶ ϵ -closure | ▶ NFA | ▶ subset construction |
| ▶ asterate | | |

homework for October 24