

# Commutation via Relative Termination\*

Nao Hirokawa  
JAIST, Japan  
hirokawa@jaist.ac.jp

Aart Middeldorp  
University of Innsbruck, Austria  
aart.middeldorp@uibk.ac.at

## Abstract

We present a generalisation of a commutation criterion by Rosen (1973), the development closedness theorem by van Oostrom (1994), and a confluence criterion by Hirokawa and Middeldorp (2011).

## 1 Introduction

This note is about sufficient conditions for the *commutation* of two *left-linear* TRSs  $\mathcal{R}$  and  $\mathcal{S}$ . Commutation means that the inclusion

$$\overset{*}{\leftarrow} \cdot \overset{*}{\rightarrow} \subseteq \overset{*}{\rightarrow} \cdot \overset{*}{\leftarrow}$$

holds. When  $\mathcal{R} = \mathcal{S}$ , commutation amounts to confluence. Furthermore, commutation is a sufficient condition for the confluence of the union of two confluent TRSs. It is well-known that  $\mathcal{R}$  and  $\mathcal{S}$  commute when there are no critical pairs between the rules of  $\mathcal{R}$  and the rules of  $\mathcal{S}$ . Toyama [6] generalized this result of Rosen [4] (and Raoult and Vuillemin [3, Proposition 10]) by admitting critical pairs between  $\mathcal{R}$  and  $\mathcal{S}$  but restricting the way in which they are joined. More precisely, Toyama showed that  $\mathcal{R}$  and  $\mathcal{S}$  commute if the following two conditions are satisfied:

$$\overset{\leftarrow}{\mathcal{R}} \times \overset{\rightarrow}{\mathcal{S}} \subseteq \overset{\#}{\mathcal{S}} \cdot \overset{*}{\leftarrow} \quad \text{and} \quad \overset{>\epsilon}{\leftarrow} \times \overset{\rightarrow}{\mathcal{R}} \subseteq \overset{\#}{\mathcal{R}}$$

As observed in [7, Corollaries 24 and 28] and [1, Proposition 7], Toyama's result is further generalized by adopting multi-steps:

$$\overset{\leftarrow}{\mathcal{R}} \times \overset{\rightarrow}{\mathcal{S}} \subseteq \overset{\circ}{\mathcal{S}} \cdot \overset{*}{\leftarrow} \quad \text{and} \quad \overset{>\epsilon}{\leftarrow} \times \overset{\rightarrow}{\mathcal{R}} \subseteq \overset{\circ}{\mathcal{R}}$$

We present a generalisation of these results by incorporating the confluence result based on relative termination [2].

## 2 Preliminaries

We assume familiarity with the conversion version of the decreasing diagrams technique for proving commutation, due to van Oostrom [8, Theorem 3]. Familiarity with proof terms witnessing multi-steps ([5, Chapter 8], [2]) is also helpful. Below we recall some basic definitions and recast a measure used in [7] as a measure on co-initial proof terms.

Let  $\mathcal{R}$  be a TRS over a signature  $\mathcal{F}$ . For each rule  $\ell \rightarrow r \in \mathcal{R}$  we introduce a fresh *rule symbol*  $\ell \rightarrow r$  whose arity is given by the number of variables in  $\ell$ . *Proof terms* are terms over functions in  $\mathcal{F}$  and rule symbols. We write  $\sqsubseteq$  for the smallest rewrite preorder on proof terms such that  $\ell \sqsubseteq \ell \rightarrow r(x_1, \dots, x_n)$  for all rules  $\ell \rightarrow r \in \mathcal{R}$  with  $\text{var}(\ell) = (x_1, \dots, x_n)$ . Here  $\text{var}(\ell)$  denotes a sequence consisting of all variables in  $\text{Var}(\ell)$  in some fixed order. Given co-initial

---

\*Supported by the JSPS KAKENHI Grant Number 25730004 and the Austrian Science Fund (FWF) P22467.

proof terms  $A$  and  $B$ , the (partial) residual operation is the proof term  $A \setminus B$  defined by the following clauses (here  $\langle A_1, \dots, A_n \rangle_\ell$  stands for the substitution  $\{x_i \mapsto A_i \mid 1 \leq i \leq n\}$ ):

$$\begin{aligned} x \setminus x &= x \\ f(A_1, \dots, A_n) \setminus f(B_1, \dots, B_n) &= f(A_1 \setminus B_1, \dots, A_n \setminus B_n) \\ \underline{\ell \rightarrow r}(A_1, \dots, A_n) \setminus \underline{\ell \rightarrow r}(B_1, \dots, B_n) &= r \langle A_1 \setminus B_1, \dots, A_n \setminus B_n \rangle_\ell \\ \underline{\ell \rightarrow r}(A_1, \dots, A_n) \setminus \underline{\ell \langle B_1, \dots, B_n \rangle}_\ell &= \underline{\ell \rightarrow r}(A_1 \setminus B_1, \dots, A_n \setminus B_n) \\ \underline{\ell \langle A_1, \dots, A_n \rangle}_\ell \setminus \underline{\ell \rightarrow r}(B_1, \dots, B_n) &= r \langle A_1 \setminus B_1, \dots, A_n \setminus B_n \rangle_\ell \end{aligned}$$

When we use  $A \setminus B$  below, it is guaranteed to be defined. (If  $A \sqsubseteq B$  then  $A \setminus B$  is a proof term without rule symbols.) A *redex* is a proof term that contains exactly one rule symbol. For a redex  $\Delta$  we write  $\Delta \in A$  if  $\Delta \sqsubseteq A$ . Given a proof term  $A$  and a redex  $\Delta \in A$ , we find it convenient to write  $A - \Delta$  for the proof term that is obtained from  $A$  by replacing the subterm occurrence  $\underline{\ell \rightarrow r}(A_1, \dots, A_n)$  corresponding to  $\Delta$  by  $\underline{\ell \langle A_1, \dots, A_n \rangle}_\ell$ .

For redexes  $\Delta_1, \Delta_2 \in A$  we write  $\Delta_1 \geq \Delta_2$  if the rule symbol of  $\Delta_1$  appears at or below that of  $\Delta_2$  in  $A$ . The set  $\blacktriangle(A)$  of positions is inductively defined on  $A$  as follows:

$$\begin{aligned} \blacktriangle(x) &= \emptyset \\ \blacktriangle(f(A_1, \dots, A_n)) &= \{iq \mid 1 \leq i \leq n \text{ and } q \in \blacktriangle(A_i)\} \\ \blacktriangle(\underline{\ell \rightarrow r}(A_1, \dots, A_n)) &= \text{Pos}_{\mathcal{F}}(\ell) \cup \{pq \mid p \in \text{Pos}_{\{x_i\}}(\ell) \text{ and } q \in \blacktriangle(A_i)\} \end{aligned}$$

where  $x \in \mathcal{V}$ ,  $f \in \mathcal{F}$ , and  $\text{var}(\ell) = (x_1, \dots, x_n)$ . So  $\blacktriangle(A)$  consists of all function positions of the redex patterns contracted in the multi-step of  $A$ . We write  $\blacktriangle(A, B)$  for  $\blacktriangle(A) \cap \blacktriangle(B)$ . The set  $\blacktriangle(A, B)$  is used below as a measure for the amount of overlap between the proof terms  $A$  and  $B$ . A pair  $(\Delta_1, \Delta_2)$  of co-initial redexes is an *overlap* if  $\blacktriangle(\Delta_1, \Delta_2) \neq \emptyset$ .

**Lemma 2.1.** *Let  $A$  and  $B$  be proof terms of left-linear TRSs  $\mathcal{R}$  and  $\mathcal{S}$  respectively. If  $\blacktriangle(A, B) \neq \emptyset$  then there exist a trivial or critical peak  $t \xleftarrow{\frac{\Delta_1}{\mathcal{R}}} s \xrightarrow{\frac{\Delta_2}{\mathcal{S}}} u$ , a context  $\mathbb{C}$ , and a substitution  $\sigma$  such that  $\mathbb{C}[\Delta_1\sigma] \in A$  and  $\mathbb{C}[\Delta_2\sigma] \in B$ . If  $\blacktriangle(A, B) = \emptyset$  then  $\xleftarrow{\frac{A}{\mathcal{R}}} \cdot \xrightarrow{\frac{B}{\mathcal{S}}} \subseteq \xrightarrow{\frac{B \setminus A}{\mathcal{S}}} \cdot \xleftarrow{\frac{A \setminus B}{\mathcal{R}}}$ .  $\square$*

### 3 Commutation by Relative Termination

Let  $\mathcal{R}$  and  $\mathcal{S}$  be TRSs. The set of critical peak steps of  $\mathcal{S}$  for  $\mathcal{R}$  is defined as follows:


$$\text{CPS}_{\mathcal{R}}(\mathcal{S}) = \{s \rightarrow u \mid t \xleftarrow{\mathcal{R}} s \rightarrow_{\mathcal{S}} u \text{ is a critical peak}\}$$

**Lemma 3.1.** *Let  $\mathcal{R}$  and  $\mathcal{S}$  be left-linear TRSs. If  $t \xleftarrow{\mathcal{R}} s \rightarrow_{\mathcal{S}} u$  then  $t \rightarrow_{\mathcal{S}} s \cdot \xleftarrow{\mathcal{R}} u$  or  $t \xleftarrow{\mathcal{R}} \cdot \text{CPS}_{\mathcal{S}}(\mathcal{R}) \leftarrow s \rightarrow_{\text{CPS}_{\mathcal{R}}(\mathcal{S})} \cdot \rightarrow_{\mathcal{S}} u$ .  $\square$*

**Theorem 3.2.** *Left-linear locally commuting TRSs  $\mathcal{R}$  and  $\mathcal{S}$  commute if  $\text{CPS}_{\mathcal{S}}(\mathcal{R}) \cup \text{CPS}_{\mathcal{R}}(\mathcal{S})$  is relatively terminating over  $\mathcal{R} \cup \mathcal{S}$ .*

*Proof.* We use decreasing diagrams with the predecessor labeling. To this end we define  $t \rightarrow_{\mathcal{R}, \mathcal{S}} u$  if and only if  $s \xrightarrow{*}_{\mathcal{R} \cup \mathcal{S}} t \rightarrow_{\mathcal{R}} u$ . The labeled relation  $\rightarrow_{\mathcal{S}, \mathcal{S}}$  is defined similarly. We write  $>$  to denote  $\rightarrow_{(\text{CPS}_{\mathcal{R}}(\mathcal{S}) \cup \text{CPS}_{\mathcal{S}}(\mathcal{R})) / (\mathcal{R} \cup \mathcal{S})}$ . By the relative termination condition,  $>$  is a well-founded order. We show that  $(\{\rightarrow_{\mathcal{R}, \mathcal{S}}\}_{s \in \mathcal{T}}, \{\rightarrow_{\mathcal{S}, \mathcal{S}}\}_{s \in \mathcal{T}})$  is decreasing with respect to  $>$ . Suppose  $t \xleftarrow{\mathcal{R}, s_1} s \rightarrow_{\mathcal{S}, s_2} u$ . By Lemma 3.1 and local commutation there are terms  $v$  and  $w$  such that (a) or (b) of Figure 1 holds. Note that  $s_i > v, w$  holds in case (b) because  $s_i \xrightarrow{*}_{\mathcal{R} \cup \mathcal{S}} s > v, w$ . Therefore, in both cases decreasingness is established. Hence  $\mathcal{R}$  and  $\mathcal{S}$  commute.  $\square$

Theorem 3.2 subsumes Rosen's commutativity criterion [4], as mutual orthogonality implies  $\text{CPS}_{\mathcal{S}}(\mathcal{R}) \cup \text{CPS}_{\mathcal{R}}(\mathcal{S}) = \emptyset$  as well as local commutation.

Figure 1: Decreasingness of  $(\dashrightarrow_{\mathcal{R}}, \dashrightarrow_{\mathcal{S}})$ .

## 4 Incorporating Development Closedness

We extend the last theorem in order to subsume the development closedness theorem of van Oostrom [7]. Let  $t \xrightarrow{\mathcal{R}}^p s \xrightarrow{q}_{\mathcal{S}} u$  be a critical peak. By definition  $p$  or  $q$  must be the root position. We say that the peak is  $(\mathcal{R}, \mathcal{S})$ -closed if  $t \xrightarrow{\mathcal{R}} u$  whenever  $p = \epsilon$  and  $t \dashrightarrow_{\mathcal{S}} u$  whenever  $q = \epsilon$ . The set of all non-closed critical peak steps of  $\mathcal{S}$  for  $\mathcal{R}$  is defined as follows:

$$\text{CPS}'_{\mathcal{R}}(\mathcal{S}) = \{s \rightarrow u \mid t \xrightarrow{\mathcal{R}} s \rightarrow_{\mathcal{S}} u \text{ is a critical peak which is not } (\mathcal{R}, \mathcal{S})\text{-closed}\}$$

**Lemma 4.1.** *Let  $\mathcal{R}$  and  $\mathcal{S}$  be left-linear TRSs. If  $t \xrightarrow{\mathcal{R}} s \dashrightarrow_{\mathcal{S}} u$  then (a)  $t \dashrightarrow_{\mathcal{S}} \cdot \xrightarrow{\mathcal{R}} u$  or (b)  $t \xrightarrow{\mathcal{R}} \cdot \text{CPS}'_{\mathcal{S}}(\mathcal{R}) \dashrightarrow s' \rightarrow_{\text{CPS}'_{\mathcal{R}}(\mathcal{S})} \cdot \dashrightarrow_{\mathcal{S}} u$  and  $s \rightarrow_{\mathcal{R} \cup \mathcal{S}}^* s'$  for some  $s'$ .*

Lemma 4.1 is a proper generalization of [2, Lemma 4]. In order to prove the lemma we introduce some more notions. Whenever  $\blacktriangle(A, B)$  is non-empty, there exists an innermost overlap. An overlap  $(\Delta_1, \Delta_2)$  in  $A \times B$  is called *innermost* if there are no other overlaps  $(\Delta'_1, \Delta'_2)$  in  $A \times B$  such that  $\Delta'_1 \geq \Delta_1$  and  $\Delta'_2 \geq \Delta_2$ . We say that  $(\Delta_1, \Delta_2)$  is  $(\mathcal{R}, \mathcal{S})$ -closed if the corresponding critical peak is  $(\mathcal{R}, \mathcal{S})$ -closed. The following is the key lemma. In order to pave the way for formalisation of advanced confluence and commutation results, we formalise the proof technique introduced in [7] using proof terms.

**Lemma 4.2.** *Suppose  $(\Delta_1, \Delta_2)$  is an innermost overlap in  $A \times B$  with  $\Delta_1 \geq \Delta_2$ . If  $(\Delta_1, \Delta_2)$  is  $(\mathcal{R}, \mathcal{S})$ -closed then  $\blacktriangle(A, B) \supseteq \blacktriangle(A \setminus \Delta_1, C)$  for some proof term  $C$  that is co-initial with  $A \setminus \Delta_1$ .*

*Proof.* Write  $t \xrightarrow[A \setminus \Delta_1]{\mathcal{R}} t_1 \xrightarrow[\mathcal{R}]{\Delta_1} s \xrightarrow[\mathcal{S}]{\Delta_2} u_1 \xrightarrow[B \setminus \Delta_2]{\mathcal{S}} u$  and let

$$t'_1 \xrightarrow[\mathcal{R}]{\Delta'_1} s' \xrightarrow[\mathcal{S}]{\Delta'_2} u'_1$$

be the corresponding critical peak. Since  $s \geq s'$ , there exist a context  $\mathbb{C}$  and a substitution  $\sigma$  such that  $s = \mathbb{C}[s'\sigma]$ ,  $t_1 = \mathbb{C}[t'_1\sigma]$ ,  $u_1 = \mathbb{C}[u'_1\sigma]$ ,  $\Delta_1 = \mathbb{C}[\Delta'_1\sigma]$ , and  $\Delta_2 = \mathbb{C}[\Delta'_2\sigma]$ . We have  $t'_1 \dashrightarrow_{\mathcal{R}} u'_1$  because the critical peak is  $(\mathcal{R}, \mathcal{S})$ -closed. Let  $D'$  be the proof term witnessing this multi-step and define  $D = \mathbb{C}[D'\sigma]$ . Clearly,  $D$  is a witness of  $t_1 \dashrightarrow_{\mathcal{R}} u_1$ . We will compose  $D$  and  $B \setminus \Delta_2$  into a single proof term  $C: t_1 \dashrightarrow_{\mathcal{R}} u$ . Let  $B' = B - \Delta_2$ . Since  $(\Delta_1, \Delta_2)$  is an innermost overlap with  $\Delta_1 \geq \Delta_2$ ,  $\blacktriangle(B', \Delta_1) = \emptyset$ . Hence  $B' \setminus \Delta_1$  is well-defined. If we show that  $\blacktriangle(B' \setminus \Delta_1, D) = \emptyset$  then we can define  $C$  as  $(B' \setminus \Delta_1) \sqcup D$ . We have  $\blacktriangle(D') \subseteq \text{Pos}_{\mathcal{F}}(t'_1)$  and thus  $\blacktriangle(D) \subseteq \{qp \mid p \in \text{Pos}_{\mathcal{F}}(t'_1)\}$  where  $q$  is the position of the hole in  $\mathbb{C}$ . In words,  $D$  affects only the  $t'_1$  part of  $\mathbb{C}[t'_1\sigma]$ . Let  $\Delta \in B'$  be an arbitrary redex. We show that  $\blacktriangle(\Delta \setminus \Delta_1, D) = \emptyset$ . We split  $B'$  into two parts:  $B' = B_1 \sqcup B_2$  with

$$B_1 = B' - B_2 \quad B_2 = \bigsqcup \{\Delta \mid \Delta \in B' \text{ and } \Delta > \Delta_2\}$$

Depending on the position of  $\Delta$ , we distinguish two cases:

- (a) If  $\Delta \in B_1$  then  $\Delta$  does not overlap  $\Delta_1$ . So  $\blacktriangle(\Delta \setminus \Delta_1)$  consists entirely of positions in  $\mathbb{C}$  and thus  $\blacktriangle(\Delta \setminus \Delta_1, D) = \emptyset$ .
- (b) Otherwise,  $\Delta \in B_2$ . Since  $(\Delta_1, \Delta_2)$  is an innermost overlap,  $\blacktriangle(\Delta, \Delta_1) = \emptyset$ . Because  $\Delta \neq \Delta_2$  we also have  $\blacktriangle(\Delta, \Delta_2) = \emptyset$ . The crucial observation is that no redex in  $\Delta \setminus \Delta_1$  overlaps with  $D$ . This follows because for the originating term  $s'$  of the critical peak we have  $\text{Pos}_{\mathcal{F}}(s') = \blacktriangle(\Delta'_1) \cup \blacktriangle(\Delta'_2)$ , due to the fact that the involved rules are left-linear. Hence all redexes in  $\Delta \setminus \Delta_1$  appear in the substitution part  $\sigma$  of  $\mathbb{C}[t'_1\sigma]$ .

We conclude that  $\blacktriangle(B' \setminus \Delta_1, D) = \emptyset$ . It remains to show that  $\blacktriangle(A, B) \supseteq \blacktriangle(A \setminus \Delta_1, C)$ . We have  $\blacktriangle(B_1 \setminus \Delta_1) = \blacktriangle(B_1)$  and hence  $\blacktriangle(C) = \blacktriangle(B_1) \cup \blacktriangle(B_2 \setminus \Delta_1) \cup \blacktriangle(D)$ . Because  $(\Delta_1, \Delta_2)$  is an innermost overlap,  $\blacktriangle(A, B_2) = \emptyset$  and therefore

$$\blacktriangle(A, B) = \blacktriangle(A, \Delta_2) \cup \blacktriangle(A, B_1)$$

and  $\blacktriangle(A \setminus \Delta_1, B_2 \setminus \Delta_1) = \emptyset$ . Because all redexes in  $B_1$  occur above  $\Delta_2$ , they do not overlap with  $\Delta_1$  and thus  $\blacktriangle(A \setminus \Delta_1, B \setminus \Delta_1) = \blacktriangle(A \setminus \Delta_1, B_1 \setminus \Delta_1) = \blacktriangle(A, B_1)$ . We obtain

$$\begin{aligned} \blacktriangle(A \setminus \Delta_1, C) &= \blacktriangle(A \setminus \Delta_1, B' \setminus \Delta_1) \cup \blacktriangle(A \setminus \Delta_1, D) \\ &= \blacktriangle(A, B_1) \cup \blacktriangle(A \setminus \Delta_1, D) \end{aligned}$$

and so it remains to show that  $\blacktriangle(A, \Delta_2) \supseteq \blacktriangle(A \setminus \Delta_1, D)$ . We split  $A$  into three parts:  $A = A_1 \sqcup \Delta_1 \sqcup A_2$  with  $A_1 = A - \Delta_1 - A_2$  and  $A_2 = \bigsqcup\{\Delta \mid \Delta \in A \text{ and } \Delta > \Delta_1\}$ . We have  $\blacktriangle(A, \Delta_2) = \blacktriangle(A_1, \Delta_2) \cup \blacktriangle(\Delta_1, \Delta_2)$  and

$$\blacktriangle(A \setminus \Delta_1, D) = \blacktriangle(A_1 \setminus \Delta_1, D) \cup \blacktriangle(\Delta_1 \setminus \Delta_1, D) \cup \blacktriangle(A_2 \setminus \Delta_1, D)$$

The second component is clearly empty but also the third component reduces to the empty set, by repeating the reasoning in case (b) above. The first component equals  $\blacktriangle(A_1, D)$  since redexes in  $A_1$  occur above  $\Delta_1$ . So  $\blacktriangle(A \setminus \Delta_1, D) = \blacktriangle(A_1, D)$ . Since  $\blacktriangle(\Delta_1, \Delta_2) \neq \emptyset$ , it suffices to show the inclusion  $\blacktriangle(A_1, D) \subseteq \blacktriangle(A_1, \Delta_2)$ , which follows from the observation that overlaps between  $A_1$  and  $D$  are restricted to positions in  $\blacktriangle(\Delta_2) - \blacktriangle(\Delta_1)$ .  $\square$

We are ready to show the aforementioned lemma.

*Proof of Lemma 4.1.* By induction on the finite set  $\blacktriangle(A, B)$  with respect to the well-founded order  $\supseteq$ . Let

$$t \xleftarrow[\mathcal{R}]{A} s \xrightarrow[\mathcal{S}]{B} u$$

If  $\blacktriangle(A, B)$  is empty then (a) holds by the second part of Lemma 2.1. Otherwise, there is an overlap in  $A \times B$ . We distinguish two cases.

- If a non-closed overlap  $(\Delta_1, \Delta_2)$  exists in  $A \times B$ , (b) holds because

$$t \xleftarrow[\mathcal{R}]{A \setminus \Delta_1} \cdot \xleftarrow[\mathcal{R}]{\Delta_1} s \xrightarrow[\mathcal{S}]{\Delta_2} \cdot \xrightarrow[\mathcal{S}]{B \setminus \Delta_2} u$$

and the two inner steps correspond to applications of rules from  $\text{CPS}'_{\mathcal{S}}(\mathcal{R})$  and  $\text{CPS}'_{\mathcal{R}}(\mathcal{S})$ .

- In the other case all overlaps are closed. Let  $(\Delta_1, \Delta_2)$  be an innermost overlap. Without loss of generality we assume  $\Delta_1 \geq \Delta_2$ . According to Lemma 4.2 there exists some proof


term  $C$  such that  $\blacktriangle(A, B) \supseteq \blacktriangle(A \setminus \Delta_1, C)$ . Let  $s'$  be the term such that  $s \xrightarrow{\Delta_1} s'$ . (In the proof of Lemma 4.2  $s'$  was called  $t_1$ .) Because

$$t \xleftarrow[\mathcal{R}]{A \setminus \Delta_1} s' \xrightarrow[\mathcal{S}]{C} u$$

the induction hypothesis applies.  $\square$

**Theorem 4.3.** *Left-linear locally commuting TRSs  $\mathcal{R}$  and  $\mathcal{S}$  commute if  $\text{CPS}'_{\mathcal{S}}(\mathcal{R}) \cup \text{CPS}'_{\mathcal{R}}(\mathcal{S})$  is relatively terminating over  $\mathcal{R} \cup \mathcal{S}$ .*

*Proof.* Use Lemma 4.1 to replace Figure 1(b) by the following diagram:



$\square$

Due to lack of space, we omit the extension that weakens the joinability requirement for overlays, but note that the result of Theorem 4.3 remains true if we strengthen the notion of  $(\mathcal{R}, \mathcal{S})$ -closedness by allowing  $t \xrightarrow{\mathcal{S}}^* \cdot \mathcal{R} \leftarrow \ominus u$  when  $p = \epsilon$ . Future work includes the extension to critical valleys [9] as well as automation. Experimental data on the confluence problem database Cops<sup>1</sup> for the presented theorems will be reported at the workshop.

## References

- [1] T. Aoto, J. Yoshida, and Y. Toyama. Proving confluence of term rewriting systems automatically. In *Proc. 21st RTA*, volume 5595 of *LNCS*, pages 93–102, 2009.
- [2] N. Hirokawa and A. Middeldorp. Decreasing diagrams and relative termination. *Journal of Automated Reasoning*, 47(4):481–501, 2011.
- [3] J.-C. Raoult and J. Vuillemin. Operational and semantic equivalence between recursive programs. *Journal of the ACM*, 27(4):772–796, 1980.
- [4] B.K. Rosen. Tree-manipulating systems and Church-Rosser theorems. *Journal of the ACM*, 20(1):160–187, 1973.
- [5] TeReSe. *Term Rewriting Systems*, volume 55 of *Cambridge Tracts in Theoretical Computer Science*. Cambridge University Press, 2003.
- [6] Y. Toyama. Commutativity of term rewriting systems. In *Programming of Future Generation Computers II*, pages 393–407. North-Holland, 1988.
- [7] V. van Oostrom. Developing developments. *Theoretical Computer Science*, 175(1):159–181, 1997.
- [8] V. van Oostrom. Confluence by decreasing diagrams – converted. In *Proc. 19th RTA*, volume 5117 of *LNCS*, pages 306–320, 2008.
- [9] V. van Oostrom. Confluence via critical valleys. In *Proc. 6th HOR*, pages 9–10, 2012.

<sup>1</sup><http://coco.nue.riec.tohoku.ac.jp/>