



Logic

Luca Campa

Philipp Dablander

Aaron Groß

Aart Middeldorp

Alexander Montag

Johannes Niederhauser

Vera Schmitt

 **articify**
ars.uibk.ac.at

with session ID **6893 6178** for anonymous questions



Outline

- 1. Summary of Previous Lecture**
- 2. Natural Deduction**
- 3. Intermezzo**
- 4. Soundness**
- 5. Further Reading**

Definition

- ▶ **Horn clause** is propositional formula

$$P_1 \wedge P_2 \wedge \dots \wedge P_n \rightarrow Q$$

with $n \geq 1$ and where P_1, \dots, P_n, Q are atoms, \perp or \top

- ▶ **Horn formula** is conjunction of Horn clauses

Theorem

satisfiability of Horn formulas is **efficiently** decidable

Remark

deciding satisfiability for **arbitrary** formulas is important and difficult problem (**SAT**)

Definition

formulas φ and ψ are **equisatisfiable** ($\varphi \approx \psi$) if

$$\varphi \text{ is satisfiable} \iff \psi \text{ is satisfiable}$$

Remark

Tseitin's transformation transforms arbitrary formula into equisatisfiable CNF in linear time

Lemma

- 1 any satisfying valuation for φ can be (uniquely) extended to satisfying valuation for $\text{TT}(\varphi)$
- 2 restriction of any satisfying valuation for $\text{TT}(\varphi)$ to atoms in φ is satisfying valuation for φ

Part I: Propositional Logic

algebraic normal forms, binary decision diagrams, conjunctive normal forms, DPLL, Horn formulas, natural deduction, Post's adequacy theorem, resolution, SAT, semantics, sorting networks, soundness and completeness, syntax, Tseitin's transformation

Part II: Predicate Logic

natural deduction, quantifier equivalences, resolution, semantics, Skolemization, syntax, undecidability, unification

Part III: Model Checking

adequacy, branching-time temporal logic, CTL*, fairness, linear-time temporal logic, model checking algorithms, symbolic model checking

Part I: Propositional Logic

algebraic normal forms, binary decision diagrams, conjunctive normal forms, DPLL, Horn formulas, **natural deduction**, Post's adequacy theorem, resolution, SAT, semantics, sorting networks, **soundness** and completeness, syntax, Tseitin's transformation

Part II: Predicate Logic

natural deduction, quantifier equivalences, resolution, semantics, Skolemization, syntax, undecidability, unification

Part III: Model Checking

adequacy, branching-time temporal logic, CTL*, fairness, linear-time temporal logic, model checking algorithms, symbolic model checking

Outline

1. Summary of Previous Lecture
- 2. Natural Deduction**
3. Intermezzo
4. Soundness
5. Further Reading

Natural Deduction

calculus for reasoning about propositions

Natural Deduction

calculus for reasoning about propositions

Definitions

► **sequent**

$$\underbrace{\varphi_1, \varphi_2, \dots, \varphi_n}_{\text{premises}} \vdash \underbrace{\psi}_{\text{conclusion}}$$

with propositional formulas $\varphi_1, \varphi_2, \dots, \varphi_n, \psi$

Natural Deduction

calculus for reasoning about propositions

Definitions

- ▶ sequent

$$\underbrace{\varphi_1, \varphi_2, \dots, \varphi_n}_{\text{premises}} \quad \vdash \quad \underbrace{\psi}_{\text{conclusion}}$$

with propositional formulas $\varphi_1, \varphi_2, \dots, \varphi_n, \psi$

- ▶ sequent $\varphi_1, \varphi_2, \dots, \varphi_n \vdash \psi$ is **valid** if ψ can be proved from premises $\varphi_1, \varphi_2, \dots, \varphi_n$ using **proof rules** of natural deduction

Natural Deduction

calculus for reasoning about propositions

Definitions

- ▶ sequent

$$\underbrace{\varphi_1, \varphi_2, \dots, \varphi_n}_{\text{premises}} \quad \vdash \quad \underbrace{\psi}_{\text{conclusion}}$$

with propositional formulas $\varphi_1, \varphi_2, \dots, \varphi_n, \psi$

- ▶ sequent $\varphi_1, \varphi_2, \dots, \varphi_n \vdash \psi$ is valid if ψ can be proved from premises $\varphi_1, \varphi_2, \dots, \varphi_n$ using proof rules of natural deduction

natural deduction consists of 17 proof rules

Definition

► and introduction

$$\frac{\varphi \quad \psi}{\varphi \wedge \psi} \wedge i$$

Definition

- ▶ and introduction

$$\frac{\varphi \quad \psi}{\varphi \wedge \psi} \wedge i$$

Example

$p, q, r \vdash (r \wedge q) \wedge p$ is valid

Definition

- ▶ and introduction

$$\frac{\varphi \quad \psi}{\varphi \wedge \psi} \wedge i$$

Example

$p, q, r \vdash (r \wedge q) \wedge p$ is valid: 1 p premise

Definition

- ▶ and introduction

$$\frac{\varphi \quad \psi}{\varphi \wedge \psi} \wedge i$$

Example

$p, q, r \vdash (r \wedge q) \wedge p$ is valid:

1	p	premise
2	q	premise

Definition

- ▶ and introduction

$$\frac{\varphi \quad \psi}{\varphi \wedge \psi} \wedge i$$

Example

$p, q, r \vdash (r \wedge q) \wedge p$ is valid:	1	p	premise
	2	q	premise
	3	r	premise

Definition

- ▶ and introduction

$$\frac{\varphi \quad \psi}{\varphi \wedge \psi} \wedge i$$

Example

$p, q, r \vdash (r \wedge q) \wedge p$ is valid:	1	p	premise
	2	q	premise
	3	r	premise
	4	$r \wedge q$	$\wedge i$ 3,2

Definition

- ▶ and introduction

$$\frac{\varphi \quad \psi}{\varphi \wedge \psi} \wedge i$$

Example

$p, q, r \vdash (r \wedge q) \wedge p$ is valid:	1	p	premise
	2	q	premise
	3	r	premise
	4	$r \wedge q$	$\wedge i$ 3,2

Definition

- ▶ and introduction

$$\frac{\varphi \quad \psi}{\varphi \wedge \psi} \wedge i$$

Example

$p, q, r \vdash (r \wedge q) \wedge p$ is valid:	1	p	premise
	2	q	premise
	3	r	premise
	4	$r \wedge q$	$\wedge i$ 3, 2

Definition

- ▶ and introduction

$$\frac{\varphi \quad \psi}{\varphi \wedge \psi} \wedge i$$

Example

$p, q, r \vdash (r \wedge q) \wedge p$ is valid:	1	p	premise
	2	q	premise
	3	r	premise
	4	$r \wedge q$	$\wedge i$ 3,2
	5	$(r \wedge q) \wedge p$	$\wedge i$ 4,1

Definition

► and elimination

$$\frac{\varphi \wedge \psi}{\varphi} \wedge e_1$$

$$\frac{\varphi \wedge \psi}{\psi} \wedge e_2$$

Definition

► and elimination

$$\frac{\varphi \wedge \psi}{\varphi} \wedge e_1$$

$$\frac{\varphi \wedge \psi}{\psi} \wedge e_2$$

Example

$p \wedge q, r \vdash r \wedge q$ is valid:

- 1 $p \wedge q$ premise
- 2 r premise

Definition

► and elimination

$$\frac{\varphi \wedge \psi}{\varphi} \wedge e_1$$

$$\frac{\varphi \wedge \psi}{\psi} \wedge e_2$$

Example

$p \wedge q, r \vdash r \wedge q$ is valid:

- 1 $p \wedge q$ premise
- 2 r premise
- 3 q $\wedge e_2$ 1

Definition

► and elimination

$$\frac{\varphi \wedge \psi}{\varphi} \wedge e_1$$

$$\frac{\varphi \wedge \psi}{\psi} \wedge e_2$$

Example

$p \wedge q, r \vdash r \wedge q$ is valid:

- 1 $p \wedge q$ premise
- 2 r premise
- 3 q $\wedge e_2$ 1
- 4 $r \wedge q$ $\wedge i$ 2, 3

Definitions

- ▶ double negation elimination

$$\frac{\neg\neg\varphi}{\varphi} \quad \neg\neg e$$

Definitions

▶ double negation elimination

$$\frac{\neg\neg\varphi}{\varphi} \quad \neg\neg e$$

▶ double negation introduction

$$\frac{\varphi}{\neg\neg\varphi} \quad \neg\neg i$$

Definitions

- ▶ double negation elimination
$$\frac{\neg\neg\varphi}{\varphi} \neg\neg e$$
- ▶ double negation introduction
$$\frac{\varphi}{\neg\neg\varphi} \neg\neg i$$

Example

$p, \neg\neg(q \wedge r) \vdash \neg\neg p \wedge r$ is valid:

- 1 p premise
- 2 $\neg\neg(q \wedge r)$ premise

Definitions

- ▶ double negation elimination
$$\frac{\neg\neg\varphi}{\varphi} \neg\neg e$$
- ▶ double negation introduction
$$\frac{\varphi}{\neg\neg\varphi} \neg\neg i$$

Example

$p, \neg\neg(q \wedge r) \vdash \neg\neg p \wedge r$ is valid:

- 1 p premise
- 2 $\neg\neg(q \wedge r)$ premise

- 6 $\neg\neg p \wedge r$

Definitions

▶ double negation elimination $\frac{\neg\neg\varphi}{\varphi} \neg\neg e$

▶ double negation introduction $\frac{\varphi}{\neg\neg\varphi} \neg\neg i$

Example

$p, \neg\neg(q \wedge r) \vdash \neg\neg p \wedge r$ is valid:

1	p	premise
2	$\neg\neg(q \wedge r)$	premise
3	$\neg\neg p$	
5	r	
6	$\neg\neg p \wedge r$	$\wedge i$ 3,5

Definitions

▶ double negation elimination $\frac{\neg\neg\varphi}{\varphi} \neg\neg e$

▶ double negation introduction $\frac{\varphi}{\neg\neg\varphi} \neg\neg i$

Example

$p, \neg\neg(q \wedge r) \vdash \neg\neg p \wedge r$ is valid:

1	p	premise
2	$\neg\neg(q \wedge r)$	premise
3	$\neg\neg p$	$\neg\neg i$ 1
5	r	
6	$\neg\neg p \wedge r$	$\wedge i$ 3,5

Definitions

▶ double negation elimination

$$\frac{\neg\neg\varphi}{\varphi} \neg\neg e$$

▶ double negation introduction

$$\frac{\varphi}{\neg\neg\varphi} \neg\neg i$$

Example

$p, \neg\neg(q \wedge r) \vdash \neg\neg p \wedge r$ is valid:

1	p	premise
2	$\neg\neg(q \wedge r)$	premise
3	$\neg\neg p$	$\neg\neg i$ 1
4	$q \wedge r$	$\neg\neg e$ 2
5	r	
6	$\neg\neg p \wedge r$	$\wedge i$ 3,5

Definitions

▶ double negation elimination

$$\frac{\neg\neg\varphi}{\varphi} \neg\neg e$$

▶ double negation introduction

$$\frac{\varphi}{\neg\neg\varphi} \neg\neg i$$

Example

$p, \neg\neg(q \wedge r) \vdash \neg\neg p \wedge r$ is valid:

1	p	premise
2	$\neg\neg(q \wedge r)$	premise
3	$\neg\neg p$	$\neg\neg i$ 1
4	$q \wedge r$	$\neg\neg e$ 2
5	r	$\wedge e_2$ 4
6	$\neg\neg p \wedge r$	$\wedge i$ 3,5

Definition

► implication elimination

$$\frac{\varphi \rightarrow \psi \quad \varphi}{\psi} \rightarrow e$$

Definition

- ▶ implication elimination (**modus ponens**)

$$\frac{\varphi \rightarrow \psi \quad \varphi}{\psi} \rightarrow e$$

Definition

- ▶ implication elimination (modus ponens)

$$\frac{\varphi \rightarrow \psi \quad \varphi}{\psi} \rightarrow e$$

Example

$p, p \rightarrow q, p \rightarrow (q \rightarrow r) \vdash r$ is valid:

1	p	premise
2	$p \rightarrow q$	premise
3	$p \rightarrow (q \rightarrow r)$	premise

Definition

- ▶ implication elimination (modus ponens)

$$\frac{\varphi \rightarrow \psi \quad \varphi}{\psi} \rightarrow e$$

Example

$p, p \rightarrow q, p \rightarrow (q \rightarrow r) \vdash r$ is valid:

1	p	premise
2	$p \rightarrow q$	premise
3	$p \rightarrow (q \rightarrow r)$	premise
4	q	$\rightarrow e$ 2, 1

Definition

- ▶ implication elimination (modus ponens)

$$\frac{\varphi \rightarrow \psi \quad \varphi}{\psi} \rightarrow e$$

Example

$p, p \rightarrow q, p \rightarrow (q \rightarrow r) \vdash r$ is valid:

1	p	premise
2	$p \rightarrow q$	premise
3	$p \rightarrow (q \rightarrow r)$	premise
4	q	$\rightarrow e$ 2, 1
5	$q \rightarrow r$	$\rightarrow e$ 3, 1

Definition

- ▶ implication elimination (modus ponens)

$$\frac{\varphi \rightarrow \psi \quad \varphi}{\psi} \rightarrow e$$

Example

$p, p \rightarrow q, p \rightarrow (q \rightarrow r) \vdash r$ is valid:

1	p	premise
2	$p \rightarrow q$	premise
3	$p \rightarrow (q \rightarrow r)$	premise
4	q	$\rightarrow e$ 2, 1
5	$q \rightarrow r$	$\rightarrow e$ 3, 1
6	r	$\rightarrow e$ 5, 4

Definition

► modus tollens

$$\frac{\varphi \rightarrow \psi \quad \neg \psi}{\neg \varphi} \quad \text{MT}$$

Definition

► modus tollens

$$\frac{\varphi \rightarrow \psi \quad \neg \psi}{\neg \varphi} \text{ MT}$$

Example

$\neg p \rightarrow q, \neg q \vdash p$ is valid:	1	$\neg p \rightarrow q$	premise
	2	$\neg q$	premise
	3	$\neg \neg p$	MT 1, 2
	4	p	$\neg \neg$ e 3

Definition

- implication introduction

$$\frac{\begin{array}{|c} \varphi \\ \vdots \\ \psi \end{array}}{\varphi \rightarrow \psi} \rightarrow i$$

Definition

► implication introduction

$$\text{box} \frac{\begin{array}{c} \varphi \\ \vdots \\ \psi \end{array}}{\varphi \rightarrow \psi} \rightarrow i$$

Definition

- implication introduction

$$\frac{\begin{array}{|l} \varphi \\ \vdots \\ \psi \end{array}}{\varphi \rightarrow \psi} \rightarrow i$$

assumption

Definition

- ▶ implication introduction

$$\frac{\begin{array}{|c|} \hline \varphi \\ \vdots \\ \psi \\ \hline \end{array}}{\varphi \rightarrow \psi} \rightarrow i$$

Example

$\neg q \rightarrow \neg p \vdash p \rightarrow q$ is valid: 1 $\neg q \rightarrow \neg p$ premise

6 $p \rightarrow q$

Definition

- implication introduction

$$\frac{\begin{array}{|c|} \hline \varphi \\ \vdots \\ \psi \\ \hline \end{array}}{\varphi \rightarrow \psi} \rightarrow i$$

Example

$\neg q \rightarrow \neg p \vdash p \rightarrow q$ is valid:

1 $\neg q \rightarrow \neg p$ premise

2 p assumption

6 $p \rightarrow q$

Definition

- ▶ implication introduction

$$\frac{\begin{array}{|c|} \hline \varphi \\ \vdots \\ \psi \\ \hline \end{array}}{\varphi \rightarrow \psi} \rightarrow i$$

Example

$\neg q \rightarrow \neg p \vdash p \rightarrow q$ is valid:

1 $\neg q \rightarrow \neg p$ premise

2 p assumption

5 q

6 $p \rightarrow q$

Definition

- implication introduction

$$\frac{\begin{array}{|c|} \hline \varphi \\ \vdots \\ \psi \\ \hline \end{array}}{\varphi \rightarrow \psi} \rightarrow i$$

Example

$\neg q \rightarrow \neg p \vdash p \rightarrow q$ is valid:

1 $\neg q \rightarrow \neg p$ premise

2 p assumption

3 $\neg\neg p$ $\neg\neg i$ 2

5 q

6 $p \rightarrow q$

Definition

- implication introduction

$$\frac{\begin{array}{|c|} \hline \varphi \\ \vdots \\ \psi \\ \hline \end{array}}{\varphi \rightarrow \psi} \rightarrow i$$

Example

$\neg q \rightarrow \neg p \vdash p \rightarrow q$ is valid:

1	$\neg q \rightarrow \neg p$	premise
2	p	assumption
3	$\neg \neg p$	$\neg \neg i$ 2
4	$\neg \neg q$	MT 1, 3
5	q	$\neg \neg e$ 4
6	$p \rightarrow q$	

Definition

- implication introduction

$$\frac{\begin{array}{|c|} \hline \varphi \\ \vdots \\ \psi \\ \hline \end{array}}{\varphi \rightarrow \psi} \rightarrow i$$

Example

$\neg q \rightarrow \neg p \vdash p \rightarrow q$ is valid:

1	$\neg q \rightarrow \neg p$	premise
2	p	assumption
3	$\neg \neg p$	$\neg \neg i$ 2
4	$\neg \neg q$	MT 1, 3
5	q	$\neg \neg e$ 4
6	$p \rightarrow q$	$\rightarrow i$ 2-5

Definition

► or introduction

$$\frac{\varphi}{\varphi \vee \psi} \quad \vee i_1$$

$$\frac{\psi}{\varphi \vee \psi} \quad \vee i_2$$

Definition

► or introduction

$$\frac{\varphi}{\varphi \vee \psi} \vee i_1$$

$$\frac{\psi}{\varphi \vee \psi} \vee i_2$$

Example

$p \wedge q \vdash \neg q \vee p$ is valid:

1 $p \wedge q$ premise

2 p $\wedge e_1$ 1

3 $\neg q \vee p$ $\vee i_2$ 2

Definition

► or elimination

$$\frac{\varphi \vee \psi \quad \begin{array}{|c|} \hline \varphi \\ \hline \vdots \\ \hline \chi \\ \hline \end{array} \quad \begin{array}{|c|} \hline \psi \\ \hline \vdots \\ \hline \chi \\ \hline \end{array}}{\chi} \vee e$$

Definition

► or elimination

$$\frac{\varphi \vee \psi \quad \begin{array}{|c|} \hline \varphi \\ \vdots \\ \chi \\ \hline \end{array} \quad \begin{array}{|c|} \hline \psi \\ \vdots \\ \chi \\ \hline \end{array}}{\chi} \vee e$$

Example

$p \vee q \vdash q \vee p$ is valid: 1 $p \vee q$ premise

Definition

► or elimination

$$\frac{\varphi \vee \psi \quad \begin{array}{|c|} \hline \varphi \\ \hline \vdots \\ \hline \chi \\ \hline \end{array} \quad \begin{array}{|c|} \hline \psi \\ \hline \vdots \\ \hline \chi \\ \hline \end{array}}{\chi} \vee e$$

Example

$p \vee q \vdash q \vee p$ is valid:

1 $p \vee q$ premise

2 p assumption

3 $q \vee p$ $\vee i_2$ 2

Definition

► or elimination

$$\frac{\varphi \vee \psi \quad \begin{array}{|c|} \hline \varphi \\ \vdots \\ \chi \\ \hline \end{array} \quad \begin{array}{|c|} \hline \psi \\ \vdots \\ \chi \\ \hline \end{array}}{\chi} \vee e$$

Example

$p \vee q \vdash q \vee p$ is valid:

1 $p \vee q$ premise

2 p assumption

3 $q \vee p$ $\vee i_2$ 2

4 q assumption

5 $q \vee p$ $\vee i_1$ 4

Definition

► or elimination

$$\frac{\varphi \vee \psi \quad \begin{array}{|c|} \hline \varphi \\ \vdots \\ \chi \\ \hline \end{array} \quad \begin{array}{|c|} \hline \psi \\ \vdots \\ \chi \\ \hline \end{array}}{\chi} \vee e$$

Example

$p \vee q \vdash q \vee p$ is valid:

1 $p \vee q$ premise

2 p assumption

3 $q \vee p$ $\vee i_2$ 2

4 q assumption

5 $q \vee p$ $\vee i_1$ 4

6 $q \vee p$ $\vee e$ 1, 2–3, 4–5

Definition

theorem is formula φ such that sequent $\vdash \varphi$ is valid

Definition

theorem is formula φ such that sequent $\vdash \varphi$ is valid

Example

$p \vee q \rightarrow q \vee p$ is theorem

Definition

theorem is formula φ such that sequent $\vdash \varphi$ is valid

Example

$p \vee q \rightarrow q \vee p$ is theorem:

1	$p \vee q$	assumption
2	p	assumption
3	$q \vee p$	$\vee i_2$ 2
4	q	assumption
5	$q \vee p$	$\vee i_1$ 4
6	$q \vee p$	$\vee e$ 1, 2-3, 4-5

Definition

theorem is formula φ such that sequent $\vdash \varphi$ is valid

Example

$p \vee q \rightarrow q \vee p$ is theorem:

1	$p \vee q$	assumption
2	p	assumption
3	$q \vee p$	$\vee i_2$ 2
4	q	assumption
5	$q \vee p$	$\vee i_1$ 4
6	$q \vee p$	$\vee e$ 1, 2-3, 4-5
7	$p \vee q \rightarrow q \vee p$	$\rightarrow i$ 1-6

- ▶ bottom elimination

$$\frac{\perp}{\phi} \perp e$$

Definitions

- ▶ bottom elimination

$$\frac{\perp}{\varphi} \perp e$$

- ▶ negation elimination

$$\frac{\varphi \quad \neg\varphi}{\perp} \neg e$$

Definitions

- ▶ bottom elimination

$$\frac{\perp}{\varphi} \perp e$$

- ▶ negation elimination

$$\frac{\varphi \quad \neg\varphi}{\perp} \neg e$$

- ▶ negation introduction

$$\frac{\boxed{\begin{array}{c} \varphi \\ \vdots \\ \perp \end{array}}}{\neg\varphi} \neg i$$

Definitions

- ▶ bottom elimination

$$\frac{\perp}{\varphi} \perp e$$

- ▶ negation elimination

$$\frac{\varphi \quad \neg\varphi}{\perp} \neg e$$

- ▶ negation introduction

$$\frac{\boxed{\begin{array}{c} \varphi \\ \vdots \\ \perp \end{array}}}{\neg\varphi} \neg i$$

- ▶ top introduction

$$\frac{}{\top} \top i$$

Examples

$p \rightarrow q, p \rightarrow \neg q \vdash \neg p$ is valid:

1	$p \rightarrow q$	premise
2	$p \rightarrow \neg q$	premise
3	p	assumption
4	q	\rightarrow e 1, 3
5	$\neg q$	\rightarrow e 2, 3
6	\perp	\neg e 4, 5

Examples

$p \rightarrow q, p \rightarrow \neg q \vdash \neg p$ is valid:

1	$p \rightarrow q$	premise
2	$p \rightarrow \neg q$	premise
3	p	assumption
4	q	\rightarrow e 1, 3
5	$\neg q$	\rightarrow e 2, 3
6	\perp	\neg e 4, 5
7	$\neg p$	\neg i 3-6

Examples

$p \rightarrow q, p \rightarrow \neg q \vdash \neg p$ is valid:

1	$p \rightarrow q$	premise
2	$p \rightarrow \neg q$	premise
3	p	assumption
4	q	\rightarrow e 1, 3
5	$\neg q$	\rightarrow e 2, 3
6	\perp	\neg e 4, 5
7	$\neg p$	\neg i 3-6

$p, p \rightarrow q, p \rightarrow \neg q \vdash r$ is valid:

1	p	premise
2	$p \rightarrow q$	premise
3	$p \rightarrow \neg q$	premise
4	q	\rightarrow e 2, 1
5	$\neg q$	\rightarrow e 3, 1
6	\perp	\neg e 4, 5
7	r	\perp e 6

Examples

$p \rightarrow q, p \rightarrow \neg q \vdash \neg p$ is valid:

1	$p \rightarrow q$	premise
2	$p \rightarrow \neg q$	premise
3	p	assumption
4	q	\rightarrow e 1, 3
5	$\neg q$	\rightarrow e 2, 3
6	\perp	\neg e 4, 5
7	$\neg p$	\neg i 3-6

$p, p \rightarrow q, p \rightarrow \neg q \vdash r$ is valid:

1	p	premise
2	$p \rightarrow q$	premise
3	$p \rightarrow \neg q$	premise
4	q	\rightarrow e 2, 1
5	$\neg q$	\rightarrow e 3, 1
6	\perp	\neg e 4, 5
7	r	\perp e 6

Example

\top is theorem: 1 \top \top i

- ▶ proof by contradiction

$$\frac{\begin{array}{c} \neg\varphi \\ \vdots \\ \perp \end{array}}{\varphi} \text{ PBC}$$

Definitions

- ▶ proof by contradiction

$$\frac{\begin{array}{|c} \neg\varphi \\ \vdots \\ \perp \end{array}}{\varphi} \text{ PBC}$$

- ▶ law of excluded middle

$$\frac{}{\varphi \vee \neg\varphi} \text{ LEM}$$

Example

$p \rightarrow q \vee r, q \rightarrow \neg p, \neg r \rightarrow p \vdash q \rightarrow r$ is valid:

1	$p \rightarrow q \vee r$	premise
2	$q \rightarrow \neg p$	premise
3	$\neg r \rightarrow p$	premise
4	q	assumption
5	$\neg p$	\rightarrow e 2, 4
6	$\neg r$	assumption
7	p	\rightarrow e 3, 6
8	\perp	\neg e 7, 5
9	r	PBC 6–8
10	$q \rightarrow r$	\rightarrow i 4–9

Example

$p \rightarrow q \vdash \neg p \vee q$ is valid:

1	$p \rightarrow q$	premise
2	$p \vee \neg p$	LEM
3	p	assumption
4	q	$\rightarrow e$ 1, 3
5	$\neg p \vee q$	$\vee i_2$ 4
6	$\neg p$	assumption
7	$\neg p \vee q$	$\vee i_1$ 6
8	$\neg p \vee q$	$\vee e$ 2, 3–5, 6–7

Summary of Natural Deduction ①

introduction

elimination

\wedge

$$\frac{\varphi \quad \psi}{\varphi \wedge \psi} \wedge i$$

$$\frac{\varphi \wedge \psi}{\varphi} \wedge e_1 \quad \frac{\varphi \wedge \psi}{\psi} \wedge e_2$$

\vee

$$\frac{\varphi}{\varphi \vee \psi} \vee i_1 \quad \frac{\psi}{\varphi \vee \psi} \vee i_2$$

$$\frac{\varphi \vee \psi \quad \begin{array}{|c|} \hline \varphi \\ \vdots \\ \chi \\ \hline \end{array} \quad \begin{array}{|c|} \hline \psi \\ \vdots \\ \chi \\ \hline \end{array}}{\chi} \vee e$$

\rightarrow

$$\frac{\begin{array}{|c|} \hline \varphi \\ \vdots \\ \psi \\ \hline \end{array}}{\varphi \rightarrow \psi} \rightarrow i$$

$$\frac{\varphi \rightarrow \psi \quad \varphi}{\psi} \rightarrow e$$

Summary of Natural Deduction ②

	introduction	elimination
\neg	$\frac{\boxed{\begin{array}{c} \varphi \\ \vdots \\ \perp \end{array}}}{\neg\varphi} \neg i$	$\frac{\varphi \quad \neg\varphi}{\perp} \neg e$
\perp		$\frac{\perp}{\varphi} \perp e$
\top	$\frac{}{\top} \top i$	
$\neg\neg$		$\frac{\neg\neg\varphi}{\varphi} \neg\neg e$

Summary of Natural Deduction ③

derived proof rules

$$\frac{\varphi \rightarrow \psi \quad \neg \psi}{\neg \varphi} \text{ MT}$$

$$\frac{\varphi}{\neg \neg \varphi} \text{ } \neg \neg \text{i}$$

$$\frac{\boxed{\begin{array}{c} \neg \varphi \\ \vdots \\ \perp \end{array}}}{\varphi} \text{ PBC}$$

$$\frac{}{\varphi \vee \neg \varphi} \text{ LEM}$$

Summary of Natural Deduction ③

derived proof rules

$$\frac{\varphi \rightarrow \psi \quad \neg \psi}{\neg \varphi} \text{ MT}$$

$$\frac{\varphi}{\neg \neg \varphi} \neg \neg \text{i}$$

$$\frac{\begin{array}{c} \neg \varphi \\ \vdots \\ \perp \end{array}}{\varphi} \text{ PBC}$$

$$\frac{}{\varphi \vee \neg \varphi} \text{ LEM}$$

Theorem

proof rules MT, $\neg \neg \text{i}$, PBC and LEM are **derivable** from other (**basic**) proof rules

Theorem

proof rules **MT**, $\neg\neg$ i, PBC and LEM are derivable from other (basic) proof rules

Proof

1 $\varphi \rightarrow \psi$ premise

2 $\neg\psi$ premise

6 $\neg\varphi$

Theorem

proof rules **MT**, $\neg\neg$ i, PBC and LEM are derivable from other (basic) proof rules

Proof

1	$\varphi \rightarrow \psi$	premise
2	$\neg\psi$	premise
3	φ	assumption
4	ψ	\rightarrow e 1, 3
5	\perp	\neg e 4, 2
6	$\neg\varphi$	\neg i 3-5

Theorem

proof rules MT, $\neg\neg$ i, PBC and LEM are derivable from other (basic) proof rules

Proof

1 φ premise

4 $\neg\neg\varphi$

Theorem

proof rules MT, $\neg\neg$ i, PBC and LEM are derivable from other (basic) proof rules

Proof

1	φ	premise
2	$\neg\varphi$	assumption
3	\perp	\neg e 1, 2
4	$\neg\neg\varphi$	\neg i 2-3

Theorem

proof rules MT, $\neg\neg$ i, **PBC** and LEM are derivable from other (basic) proof rules

Proof

1

$\neg\varphi$

hypothesis

\vdots

n

\perp

φ

conclusion

Theorem

proof rules MT, \neg -i, **PBC** and LEM are derivable from other (basic) proof rules

Proof

1	$\neg\varphi$	hypothesis
	\vdots	
n	\perp	
$n+1$	$\neg\neg\varphi$ \neg i 1- n	conclusion
	φ	

Theorem

proof rules MT, \neg -i, **PBC** and LEM are derivable from other (basic) proof rules

Proof

1	$\neg\varphi$	hypothesis
	\vdots	
n	\perp	
$n+1$	$\neg\neg\varphi$ \neg i 1- n	
$n+2$	φ \neg e $n+1$	conclusion

Theorem

proof rules MT, $\neg\neg$ i, PBC and **LEM** are derivable from other (basic) proof rules

Proof

$$\varphi \vee \neg\varphi$$

Theorem

proof rules MT, $\neg\neg$ i, PBC and **LEM** are derivable from other (basic) proof rules

Proof

1	$\neg(\varphi \vee \neg\varphi)$	assumption
2	φ	assumption
3	$\varphi \vee \neg\varphi$	$\vee i_1$ 2
4	\perp	$\neg e$ 3,1
5	$\neg\varphi$	$\neg i$ 2-4
6	$\varphi \vee \neg\varphi$	$\vee i_2$ 5
7	\perp	$\neg e$ 6,1
8	$\neg\neg(\varphi \vee \neg\varphi)$	$\neg i$ 1-7
9	$\varphi \vee \neg\varphi$	$\neg\neg e$ 8

Theorem

proof rules LEM, PBC and $\neg\neg e$ are **inter-derivable** (with respect to other basic proof rules)

Theorem

proof rules LEM, PBC and $\neg\neg e$ are inter-derivable (with respect to other basic proof rules)

Remark

- ▶ LEM, PBC and $\neg\neg e$ are **controversial** because they are not **constructive**

Theorem

proof rules LEM, PBC and $\neg\neg e$ are inter-derivable (with respect to other basic proof rules)

Remark

- ▶ LEM, PBC and $\neg\neg e$ are controversial because they are not constructive
- ▶ **classical** logicians use all proof rules

Theorem

proof rules LEM, PBC and $\neg\neg e$ are inter-derivable (with respect to other basic proof rules)

Remark

- ▶ LEM, PBC and $\neg\neg e$ are controversial because they are not constructive
- ▶ classical logicians use all proof rules
- ▶ **intuitionistic** logicians do not use LEM, PBC and $\neg\neg e$

Theorem

proof rules LEM, PBC and $\neg\neg e$ are inter-derivable (with respect to other basic proof rules)

Remark

- ▶ LEM, PBC and $\neg\neg e$ are controversial because they are not constructive
- ▶ classical logicians use all proof rules
- ▶ intuitionistic logicians do not use LEM, PBC and $\neg\neg e$

Example

- ▶ formula $((p \rightarrow q) \rightarrow p) \rightarrow p$ is valid

Theorem

proof rules LEM, PBC and $\neg\neg e$ are inter-derivable (with respect to other basic proof rules)

Remark

- ▶ LEM, PBC and $\neg\neg e$ are controversial because they are not constructive
- ▶ classical logicians use all proof rules
- ▶ intuitionistic logicians do not use LEM, PBC and $\neg\neg e$

Example

- ▶ formula $((p \rightarrow q) \rightarrow p) \rightarrow p$ is valid
- ▶ sequent $\vdash ((p \rightarrow q) \rightarrow p) \rightarrow p$ is valid but proof requires LEM, PBC or $\neg\neg e$

Outline

1. Summary of Previous Lecture
2. Natural Deduction
- 3. Intermezzo**
4. Soundness
5. Further Reading

Question

Which of the following statements are true ?

- A** The sequent $p \rightarrow q \vdash \neg q \rightarrow \neg p$ is provable in natural deduction.
- B** The sequent $(p \rightarrow q) \rightarrow p \vdash p$ is only derivable in classical logic.
- C** Valid sequents can have arbitrarily long natural deduction proofs.
- D** Every valid sequent has a proof without $\neg\neg$ i.
- E** Intuitionistic logicians can prove more statements than classical logicians.
- F** Every natural deduction proof of sequent with n premises has at least n lines.



Question

Which of the following statements are true ?

- ✓ The sequent $p \rightarrow q \vdash \neg q \rightarrow \neg p$ is provable in natural deduction.
- ✓ The sequent $(p \rightarrow q) \rightarrow p \vdash p$ is only derivable in classical logic.
- ✓ Valid sequents can have arbitrarily long natural deduction proofs.
- ✓ Every valid sequent has a proof without $\neg\neg$ i.
- E Intuitionistic logicians can prove more statements than classical logicians.
- F Every natural deduction proof of sequent with n premises has at least n lines.



Outline

1. Summary of Previous Lecture
2. Natural Deduction
3. Intermezzo
- 4. Soundness**
5. Further Reading

Theorem

propositional logic is **sound**:

$$\varphi_1, \varphi_2, \dots, \varphi_n \vdash \psi \text{ is valid} \implies \varphi_1, \varphi_2, \dots, \varphi_n \models \psi$$

Theorem

propositional logic is **sound**:

$$\varphi_1, \varphi_2, \dots, \varphi_n \vdash \psi \text{ is valid} \implies \varphi_1, \varphi_2, \dots, \varphi_n \vDash \psi$$

only true statements can be proved

Theorem

natural deduction is sound:

$$\varphi_1, \varphi_2, \dots, \varphi_n \vdash \psi \text{ is valid} \implies \varphi_1, \varphi_2, \dots, \varphi_n \vDash \psi$$

only true statements can be proved

Theorem

natural deduction is sound:

$$\varphi_1, \varphi_2, \dots, \varphi_n \vdash \psi \text{ is valid} \implies \varphi_1, \varphi_2, \dots, \varphi_n \vDash \psi$$

only true statements can be proved

Corollary

theorems are valid

Idea

use **induction** on length of natural deduction proof and **case analysis** of last proof step

Idea

use **induction** on length of natural deduction proof and **case analysis** of last proof step

Problem

initial part of proof need not correspond to sequent with same premises

1	$p \rightarrow q$	premise
2	$p \rightarrow \neg q$	premise
3	p	assumption
4	q	\rightarrow e 1, 3
5	$\neg q$	\rightarrow e 2, 3
6	\perp	\neg e 4, 5
7	$\neg p$	\neg i 3-6

Idea

use **induction** on length of natural deduction proof and **case analysis** of last proof step

Problem

initial part of proof need not correspond to sequent with same premises

1 $p \rightarrow q$ premise

2 $p \rightarrow \neg q$ premise

3 p assumption

4 q \rightarrow e 1, 3

5 $\neg q$ \rightarrow e 2, 3

6 \perp \neg e 4, 5

7 $\neg p$ \neg i 3-6

$p \rightarrow q, p \rightarrow \neg q \vdash p \rightarrow q$

$p \rightarrow q, p \rightarrow \neg q \vdash p \rightarrow \neg q$

$p \rightarrow q, p \rightarrow \neg q \vdash \neg p$

Idea

use **induction** on length of natural deduction proof and **case analysis** of last proof step

Problem

initial part of proof need not correspond to sequent with same premises

1	$p \rightarrow q$	premise	$p \rightarrow q, p \rightarrow \neg q \vdash p \rightarrow q$
2	$p \rightarrow \neg q$	premise	$p \rightarrow q, p \rightarrow \neg q \vdash p \rightarrow \neg q$
3	p	assumption	?
4	q	\rightarrow e 1, 3	?
5	$\neg q$	\rightarrow e 2, 3	?
6	\perp	\neg e 4, 5	?
7	$\neg p$	\neg i 3-6	$p \rightarrow q, p \rightarrow \neg q \vdash \neg p$

Solution

add assumptions to sequents

1 $p \rightarrow q$ premise

2 $p \rightarrow \neg q$ premise

3 p assumption

4 q \rightarrow e 1, 3

5 $\neg q$ \rightarrow e 2, 3

6 \perp \neg e 4, 5

7 $\neg p$ \neg i 3-6

$p \rightarrow q, p \rightarrow \neg q \vdash p \rightarrow q$

$p \rightarrow q, p \rightarrow \neg q \vdash p \rightarrow \neg q$

$p \rightarrow q, p \rightarrow \neg q \vdash \neg p$

Solution

add assumptions to sequents

1 $p \rightarrow q$ premise

2 $p \rightarrow \neg q$ premise

3 p assumption

4 q \rightarrow e 1, 3

5 $\neg q$ \rightarrow e 2, 3

6 \perp \neg e 4, 5

7 $\neg p$ \neg i 3-6

$p \rightarrow q, p \rightarrow \neg q \vdash p \rightarrow q$

$p \rightarrow q, p \rightarrow \neg q \vdash p \rightarrow \neg q$

$p \rightarrow q, p \rightarrow \neg q; p \vdash p$

$p \rightarrow q, p \rightarrow \neg q; p \vdash q$

$p \rightarrow q, p \rightarrow \neg q; p \vdash \neg q$

$p \rightarrow q, p \rightarrow \neg q; p \vdash \perp$

$p \rightarrow q, p \rightarrow \neg q \vdash \neg p$

Solution

add assumptions to sequents

1	$p \rightarrow q$	premise	$p \rightarrow q, p \rightarrow \neg q \vdash p \rightarrow q$
2	$p \rightarrow \neg q$	premise	$p \rightarrow q, p \rightarrow \neg q \vdash p \rightarrow \neg q$
3	p	assumption	$p \rightarrow q, p \rightarrow \neg q; p \vdash p$
4	q	$\rightarrow e$ 1, 3	$p \rightarrow q, p \rightarrow \neg q; p \vdash q$
5	$\neg q$	$\rightarrow e$ 2, 3	$p \rightarrow q, p \rightarrow \neg q; p \vdash \neg q$
6	\perp	$\neg e$ 4, 5	$p \rightarrow q, p \rightarrow \neg q; p \vdash \perp$
7	$\neg p$	$\neg i$ 3-6	$p \rightarrow q, p \rightarrow \neg q \vdash \neg p$

Definition

extended sequent

$$\underbrace{\varphi_1, \varphi_2, \dots, \varphi_n}_{\text{premises}}; \underbrace{\psi_1, \psi_2, \dots, \psi_m}_{\text{assumptions}} \vdash \underbrace{\chi}_{\text{conclusion}}$$

Φ_1 Φ_2

Example

1 $p \wedge q \rightarrow r$ premise

2 p assumption

3 q assumption

4 $p \wedge q$ \wedge i 2, 3

5 r \rightarrow e 1, 4

6 $q \rightarrow r$ \rightarrow i 3-5

7 $p \rightarrow (q \rightarrow r)$ \rightarrow i 2-6

$p \wedge q \rightarrow r \vdash p \wedge q \rightarrow r$

$p \wedge q \rightarrow r; p \vdash p$

$p \wedge q \rightarrow r; p, q \vdash q$

$p \wedge q \rightarrow r; p, q \vdash p \wedge q$

$p \wedge q \rightarrow r; p, q \vdash r$

$p \wedge q \rightarrow r; p \vdash q \rightarrow r$

$p \wedge q \rightarrow r \vdash p \rightarrow (q \rightarrow r)$

Example

1	$p \wedge q \rightarrow r$	premise	$p \wedge q \rightarrow r \vdash p \wedge q \rightarrow r$
2	p	assumption	$p \wedge q \rightarrow r; p \vdash p$
3	q	assumption	$p \wedge q \rightarrow r; p, q \vdash q$
4	$p \wedge q$	\wedge i 2, 3	$p \wedge q \rightarrow r; p, q \vdash p \wedge q$
5	r	\rightarrow e 1, 4	$p \wedge q \rightarrow r; p, q \vdash r$
6	$q \rightarrow r$	\rightarrow i 3-5	$p \wedge q \rightarrow r; p \vdash q \rightarrow r$
7	$p \rightarrow (q \rightarrow r)$	\rightarrow i 2-6	$p \wedge q \rightarrow r \vdash p \rightarrow (q \rightarrow r)$

Soundness Proof

by induction on length of proof of

$$\Phi_1; \Phi_2 \vdash \psi$$

we prove

$$\Phi_1, \Phi_2 \models \psi$$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Base Case

► premise $\psi \in \Phi_1 \implies \Phi_1, \Phi_2 \models \psi$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Base Cases

- ▶ premise $\psi \in \Phi_1 \implies \Phi_1, \Phi_2 \models \psi$
- ▶ assumption $\psi \in \Phi_2 \implies \Phi_1, \Phi_2 \models \psi$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Base Cases

- ▶ premise $\psi \in \Phi_1 \implies \Phi_1, \Phi_2 \models \psi$
- ▶ assumption $\psi \in \Phi_2 \implies \Phi_1, \Phi_2 \models \psi$
- ▶ \top $\psi = \top \implies \Phi_1, \Phi_2 \models \psi$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step)

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step) \wedge i

$$\psi = \psi_1 \wedge \psi_2$$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step) \wedge i

$\psi = \psi_1 \wedge \psi_2$

shorter proofs $\Phi_1; \Phi_2^1 \vdash \psi_1$ and $\Phi_1; \Phi_2^2 \vdash \psi_2$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \vDash \psi$

Induction Step (case analysis of last proof step) \wedge i

$\psi = \psi_1 \wedge \psi_2$

shorter proofs $\Phi_1; \Phi_2^1 \vdash \psi_1$ and $\Phi_1; \Phi_2^2 \vdash \psi_2$ with $\Phi_2^1, \Phi_2^2 \subseteq \Phi_2$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step) $\wedge i$

$\psi = \psi_1 \wedge \psi_2$

shorter proofs $\Phi_1; \Phi_2^1 \vdash \psi_1$ and $\Phi_1; \Phi_2^2 \vdash \psi_2$ with $\Phi_2^1, \Phi_2^2 \subseteq \Phi_2$

induction hypothesis: $\Phi_1, \Phi_2^1 \models \psi_1$ and $\Phi_1, \Phi_2^2 \models \psi_2$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step) \wedge i

$$\psi = \psi_1 \wedge \psi_2$$

shorter proofs $\Phi_1; \Phi_2^1 \vdash \psi_1$ and $\Phi_1; \Phi_2^2 \vdash \psi_2$ with $\Phi_2^1, \Phi_2^2 \subseteq \Phi_2$

induction hypothesis: $\Phi_1, \Phi_2^1 \models \psi_1$ and $\Phi_1, \Phi_2^2 \models \psi_2$

$$\bar{v}(\varphi) = \top \text{ for all } \varphi \in \Phi_1, \Phi_2$$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step) \wedge i

$\psi = \psi_1 \wedge \psi_2$

shorter proofs $\Phi_1; \Phi_2^1 \vdash \psi_1$ and $\Phi_1; \Phi_2^2 \vdash \psi_2$ with $\Phi_2^1, \Phi_2^2 \subseteq \Phi_2$

induction hypothesis: $\Phi_1, \Phi_2^1 \models \psi_1$ and $\Phi_1, \Phi_2^2 \models \psi_2$

$\bar{v}(\varphi) = \top$ for all $\varphi \in \Phi_1, \Phi_2 \implies \bar{v}(\varphi) = \top$ for all $\varphi \in \Phi_1, \Phi_2^1, \Phi_2^2$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step) \wedge i

$\psi = \psi_1 \wedge \psi_2$

shorter proofs $\Phi_1; \Phi_2^1 \vdash \psi_1$ and $\Phi_1; \Phi_2^2 \vdash \psi_2$ with $\Phi_2^1, \Phi_2^2 \subseteq \Phi_2$

induction hypothesis: $\Phi_1, \Phi_2^1 \models \psi_1$ and $\Phi_1, \Phi_2^2 \models \psi_2$

$$\begin{aligned} \bar{v}(\varphi) = \top \text{ for all } \varphi \in \Phi_1, \Phi_2 &\implies \bar{v}(\varphi) = \top \text{ for all } \varphi \in \Phi_1, \Phi_2^1, \Phi_2^2 \\ &\implies \bar{v}(\psi_1) = \bar{v}(\psi_2) = \top \end{aligned}$$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step) \wedge i

$\psi = \psi_1 \wedge \psi_2$

shorter proofs $\Phi_1; \Phi_2^1 \vdash \psi_1$ and $\Phi_1; \Phi_2^2 \vdash \psi_2$ with $\Phi_2^1, \Phi_2^2 \subseteq \Phi_2$

induction hypothesis: $\Phi_1, \Phi_2^1 \models \psi_1$ and $\Phi_1, \Phi_2^2 \models \psi_2$

$\bar{v}(\varphi) = \text{T}$ for all $\varphi \in \Phi_1, \Phi_2 \implies \bar{v}(\varphi) = \text{T}$ for all $\varphi \in \Phi_1, \Phi_2^1, \Phi_2^2$

$\implies \bar{v}(\psi_1) = \bar{v}(\psi_2) = \text{T}$

$\implies \bar{v}(\psi_1 \wedge \psi_2) = \text{T}$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step) \wedge i

$\psi = \psi_1 \wedge \psi_2$

shorter proofs $\Phi_1; \Phi_2^1 \vdash \psi_1$ and $\Phi_1; \Phi_2^2 \vdash \psi_2$ with $\Phi_2^1, \Phi_2^2 \subseteq \Phi_2$

induction hypothesis: $\Phi_1, \Phi_2^1 \models \psi_1$ and $\Phi_1, \Phi_2^2 \models \psi_2$

$\bar{v}(\varphi) = \top$ for all $\varphi \in \Phi_1, \Phi_2 \implies \bar{v}(\varphi) = \top$ for all $\varphi \in \Phi_1, \Phi_2^1, \Phi_2^2$

$\implies \bar{v}(\psi_1) = \bar{v}(\psi_2) = \top$

$\implies \bar{v}(\psi_1 \wedge \psi_2) = \top$

hence $\Phi_1, \Phi_2 \models \psi$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step) $\rightarrow i$

$\psi = \psi_1 \rightarrow \psi_2$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step) $\rightarrow i$

$\psi = \psi_1 \rightarrow \psi_2$

shorter proof $\Phi_1; \Phi_2^1, \psi_1 \vdash \psi_2$ with $\Phi_2^1 \subseteq \Phi_2$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step) $\rightarrow i$

$\psi = \psi_1 \rightarrow \psi_2$

shorter proof $\Phi_1; \Phi_2^1, \psi_1 \vdash \psi_2$ with $\Phi_2^1 \subseteq \Phi_2$

induction hypothesis: $\Phi_1, \Phi_2^1, \psi_1 \models \psi_2$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step) $\rightarrow i$

$\psi = \psi_1 \rightarrow \psi_2$

shorter proof $\Phi_1; \Phi_2^1, \psi_1 \vdash \psi_2$ with $\Phi_2^1 \subseteq \Phi_2$

induction hypothesis: $\Phi_1, \Phi_2^1, \psi_1 \models \psi_2$

$\bar{v}(\varphi) = \top$ for all $\varphi \in \Phi_1, \Phi_2$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step) $\rightarrow i$

$\psi = \psi_1 \rightarrow \psi_2$

shorter proof $\Phi_1; \Phi_2^1, \psi_1 \vdash \psi_2$ with $\Phi_2^1 \subseteq \Phi_2$

induction hypothesis: $\Phi_1, \Phi_2^1, \psi_1 \models \psi_2$

$\bar{v}(\varphi) = T$ for all $\varphi \in \Phi_1, \Phi_2$

two cases

$$\bar{v}(\psi_1) = F$$

$$\bar{v}(\psi_1) = T$$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step) $\rightarrow i$

$\psi = \psi_1 \rightarrow \psi_2$

shorter proof $\Phi_1; \Phi_2^1, \psi_1 \vdash \psi_2$ with $\Phi_2^1 \subseteq \Phi_2$

induction hypothesis: $\Phi_1, \Phi_2^1, \psi_1 \models \psi_2$

$\bar{v}(\varphi) = \text{T}$ for all $\varphi \in \Phi_1, \Phi_2$

two cases

$$\bar{v}(\psi_1) = \text{F} \implies \bar{v}(\psi_1 \rightarrow \psi_2) = \text{T}$$

$$\bar{v}(\psi_1) = \text{T}$$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step) $\rightarrow i$

$\psi = \psi_1 \rightarrow \psi_2$

shorter proof $\Phi_1; \Phi_2^1, \psi_1 \vdash \psi_2$ with $\Phi_2^1 \subseteq \Phi_2$

induction hypothesis: $\Phi_1, \Phi_2^1, \psi_1 \models \psi_2$

$\bar{v}(\varphi) = T$ for all $\varphi \in \Phi_1, \Phi_2$

two cases

$$\bar{v}(\psi_1) = F \implies \bar{v}(\psi_1 \rightarrow \psi_2) = T$$

$$\bar{v}(\psi_1) = T \implies \bar{v}(\varphi) = T \text{ for all } \varphi \in \Phi_1, \Phi_2^1, \psi_1$$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step) $\rightarrow i$

$\psi = \psi_1 \rightarrow \psi_2$

shorter proof $\Phi_1; \Phi_2^1, \psi_1 \vdash \psi_2$ with $\Phi_2^1 \subseteq \Phi_2$

induction hypothesis: $\Phi_1, \Phi_2^1, \psi_1 \models \psi_2$

$\bar{v}(\varphi) = T$ for all $\varphi \in \Phi_1, \Phi_2$

two cases

$$\bar{v}(\psi_1) = F \implies \bar{v}(\psi_1 \rightarrow \psi_2) = T$$

$$\bar{v}(\psi_1) = T \implies \bar{v}(\varphi) = T \text{ for all } \varphi \in \Phi_1, \Phi_2^1, \psi_1 \implies \bar{v}(\psi_2) = T$$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step) $\rightarrow i$

$\psi = \psi_1 \rightarrow \psi_2$

shorter proof $\Phi_1; \Phi_2^1, \psi_1 \vdash \psi_2$ with $\Phi_2^1 \subseteq \Phi_2$

induction hypothesis: $\Phi_1, \Phi_2^1, \psi_1 \models \psi_2$

$\bar{v}(\varphi) = T$ for all $\varphi \in \Phi_1, \Phi_2$

two cases

$$\bar{v}(\psi_1) = F \implies \bar{v}(\psi_1 \rightarrow \psi_2) = T$$

$$\bar{v}(\psi_1) = T \implies \bar{v}(\varphi) = T \text{ for all } \varphi \in \Phi_1, \Phi_2^1, \psi_1 \implies \bar{v}(\psi_2) = T \implies \bar{v}(\psi_1 \rightarrow \psi_2) = T$$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step) $\rightarrow i$

$\psi = \psi_1 \rightarrow \psi_2$

shorter proof $\Phi_1; \Phi_2^1, \psi_1 \vdash \psi_2$ with $\Phi_2^1 \subseteq \Phi_2$

induction hypothesis: $\Phi_1, \Phi_2^1, \psi_1 \models \psi_2$

$\bar{v}(\varphi) = \text{T}$ for all $\varphi \in \Phi_1, \Phi_2$

two cases

$$\bar{v}(\psi_1) = \text{F} \implies \bar{v}(\psi_1 \rightarrow \psi_2) = \text{T}$$

$$\bar{v}(\psi_1) = \text{T} \implies \bar{v}(\varphi) = \text{T} \text{ for all } \varphi \in \Phi_1, \Phi_2^1, \psi_1 \implies \bar{v}(\psi_2) = \text{T} \implies \bar{v}(\psi_1 \rightarrow \psi_2) = \text{T}$$

hence $\Phi_1, \Phi_2 \models \psi$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step) \neg e

$\psi = \perp$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step) $\neg e$

$\psi = \perp$

shorter proofs $\Phi_1; \Phi_2^1 \vdash \psi'$ and $\Phi_1; \Phi_2^2 \vdash \neg\psi'$ with $\Phi_2^1, \Phi_2^2 \subseteq \Phi_2$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step) $\neg e$

$\psi = \perp$

shorter proofs $\Phi_1; \Phi_2^1 \vdash \psi'$ and $\Phi_1; \Phi_2^2 \vdash \neg\psi'$ with $\Phi_2^1, \Phi_2^2 \subseteq \Phi_2$

induction hypothesis: $\Phi_1, \Phi_2^1 \models \psi'$ and $\Phi_1, \Phi_2^2 \models \neg\psi'$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step) $\neg e$

$\psi = \perp$

shorter proofs $\Phi_1; \Phi_2^1 \vdash \psi'$ and $\Phi_1; \Phi_2^2 \vdash \neg\psi'$ with $\Phi_2^1, \Phi_2^2 \subseteq \Phi_2$

induction hypothesis: $\Phi_1, \Phi_2^1 \models \psi'$ and $\Phi_1, \Phi_2^2 \models \neg\psi'$

$\bar{v}(\varphi) = \top$ for all $\varphi \in \Phi_1, \Phi_2$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step) $\neg e$

$\psi = \perp$

shorter proofs $\Phi_1; \Phi_2^1 \vdash \psi'$ and $\Phi_1; \Phi_2^2 \vdash \neg\psi'$ with $\Phi_2^1, \Phi_2^2 \subseteq \Phi_2$

induction hypothesis: $\Phi_1, \Phi_2^1 \models \psi'$ and $\Phi_1, \Phi_2^2 \models \neg\psi'$

$\bar{v}(\varphi) = T$ for all $\varphi \in \Phi_1, \Phi_2 \implies \bar{v}(\varphi) = T$ for all $\varphi \in \Phi_1, \Phi_2^1, \Phi_2^2$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step) $\neg e$

$\psi = \perp$

shorter proofs $\Phi_1; \Phi_2^1 \vdash \psi'$ and $\Phi_1; \Phi_2^2 \vdash \neg\psi'$ with $\Phi_2^1, \Phi_2^2 \subseteq \Phi_2$

induction hypothesis: $\Phi_1, \Phi_2^1 \models \psi'$ and $\Phi_1, \Phi_2^2 \models \neg\psi'$

$$\begin{aligned} \bar{v}(\varphi) = \text{T for all } \varphi \in \Phi_1, \Phi_2 &\implies \bar{v}(\varphi) = \text{T for all } \varphi \in \Phi_1, \Phi_2^1, \Phi_2^2 \\ &\implies \bar{v}(\psi') = \text{T and } \bar{v}(\neg\psi') = \text{T} \end{aligned}$$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step) $\neg e$

$\psi = \perp$

shorter proofs $\Phi_1; \Phi_2^1 \vdash \psi'$ and $\Phi_1; \Phi_2^2 \vdash \neg\psi'$ with $\Phi_2^1, \Phi_2^2 \subseteq \Phi_2$

induction hypothesis: $\Phi_1, \Phi_2^1 \models \psi'$ and $\Phi_1, \Phi_2^2 \models \neg\psi'$

$\bar{v}(\varphi) = \text{T}$ for all $\varphi \in \Phi_1, \Phi_2 \implies \bar{v}(\varphi) = \text{T}$ for all $\varphi \in \Phi_1, \Phi_2^1, \Phi_2^2$

$\implies \bar{v}(\psi') = \text{T}$ and $\bar{v}(\neg\psi') = \text{T}$ ⚡

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step) $\neg e$

$\psi = \perp$

shorter proofs $\Phi_1; \Phi_2^1 \vdash \psi'$ and $\Phi_1; \Phi_2^2 \vdash \neg\psi'$ with $\Phi_2^1, \Phi_2^2 \subseteq \Phi_2$

induction hypothesis: $\Phi_1, \Phi_2^1 \models \psi'$ and $\Phi_1, \Phi_2^2 \models \neg\psi'$

$\bar{v}(\varphi) = \text{T}$ for all $\varphi \in \Phi_1, \Phi_2 \implies \bar{v}(\varphi) = \text{T}$ for all $\varphi \in \Phi_1, \Phi_2^1, \Phi_2^2$

$\implies \bar{v}(\psi') = \text{T}$ and $\bar{v}(\neg\psi') = \text{T}$ ⚡

hence $\Phi_1, \Phi_2 \models \perp$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step) $\forall e$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step) $\forall e$

shorter proofs $\Phi_1; \Phi_2^1 \vdash \psi_1 \vee \psi_2$ and $\Phi_1; \Phi_2^2, \psi_1 \vdash \psi$ and $\Phi_1; \Phi_2^3, \psi_2 \vdash \psi$ with $\Phi_2^1, \Phi_2^2, \Phi_2^3 \subseteq \Phi_2$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \vDash \psi$

Induction Step (case analysis of last proof step) $\vee e$

shorter proofs $\Phi_1; \Phi_2^1 \vdash \psi_1 \vee \psi_2$ and $\Phi_1; \Phi_2^2, \psi_1 \vdash \psi$ and $\Phi_1; \Phi_2^3, \psi_2 \vdash \psi$ with $\Phi_2^1, \Phi_2^2, \Phi_2^3 \subseteq \Phi_2$

induction hypothesis: $\Phi_1, \Phi_2^1 \vDash \psi_1 \vee \psi_2$ and $\Phi_1; \Phi_2^2, \psi_1 \vDash \psi$ and $\Phi_1; \Phi_2^3, \psi_2 \vDash \psi$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step) $\forall e$

shorter proofs $\Phi_1; \Phi_2^1 \vdash \psi_1 \vee \psi_2$ and $\Phi_1; \Phi_2^2, \psi_1 \vdash \psi$ and $\Phi_1; \Phi_2^3, \psi_2 \vdash \psi$ with $\Phi_2^1, \Phi_2^2, \Phi_2^3 \subseteq \Phi_2$

induction hypothesis: $\Phi_1, \Phi_2^1 \models \psi_1 \vee \psi_2$ and $\Phi_1; \Phi_2^2, \psi_1 \models \psi$ and $\Phi_1; \Phi_2^3, \psi_2 \models \psi$

$\bar{v}(\varphi) = \top$ for all $\varphi \in \Phi_1, \Phi_2$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step) $\vee e$

shorter proofs $\Phi_1; \Phi_2^1 \vdash \psi_1 \vee \psi_2$ and $\Phi_1; \Phi_2^2, \psi_1 \vdash \psi$ and $\Phi_1; \Phi_2^3, \psi_2 \vdash \psi$ with $\Phi_2^1, \Phi_2^2, \Phi_2^3 \subseteq \Phi_2$

induction hypothesis: $\Phi_1, \Phi_2^1 \models \psi_1 \vee \psi_2$ and $\Phi_1; \Phi_2^2, \psi_1 \models \psi$ and $\Phi_1; \Phi_2^3, \psi_2 \models \psi$

$\bar{v}(\varphi) = \text{T}$ for all $\varphi \in \Phi_1, \Phi_2 \implies v(\psi_1 \vee \psi_2) = \text{T}$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step) $\vee e$

shorter proofs $\Phi_1; \Phi_2^1 \vdash \psi_1 \vee \psi_2$ and $\Phi_1; \Phi_2^2, \psi_1 \vdash \psi$ and $\Phi_1; \Phi_2^3, \psi_2 \vdash \psi$ with $\Phi_2^1, \Phi_2^2, \Phi_2^3 \subseteq \Phi_2$

induction hypothesis: $\Phi_1, \Phi_2^1 \models \psi_1 \vee \psi_2$ and $\Phi_1; \Phi_2^2, \psi_1 \models \psi$ and $\Phi_1; \Phi_2^3, \psi_2 \models \psi$

$\bar{v}(\varphi) = \top$ for all $\varphi \in \Phi_1, \Phi_2 \implies v(\psi_1 \vee \psi_2) = \top$

two cases

$$\bar{v}(\psi_1) = \top$$

$$\bar{v}(\psi_2) = \top$$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step) $\vee e$

shorter proofs $\Phi_1; \Phi_2^1 \vdash \psi_1 \vee \psi_2$ and $\Phi_1; \Phi_2^2, \psi_1 \vdash \psi$ and $\Phi_1; \Phi_2^3, \psi_2 \vdash \psi$ with $\Phi_2^1, \Phi_2^2, \Phi_2^3 \subseteq \Phi_2$

induction hypothesis: $\Phi_1, \Phi_2^1 \models \psi_1 \vee \psi_2$ and $\Phi_1; \Phi_2^2, \psi_1 \models \psi$ and $\Phi_1; \Phi_2^3, \psi_2 \models \psi$

$\bar{v}(\varphi) = \top$ for all $\varphi \in \Phi_1, \Phi_2 \implies v(\psi_1 \vee \psi_2) = \top$

two cases

$$\bar{v}(\psi_1) = \top \implies \bar{v}(\psi) = \top$$

$$\bar{v}(\psi_2) = \top$$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step) $\vee e$

shorter proofs $\Phi_1; \Phi_2^1 \vdash \psi_1 \vee \psi_2$ and $\Phi_1; \Phi_2^2, \psi_1 \vdash \psi$ and $\Phi_1; \Phi_2^3, \psi_2 \vdash \psi$ with $\Phi_2^1, \Phi_2^2, \Phi_2^3 \subseteq \Phi_2$

induction hypothesis: $\Phi_1, \Phi_2^1 \models \psi_1 \vee \psi_2$ and $\Phi_1; \Phi_2^2, \psi_1 \models \psi$ and $\Phi_1; \Phi_2^3, \psi_2 \models \psi$

$\bar{v}(\varphi) = \text{T}$ for all $\varphi \in \Phi_1, \Phi_2 \implies v(\psi_1 \vee \psi_2) = \text{T}$

two cases

$$\bar{v}(\psi_1) = \text{T} \implies \bar{v}(\psi) = \text{T}$$

$$\bar{v}(\psi_2) = \text{T} \implies \bar{v}(\psi) = \text{T}$$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step) $\vee e$

shorter proofs $\Phi_1; \Phi_2^1 \vdash \psi_1 \vee \psi_2$ and $\Phi_1; \Phi_2^2, \psi_1 \vdash \psi$ and $\Phi_1; \Phi_2^3, \psi_2 \vdash \psi$ with $\Phi_2^1, \Phi_2^2, \Phi_2^3 \subseteq \Phi_2$

induction hypothesis: $\Phi_1, \Phi_2^1 \models \psi_1 \vee \psi_2$ and $\Phi_1; \Phi_2^2, \psi_1 \models \psi$ and $\Phi_1; \Phi_2^3, \psi_2 \models \psi$

$\bar{v}(\varphi) = \text{T}$ for all $\varphi \in \Phi_1, \Phi_2 \implies v(\psi_1 \vee \psi_2) = \text{T}$

two cases

$$\bar{v}(\psi_1) = \text{T} \implies \bar{v}(\psi) = \text{T}$$

$$\bar{v}(\psi_2) = \text{T} \implies \bar{v}(\psi) = \text{T}$$

hence $\Phi_1, \Phi_2 \models \psi$

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step)

remaining (basic) proof rules

$\wedge e_1 \quad \wedge e_2 \quad \forall i_1 \quad \forall i_2 \quad \rightarrow e \quad \neg i \quad \perp e \quad \neg \neg e$

are similar

Claim

$\Phi_1; \Phi_2 \vdash \psi$ is valid $\implies \Phi_1, \Phi_2 \models \psi$

Induction Step (case analysis of last proof step)

remaining (basic) proof rules

$\wedge e_1 \quad \wedge e_2 \quad \forall i_1 \quad \forall i_2 \quad \rightarrow e \quad \neg i \quad \perp e \quad \neg \neg e$

are similar

Corollary

natural deduction is sound

Outline

1. Summary of Previous Lecture
2. Natural Deduction
3. Intermezzo
4. Soundness
- 5. Further Reading**

- ▶ Section 1.2
- ▶ Sections 1.4.2 and 1.4.3

Huth and Ryan

- ▶ Section 1.2
- ▶ Sections 1.4.2 and 1.4.3

Intuitionistic Logic

- ▶ Wikipedia

[accessed December 27, 2024]

Huth and Ryan

- ▶ Section 1.2
- ▶ Sections 1.4.2 and 1.4.3

Intuitionistic Logic

- ▶ Wikipedia

[accessed December 27, 2024]

Differences (slides – book)

- ▶ top introduction proof rule
- ▶ order of premises in $\rightarrow e$

Important Concepts

- ▶ and elimination
- ▶ and introduction
- ▶ assumption
- ▶ bottom elimination
- ▶ derived proof rule
- ▶ double negation introduction
- ▶ double negation elimination
- ▶ extended sequent
- ▶ implication elimination
- ▶ implication introduction
- ▶ intuitionistic logic
- ▶ law of excluded middle
- ▶ modus ponens
- ▶ modus tollens
- ▶ natural deduction
- ▶ negation elimination
- ▶ negation introduction
- ▶ or elimination
- ▶ or introduction
- ▶ premise
- ▶ proof by contradiction
- ▶ sequent
- ▶ soundness
- ▶ theorem
- ▶ top introduction
- ▶ validity of sequents

Important Concepts

- ▶ and elimination
- ▶ and introduction
- ▶ assumption
- ▶ bottom elimination
- ▶ derived proof rule
- ▶ double negation introduction
- ▶ double negation elimination
- ▶ extended sequent
- ▶ implication elimination
- ▶ implication introduction
- ▶ intuitionistic logic
- ▶ law of excluded middle
- ▶ modus ponens
- ▶ modus tollens
- ▶ natural deduction
- ▶ negation elimination
- ▶ negation introduction
- ▶ or elimination
- ▶ or introduction
- ▶ premise
- ▶ proof by contradiction
- ▶ sequent
- ▶ soundness
- ▶ theorem
- ▶ top introduction
- ▶ validity of sequents

homework for March 26