



# Term Rewriting

Philipp Dablander and **Aart Middeldorp**

## Initial Remarks

- ▶ **Term Rewriting** is part of compulsory module 20 ("Ausgewählte Kapitel") in bachelor program Computer Science

## Initial Remarks

- ▶ **Term Rewriting** is part of compulsory module 20 ("Ausgewählte Kapitel") in bachelor program Computer Science
- ▶ **Term Rewriting** is part of elective module 20 ("individual choice of specialization") in master program Computer Science

## Initial Remarks

- ▶ **Term Rewriting** is part of compulsory module 20 ("Ausgewählte Kapitel") in bachelor program Computer Science
- ▶ **Term Rewriting** is part of elective module 20 ("individual choice of specialization") in master program Computer Science
- ▶ any course offered by MIP can be used for "Ausgewählte Kapitel"

## Initial Remarks

- ▶ **Term Rewriting** is part of compulsory module 20 ("Ausgewählte Kapitel") in bachelor program Computer Science
- ▶ **Term Rewriting** is part of elective module 20 ("individual choice of specialization") in master program Computer Science
- ▶ any course offered by MIP can be used for "Ausgewählte Kapitel"
- ▶ other master courses with theory content (**Logic and Learning** specialization):
  - ▶ WM 2: Constraint Solving
  - ▶ WM 7: Interactive Theorem Proving in Isabelle/HOL
  - ▶ WM 7: Current Challenges in Probabilistic Learning, Inference, and their Applications
  - ▶ WM 8: Advanced Logic and Quantum Logic
  - ▶ WM 9: Research Seminar CL/TCS

# Outline

- 1. Organisation**
- 2. Examples**
- 3. Terms**
- 4. Exercises**
- 5. Further Reading**

- ▶ LVA 703141

- ▶ LVA 703141 VU 3 – 5 ECTS

## Organisation

- ▶ LVA 703141    VU 3 – 5 ECTS    15:30–18:00 in HS 11

## Organisation

- ▶ LVA 703141    VU 3 – 5 ECTS    15:30–18:00 in HS 11
- ▶ <http://cl-informatik.uibk.ac.at/teaching/ss26/trs>

## Organisation

▶ LVA 703141 VU 3 – 5 ECTS 15:30–18:00 in HS 11

▶ <http://cl-informatik.uibk.ac.at/teaching/ss26/trs>

OLAT

## Organisation

- ▶ LVA 703141    VU 3 – 5 ECTS    15:30–18:00 in HS 11
- ▶ <http://cl-informatik.uibk.ac.at/teaching/ss26/trs>    OLAT

## Consultation Hours

Philipp Dablander	3M03	Wednesday	10:00–11:30
Aart Middeldorp	3M07	Monday	12:00–13:30

## Organisation

- ▶ LVA 703141 VU 3 – 5 ECTS 15:30–18:00 in HS 11
- ▶ <http://cl-informatik.uibk.ac.at/teaching/ss26/trs> OLAT

## Consultation Hours

Philipp Dablander	3M03	Wednesday	10:00–11:30
Aart Middeldorp	3M07	Monday	12:00–13:30

## Schedule

lecture 1	March 2	lecture 6	April 20	lecture 11	June 1
lecture 2	March 9	lecture 7	April 27	lecture 12	June 8
lecture 3	March 16	lecture 8	May 4	lecture 13	June 15
lecture 4	March 23	lecture 9	May 11	lecture 14	June 22
lecture 5	April 13	lecture 10	May 18		

## Organisation

- ▶ LVA 703141 VU 3 – 5 ECTS 15:30–18:00 in HS 11
- ▶ <http://cl-informatik.uibk.ac.at/teaching/ss26/trs> OLAT

## Consultation Hours

Philipp Dablander	3M03	Wednesday	10:00–11:30
Aart Middeldorp	3M07	Monday	12:00–13:30

## Schedule

lecture 1	March 2	lecture 6	April 20	lecture 11	June 1
lecture 2	March 9	lecture 7	April 27	lecture 12	June 8
lecture 3	March 16	lecture 8	May 4	lecture 13	June 15
lecture 4	March 23	lecture 9	May 11	lecture 14	June 22 (test)
lecture 5	April 13	lecture 10	May 18		

$$\text{score} = \min\left(\max\left(\frac{50}{69}(E + P) + \frac{1}{3}T + B, T + B\right), 100\right)$$

$$\text{score} = \min\left(\max\left(\frac{50}{69}(E + P) + \frac{1}{3}T + B, T + B\right), 100\right)$$

$E$ : points for solved **exercises** (at most 84)

$$\text{score} = \min\left(\max\left(\frac{50}{69}(E + P) + \frac{1}{3}T + B, T + B\right), 100\right)$$

$E$ : points for solved exercises (at most 84)

$P$ : points for **presentation** of solutions (at most 8)

$$\text{score} = \min\left(\max\left(\frac{50}{69}(E + P) + \frac{1}{3}T + B, T + B\right), 100\right)$$

$E$ : points for solved exercises (at most 84)

$P$ : points for presentation of solutions (at most 8)

$T$ : points for **test** (at most 100)

$$\text{score} = \min\left(\max\left(\frac{50}{69}(E + P) + \frac{1}{3}T + B, T + B\right), 100\right)$$

$E$ : points for solved exercises (at most 84)

$P$ : points for presentation of solutions (at most 8)

$T$ : points for test (at most 100)

$B$ : points for **bonus exercises** (at most 20)

$$\text{score} = \min\left(\max\left(\frac{50}{69}(E + P) + \frac{1}{3}T + B, T + B\right), 100\right)$$

$E$ : points for solved exercises (at most 84)

$P$ : points for presentation of solutions (at most 8)

$T$ : points for test (at most 100)

$B$ : points for bonus exercises (at most 20)

$$\text{grade} = \text{score} \in (-50) \rightarrow 5 \quad [50 - 63) \rightarrow 4 \quad [63 - 75) \rightarrow 3 \quad [75 - 88) \rightarrow 2 \quad [88 -) \rightarrow 1$$

$$\text{score} = \min\left(\max\left(\frac{50}{69}(E + P) + \frac{1}{3}T + B, T + B\right), 100\right)$$

$E$ : points for solved **exercises** (at most 84)

$P$ : points for presentation of solutions (at most 8)

$T$ : points for test (at most 100)

$B$ : points for **bonus exercises** (at most 20)

$$\text{grade} = \text{score} \in (-50) \rightarrow 5 \quad [50 - 63) \rightarrow 4 \quad [63 - 75) \rightarrow 3 \quad [75 - 88) \rightarrow 2 \quad [88 -) \rightarrow 1$$

- ▶ solved exercises must be marked and solutions (**PDF format**) must be uploaded in **OLAT** before **10 am on Monday**

$$\text{score} = \min\left(\max\left(\frac{50}{69}(E + P) + \frac{1}{3}T + B, T + B\right), 100\right)$$

$E$ : points for solved exercises (at most 84)

$P$ : points for **presentation** of solutions (at most 8)

$T$ : points for test (at most 100)

$B$ : points for bonus exercises (at most 20)

$$\text{grade} = \text{score} \in (-50) \rightarrow 5 \quad [50 - 63) \rightarrow 4 \quad [63 - 75) \rightarrow 3 \quad [75 - 88) \rightarrow 2 \quad [88 -) \rightarrow 1$$

- ▶ solved exercises must be marked and solutions (PDF format) must be uploaded in OLAT before 10 am on Monday
- ▶ presentations are optional

$$\text{score} = \min\left(\max\left(\frac{50}{69}(E + P) + \frac{1}{3}T + B, T + B\right), 100\right)$$

$E$ : points for solved exercises (at most 84)

$P$ : points for presentation of solutions (at most 8)

$T$ : points for **test** (at most 100)

$B$ : points for bonus exercises (at most 20)

$$\text{grade} = \text{score} \in (-50) \rightarrow 5 \quad [50 - 63) \rightarrow 4 \quad [63 - 75) \rightarrow 3 \quad [75 - 88) \rightarrow 2 \quad [88 -) \rightarrow 1$$

- ▶ solved exercises must be marked and solutions (PDF format) must be uploaded in OLAT before 10 am on Monday
- ▶ presentations are optional
- ▶ (optional) test on June 22

## Grading (cont'd)

- ▶ VU  $\implies$  attendance is mandatory

## Grading (cont'd)

- ▶ VU  $\implies$  attendance is mandatory
- ▶ unexcused attendance is allowed twice  $\implies$  zero points

## Grading (cont'd)

- ▶ VU  $\implies$  attendance is mandatory
- ▶ unexcused attendance is allowed twice  $\implies$  zero points
- ▶ dropping out without grade is possible until March 16

## Grading (cont'd)

- ▶ VU  $\implies$  attendance is mandatory
- ▶ unexcused attendance is allowed twice  $\implies$  zero points
- ▶ **dropping out without grade is possible until March 16**

## Presentations

- ▶ solutions will be presented in first part of every subsequent lecture

## Grading (cont'd)

- ▶ VU  $\implies$  attendance is mandatory
- ▶ unexcused attendance is allowed twice  $\implies$  zero points
- ▶ **dropping out without grade is possible until March 16**

## Presentations

- ▶ solutions will be presented in first part of every subsequent lecture
- ▶ scheduling of presentations in OLAT

## Grading (cont'd)

- ▶ VU  $\implies$  attendance is mandatory
- ▶ unexcused attendance is allowed twice  $\implies$  zero points
- ▶ **dropping out without grade is possible until March 16**


## Presentations

- ▶ solutions will be presented in first part of every subsequent lecture
- ▶ scheduling of presentations in OLAT

evaluation 25S


# Outline

1. Organisation
- 2. Examples**
3. Terms
4. Exercises
5. Further Reading






A colony of chameleons includes 20 red, 18 blue, and 16 green individuals. Whenever two chameleons of different color meet, each changes to the third color. Some time passes during which no chameleons are born or die nor do any enter or leave the colony. Is it possible that at the end of this period, all 54 chameleons are the same color?










A colony of chameleons includes 20 red, 18 blue, and 16 green individuals. Whenever two chameleons of different color meet, each changes to the third color. Some time passes during which no chameleons are born or die nor do any enter or leave the colony. Is it possible that at the end of this period, all 54 chameleons are the same color?









A colony of chameleons includes 20 red, 18 blue, and 16 green individuals. Whenever two chameleons of different color meet, each changes to the third color. Some time passes during which no chameleons are born or die nor do any enter or leave the colony. Is it possible that at the end of this period, all 54 chameleons are the same color?






A colony of chameleons includes 20 red, 18 blue, and 16 green individuals. Whenever two chameleons of different color meet, each changes to the third color. Some time passes during which no chameleons are born or die nor do any enter or leave the colony. Is it possible that at the end of this period, all 54 chameleons are the same color?









A colony of chameleons includes 20 red, 18 blue, and 16 green individuals. Whenever two chameleons of different color meet, each changes to the third color. Some time passes during which no chameleons are born or die nor do any enter or leave the colony. Is it possible that at the end of this period, all 54 chameleons are the same color?





A colony of chameleons includes 20 red, 18 blue, and 16 green individuals. Whenever two chameleons of different color meet, each changes to the third color. Some time passes during which no chameleons are born or die nor do any enter or leave the colony. Is it possible that at the end of this period, all 54 chameleons are the same color?



A team of genetic engineers decides to create cows that produce cola instead of milk. To that end they have to transform the DNA of the milk gene

TAGCTAGCTAGCT

in every fertilized egg into the cola gene

CTGACTGACT



A team of genetic engineers decides to create cows that produce cola instead of milk. To that end they have to transform the DNA of the milk gene

TAGCTAGCTAGCT

in every fertilized egg into the cola gene

CTGACTGACT



Techniques exist to perform the following DNA substitutions

TCAT ↔ T

GAG ↔ AG

CTC ↔ TC

AGTA ↔ A

TAT ↔ CT

A team of genetic engineers decides to create cows that produce cola instead of milk. To that end they have to transform the DNA of the milk gene

TAGCTAGCTAGCT

in every fertilized egg into the cola gene

CTGACTGACT



Techniques exist to perform the following DNA substitutions

TCAT ↔ T

GAG ↔ AG

CTC ↔ TC

AGTA ↔ A

TAT ↔ CT

Recently it has been discovered that the mad cow disease is caused by a retrovirus with the following DNA sequence **CTGCTACTGACT**. What now, if accidentally cows with this virus are created? According to the engineers there is little risk because this never happened in their experiments, but various action groups demand absolute assurance.

Coffee beans come in two kinds called black (●) and white (○). A two-player game starts with a random sequence of black and white beans. In a move, a player must take two adjacent beans and put back one bean, according to the following set of rules:

● ● → ○

○ ○ → ○

● ○ → ●

○ ● → ●

The player who puts the last black bean wins.



Coffee beans come in two kinds called black (●) and white (○). A two-player game starts with a random sequence of black and white beans. In a move, a player must take two adjacent beans and put back one bean, according to the following set of rules:

● ● → ○

○ ○ → ○

● ○ → ●

○ ● → ●

The player who puts the last black bean wins.

- 
- 
- 
- 
- 
- 
- 
- 
- 
- 
- 
- 
- 
- 
- 
- 



Coffee beans come in two kinds called black (●) and white (○). A two-player game starts with a random sequence of black and white beans. In a move, a player must take two adjacent beans and put back one bean, according to the following set of rules:


● ● → ○

○ ○ → ○

● ○ → ●

○ ● → ●

The player who puts the last black bean wins.



Coffee beans come in two kinds called black (●) and white (○). A two-player game starts with a random sequence of black and white beans. In a move, a player must take two adjacent beans and put back one bean, according to the following set of rules:


● ● → ○

○ ○ → ○

● ○ → ●

○ ● → ●

The player who puts the last black bean wins.



Coffee beans come in two kinds called black (●) and white (○). A two-player game starts with a random sequence of black and white beans. In a move, a player must take two adjacent beans and put back one bean, according to the following set of rules:


$$\bullet \bullet \rightarrow \circ$$

$$\circ \circ \rightarrow \bullet$$

$$\bullet \circ \rightarrow \bullet$$

$$\circ \bullet \rightarrow \bullet$$

The player who puts the last black bean wins.



Coffee beans come in two kinds called black (●) and white (○). A two-player game starts with a random sequence of black and white beans. In a move, a player must take two adjacent beans and put back one bean, according to the following set of rules:


● ● → ○

○ ○ → ○

● ○ → ●

○ ● → ●

The player who puts the last black bean wins.



Coffee beans come in two kinds called black (●) and white (○). A two-player game starts with a random sequence of black and white beans. In a move, a player must take two adjacent beans and put back one bean, according to the following set of rules:


● ● → ○

○ ○ → ○

● ○ → ●

○ ● → ●

The player who puts the last black bean wins.



Coffee beans come in two kinds called black (●) and white (○). A two-player game starts with a random sequence of black and white beans. In a move, a player must take two adjacent beans and put back one bean, according to the following set of rules:


$$\bullet \bullet \rightarrow \circ$$

$$\circ \circ \rightarrow \bullet$$

$$\bullet \circ \rightarrow \bullet$$

$$\circ \bullet \rightarrow \bullet$$

The player who puts the last black bean wins.



Coffee beans come in two kinds called black (●) and white (○). A two-player game starts with a random sequence of black and white beans. In a move, a player must take two adjacent beans and put back one bean, according to the following set of rules:


● ● → ○

○ ○ → ○

● ○ → ●

○ ● → ●

The player who puts the last black bean wins.



Coffee beans come in two kinds called black (●) and white (○). A two-player game starts with a random sequence of black and white beans. In a move, a player must take two adjacent beans and put back one bean, according to the following set of rules:


● ● → ○

○ ○ → ○

● ○ → ●

○ ● → ●

The player who puts the last black bean wins.



Coffee beans come in two kinds called black (●) and white (○). A two-player game starts with a random sequence of black and white beans. In a move, a player must take two adjacent beans and put back one bean, according to the following set of rules:


● ● → ○

○ ○ → ○

● ○ → ●

○ ● → ●

The player who puts the last black bean wins.



Coffee beans come in two kinds called black (●) and white (○). A two-player game starts with a random sequence of black and white beans. In a move, a player must take two adjacent beans and put back one bean, according to the following set of rules:


$$\bullet \bullet \rightarrow \circ$$

$$\circ \circ \rightarrow \circ$$

$$\bullet \circ \rightarrow \bullet$$

$$\circ \bullet \rightarrow \bullet$$

The player who puts the last black bean wins.



Coffee beans come in two kinds called black (●) and white (○). A two-player game starts with a random sequence of black and white beans. In a move, a player must take two adjacent beans and put back one bean, according to the following set of rules:


● ● → ○

○ ○ → ○

● ○ → ●

○ ● → ●

The player who puts the last black bean wins.



Coffee beans come in two kinds called black (●) and white (○). A two-player game starts with a random sequence of black and white beans. In a move, a player must take two adjacent beans and put back one bean, according to the following set of rules:


● ● → ○

○ ○ → ○

● ○ → ●

○ ● → ●

The player who puts the last black bean wins.



Coffee beans come in two kinds called black (●) and white (○). A two-player game starts with a random sequence of black and white beans. In a move, a player must take two adjacent beans and put back one bean, according to the following set of rules:


● ● → ○

○ ○ → ○

● ○ → ●

○ ● → ●

The player who puts the last black bean wins.



Coffee beans come in two kinds called black (●) and white (○). A two-player game starts with a random sequence of black and white beans. In a move, a player must take two adjacent beans and put back one bean, according to the following set of rules:


● ● → ○

○ ○ → ○

● ○ → ●

○ ● → ●

The player who puts the last black bean wins.



Coffee beans come in two kinds called black (●) and white (○). A two-player game starts with a random sequence of black and white beans. In a move, a player must take two adjacent beans and put back one bean, according to the following set of rules:


● ● → ○

○ ○ → ○

● ○ → ●

○ ● → ●

The player who puts the last black bean wins.



Coffee beans come in two kinds called black (●) and white (○). A two-player game starts with a random sequence of black and white beans. In a move, a player must take two adjacent beans and put back one bean, according to the following set of rules:


● ● → ○

○ ○ → ○

● ○ → ●

○ ● → ●

The player who puts the last black bean wins.




Does the game terminate?

Coffee beans come in two kinds called black (●) and white (○). A two-player game starts with a random sequence of black and white beans. In a move, a player must take two adjacent beans and put back one bean, according to the following set of rules:

● ● → ○      ○ ○ → ○      ● ○ → ●      ○ ● → ●

The player who puts the last black bean wins.




Does the game terminate? Does one of the players have a winning strategy?

Coffee beans come in two kinds called black (●) and white (○). A two-player game starts with a random sequence of black and white beans. In a move, a player must take two adjacent beans and put back one bean, according to the following set of rules:


● ● → ○      ○ ○ → ○      ● ○ → ●      ○ ● → ●

The player who puts the last black bean wins.




Does the game terminate? Does one of the players have a winning strategy? What if the rules are changed to ● ● → ○ ○ ○ ○      ○ ○ → ○      ● ○ → ○ ○ ○ ●      ○ ● → ● ?



The mythological monster **Hydra** is a dragon-like creature with multiple heads. Whenever **Hercules** in his fight chops off a head, more and more new heads can grow instead, since the beast gets increasingly angry. Hydra dies and Hercules wins if there are no heads left.



The mythological monster **Hydra** is a dragon-like creature with multiple heads. Whenever **Hercules** in his fight chops off a head, more and more new heads can grow instead, since the beast gets increasingly angry. Hydra dies and Hercules wins if there are no heads left. Here we model a Hydra as an unordered tree. If Hercules cuts off a leaf  $l$  that has a grandparent  $g$ , the branch from  $g$  to the parent of  $l$  gets multiplied, with the number of copies depending on the number of decapitations so far.





The mythological monster **Hydra** is a dragon-like creature with multiple heads. Whenever **Hercules** in his fight chops off a head, more and more new heads can grow instead, since the beast gets increasingly angry. Hydra dies and Hercules wins if there are no heads left. Here we model a Hydra as an unordered tree. If Hercules cuts off a leaf  $l$  that has a grandparent  $g$ , the branch from  $g$  to the parent of  $l$  gets multiplied, with the number of copies depending on the number of decapitations so far.





0

The mythological monster **Hydra** is a dragon-like creature with multiple heads. Whenever **Hercules** in his fight chops off a head, more and more new heads can grow instead, since the beast gets increasingly angry. Hydra dies and Hercules wins if there are no heads left. Here we model a Hydra as an unordered tree. If Hercules cuts off a leaf  $l$  that has a grandparent  $g$ , the branch from  $g$  to the parent of  $l$  gets multiplied, with the number of copies depending on the number of decapitations so far.





0



The mythological monster **Hydra** is a dragon-like creature with multiple heads. Whenever **Hercules** in his fight chops off a head, more and more new heads can grow instead, since the beast gets increasingly angry. Hydra dies and Hercules wins if there are no heads left. Here we model a Hydra as an unordered tree. If Hercules cuts off a leaf  $l$  that has a grandparent  $g$ , the branch from  $g$  to the parent of  $l$  gets multiplied, with the number of copies depending on the number of decapitations so far.





The mythological monster **Hydra** is a dragon-like creature with multiple heads. Whenever **Hercules** in his fight chops off a head, more and more new heads can grow instead, since the beast gets increasingly angry. Hydra dies and Hercules wins if there are no heads left. Here we model a Hydra as an unordered tree. If Hercules cuts off a leaf  $l$  that has a grandparent  $g$ , the branch from  $g$  to the parent of  $l$  gets multiplied, with the number of copies depending on the number of decapitations so far.




The mythological monster **Hydra** is a dragon-like creature with multiple heads. Whenever **Hercules** in his fight chops off a head, more and more new heads can grow instead, since the beast gets increasingly angry. Hydra dies and Hercules wins if there are no heads left. Here we model a Hydra as an unordered tree. If Hercules cuts off a leaf  $l$  that has a grandparent  $g$ , the branch from  $g$  to the parent of  $l$  gets multiplied, with the number of copies depending on the number of decapitations so far.





The mythological monster **Hydra** is a dragon-like creature with multiple heads. Whenever **Hercules** in his fight chops off a head, more and more new heads can grow instead, since the beast gets increasingly angry. Hydra dies and Hercules wins if there are no heads left. Here we model a Hydra as an unordered tree. If Hercules cuts off a leaf  $l$  that has a grandparent  $g$ , the branch from  $g$  to the parent of  $l$  gets multiplied, with the number of copies depending on the number of decapitations so far.




The mythological monster **Hydra** is a dragon-like creature with multiple heads. Whenever **Hercules** in his fight chops off a head, more and more new heads can grow instead, since the beast gets increasingly angry. Hydra dies and Hercules wins if there are no heads left. Here we model a Hydra as an unordered tree. If Hercules cuts off a leaf  $l$  that has a grandparent  $g$ , the branch from  $g$  to the parent of  $l$  gets multiplied, with the number of copies depending on the number of decapitations so far.




The mythological monster **Hydra** is a dragon-like creature with multiple heads. Whenever **Hercules** in his fight chops off a head, more and more new heads can grow instead, since the beast gets increasingly angry. Hydra dies and Hercules wins if there are no heads left. Here we model a Hydra as an unordered tree. If Hercules cuts off a leaf  $l$  that has a grandparent  $g$ , the branch from  $g$  to the parent of  $l$  gets multiplied, with the number of copies depending on the number of decapitations so far.




The mythological monster **Hydra** is a dragon-like creature with multiple heads. Whenever **Hercules** in his fight chops off a head, more and more new heads can grow instead, since the beast gets increasingly angry. Hydra dies and Hercules wins if there are no heads left. Here we model a Hydra as an unordered tree. If Hercules cuts off a leaf  $l$  that has a grandparent  $g$ , the branch from  $g$  to the parent of  $l$  gets multiplied, with the number of copies depending on the number of decapitations so far.




0




1



2




3




4


The mythological monster **Hydra** is a dragon-like creature with multiple heads. Whenever **Hercules** in his fight chops off a head, more and more new heads can grow instead, since the beast gets increasingly angry. Hydra dies and Hercules wins if there are no heads left. Here we model a Hydra as an unordered tree. If Hercules cuts off a leaf  $l$  that has a grandparent  $g$ , the branch from  $g$  to the parent of  $l$  gets multiplied, with the number of copies depending on the number of decapitations so far.




0




1



2



3



4

Can Hercules win the battle?

## Example (Addition on Natural Numbers in Unary Notation)

signature      0 (constant)    s (unary)    a (binary)

## Example (Addition on Natural Numbers in Unary Notation)

signature      0 (constant)    s (unary)    a (binary)

terms            0

## Example (Addition on Natural Numbers in Unary Notation)

signature      0 (constant)    s (unary)    a (binary)

terms            0    s(0)

## Example (Addition on Natural Numbers in Unary Notation)

signature      0 (constant)    s (unary)    a (binary)

terms            0    s(0)    s(s(0))

## Example (Addition on Natural Numbers in Unary Notation)

signature      0 (constant)    s (unary)    a (binary)

terms            0    s(0)    s(s(0))    a(s(0),s(0))

## Example (Addition on Natural Numbers in Unary Notation)

signature      0 (constant)   s (unary)   a (binary)

terms            0   s(0)   s(s(0))   a(s(0),s(0))   s(a(s(0),s(a(0,s(s(0))))))

## Example (Addition on Natural Numbers in Unary Notation)

signature      0 (constant)    s (unary)    a (binary)

terms            0    s(0)    s(s(0))    a(s(0),s(0))    s(a(s(0),s(a(0,s(s(0))))))

rewrite rules

$$a(0, y) \rightarrow y$$

$$a(s(x), y) \rightarrow s(a(x, y))$$

## Example (Addition on Natural Numbers in Unary Notation)

signature      0 (constant)    s (unary)    a (binary)

terms            0    s(0)    s(s(0))    a(s(0),s(0))    s(a(s(0),s(a(0,s(s(0))))))

rewrite rules      a(0,y) → y                      variables

a(s(x),y) → s(a(x,y))

## Example (Addition on Natural Numbers in Unary Notation)

signature      0 (constant)    s (unary)    a (binary)

terms            0    s(0)    s(s(0))    a(s(0),s(0))    s(a(s(0),s(a(0,s(s(0))))))

rewrite rules       $a(0,y) \rightarrow y$   
 $a(s(x),y) \rightarrow s(a(x,y))$

rewriting         $a(s(s(s(0))),s(s(0)))$

## Example (Addition on Natural Numbers in Unary Notation)

signature      0 (constant)    s (unary)    a (binary)

terms            0    s(0)    s(s(0))    a(s(0),s(0))    s(a(s(0),s(a(0,s(s(0))))))

rewrite rules      a(0,y)  $\rightarrow$  y

a(s(x),y)  $\rightarrow$  s(a(x,y))

rewriting        a(s(s(s(0))),s(s(0)))

## Example (Addition on Natural Numbers in Unary Notation)

signature      0 (constant)    s (unary)    a (binary)

terms            0    s(0)    s(s(0))    a(s(0),s(0))    s(a(s(0),s(a(0,s(s(0))))))

rewrite rules      a(0,y)  $\rightarrow$  y

a(s(x),y)  $\rightarrow$  s(a(x,y))      x  $\mapsto$  s(s(0))    y  $\mapsto$  s(s(0))

rewriting        a(s(s(s(0))),s(s(0)))

## Example (Addition on Natural Numbers in Unary Notation)

signature      0 (constant)    s (unary)    a (binary)

terms            0    s(0)    s(s(0))    a(s(0),s(0))    s(a(s(0),s(a(0,s(s(0))))))

rewrite rules      a(0,y) → y

a(s(x),y) → s(a(x,y))      x ↦ s(s(0))    y ↦ s(s(0))

rewriting        a(s(s(s(0))),s(s(0))) → s(a(s(s(0)),s(s(0))))

## Example (Addition on Natural Numbers in Unary Notation)

signature      0 (constant)    s (unary)    a (binary)

terms            0    s(0)    s(s(0))    a(s(0),s(0))    s(a(s(0),s(a(0,s(s(0))))))

rewrite rules      a(0,y) → y

a(s(x),y) → s(a(x,y))      x ↦ s(0)    y ↦ s(s(0))

rewriting          a(s(s(s(0))),s(s(0))) → s(a(s(s(0)),s(s(0))))

## Example (Addition on Natural Numbers in Unary Notation)

signature      0 (constant)    s (unary)    a (binary)

terms            0    s(0)    s(s(0))    a(s(0),s(0))    s(a(s(0),s(a(0,s(s(0))))))

rewrite rules      a(0,y)  $\rightarrow$  y

a(s(x),y)  $\rightarrow$  s(a(x,y))      x  $\mapsto$  s(0)    y  $\mapsto$  s(s(0))

rewriting          a(s(s(s(0))),s(s(0)))  $\rightarrow$  s(a(s(s(0)),s(s(0))))  
 $\rightarrow$  s(s(a(s(0),s(s(0)))))

## Example (Addition on Natural Numbers in Unary Notation)

signature      0 (constant)    s (unary)    a (binary)

terms            0    s(0)    s(s(0))    a(s(0),s(0))    s(a(s(0),s(a(0,s(s(0))))))

rewrite rules      a(0,y)  $\rightarrow$  y

a(s(x),y)  $\rightarrow$  s(a(x,y))      x  $\mapsto$  0    y  $\mapsto$  s(s(0))

rewriting          a(s(s(s(0))),s(s(0)))  $\rightarrow$  s(a(s(s(0)),s(s(0))))  
 $\rightarrow$  s(s(a(s(0),s(s(0)))))



## Example (Addition on Natural Numbers in Unary Notation)

signature      0 (constant)    s (unary)    a (binary)

terms            0    s(0)    s(s(0))    a(s(0),s(0))    s(a(s(0),s(a(0,s(s(0))))))

rewrite rules       $a(0,y) \rightarrow y$                        $y \mapsto s(s(0))$   
 $a(s(x),y) \rightarrow s(a(x,y))$

rewriting           $a(s(s(s(0))),s(s(0))) \rightarrow s(a(s(s(0)),s(s(0))))$   
                          $\rightarrow s(s(a(s(0),s(s(0)))))$   
                          $\rightarrow s(s(s(a(0,s(s(0))))))$

## Example (Addition on Natural Numbers in Unary Notation)

signature      0 (constant)    s (unary)    a (binary)

terms            0    s(0)    s(s(0))    a(s(0),s(0))    s(a(s(0),s(a(0,s(s(0))))))

rewrite rules       $a(0, y) \rightarrow y$                        $y \mapsto s(s(0))$   
 $a(s(x), y) \rightarrow s(a(x, y))$

rewriting           $a(s(s(s(0))), s(s(0))) \rightarrow s(a(s(s(0)), s(s(0))))$   
                                  $\rightarrow s(s(a(s(0), s(s(0)))))$   
                                  $\rightarrow s(s(s(a(0, s(s(0))))))$   
                                  $\rightarrow s(s(s(s(s(0)))))$

## Example (Addition on Natural Numbers in Unary Notation)

signature      0 (constant)    s (unary)    a (binary)

terms            0    s(0)    s(s(0))    a(s(0),s(0))    s(a(s(0),s(a(0,s(s(0))))))

rewrite rules      a(0,y) → y

a(s(x),y) → s(a(x,y))

rewriting          a(s(s(s(0))),s(s(0))) → s(a(s(s(0)),s(s(0))))

→ s(s(a(s(0),s(s(0)))))

→ s(s(s(a(0,s(s(0))))))

→ s(s(s(s(0))))

normal form

## Example (Addition on Natural Numbers in Decimal Notation)

signature      0 1 ... 9 (constants) + : (binary, infix)

## Example (Addition on Natural Numbers in Decimal Notation)

signature      0 1 ... 9 (constants) + : (binary, infix)

terms            1 + 3   2 + (7 : 3)   (2 : (3 : x)) + ((1 + 7) : 2)

## Example (Addition on Natural Numbers in Decimal Notation)

signature       $0\ 1\ \dots\ 9$  (constants)     $+$  : (binary, infix)

terms             $1 + 3$      $2 + (7 : 3)$      $(2 : (3 : x)) + ((1 + 7) : 2)$

rewrite rules

$0 + 0 \rightarrow 0$	$1 + 0 \rightarrow 1$	$\dots$	$9 + 0 \rightarrow 9$	$0 : x \rightarrow x$
$0 + 1 \rightarrow 1$	$1 + 1 \rightarrow 2$	$\dots$	$9 + 1 \rightarrow 1 : 0$	$x + (y : z) \rightarrow y : (x + z)$
$0 + 2 \rightarrow 2$	$1 + 2 \rightarrow 3$	$\dots$	$9 + 2 \rightarrow 1 : 1$	$(x : y) + z \rightarrow x : (y + z)$
$0 + 3 \rightarrow 3$	$1 + 3 \rightarrow 4$	$\dots$	$9 + 3 \rightarrow 1 : 2$	$x : (y : z) \rightarrow (x + y) : z$
$0 + 4 \rightarrow 4$	$1 + 4 \rightarrow 5$	$\dots$	$9 + 4 \rightarrow 1 : 3$	
$0 + 5 \rightarrow 5$	$1 + 5 \rightarrow 6$	$\dots$	$9 + 5 \rightarrow 1 : 4$	
$0 + 6 \rightarrow 6$	$1 + 6 \rightarrow 7$	$\dots$	$9 + 6 \rightarrow 1 : 5$	
$0 + 7 \rightarrow 7$	$1 + 7 \rightarrow 8$	$\dots$	$9 + 7 \rightarrow 1 : 6$	
$0 + 8 \rightarrow 8$	$1 + 8 \rightarrow 9$	$\dots$	$9 + 8 \rightarrow 1 : 7$	
$0 + 9 \rightarrow 9$	$1 + 9 \rightarrow 1 : 0$	$\dots$	$9 + 9 \rightarrow 1 : 8$	

## Example (Addition on Natural Numbers in Decimal Notation)

signature       $0 \ 1 \ \dots \ 9$  (constants)     $+$      $:$  (binary, infix)

terms             $1 + 3 \quad 2 + (7 : 3) \quad (2 : (3 : x)) + ((1 + 7) : 2)$

rewrite rules     $0 + 0 \rightarrow 0 \quad 1 + 0 \rightarrow 1 \quad \dots \quad 9 + 0 \rightarrow 9 \quad 0 : x \rightarrow x$   
 $0 + 1 \rightarrow 1 \quad 1 + 1 \rightarrow 2 \quad \dots \quad 9 + 1 \rightarrow 1 : 0 \quad x + (y : z) \rightarrow y : (x + z)$   
 $0 + 2 \rightarrow 2 \quad 1 + 2 \rightarrow 3 \quad \dots \quad 9 + 2 \rightarrow 1 : 1 \quad (x : y) + z \rightarrow x : (y + z)$   
 $0 + 3 \rightarrow 3 \quad 1 + 3 \rightarrow 4 \quad \dots \quad 9 + 3 \rightarrow 1 : 2 \quad x : (y : z) \rightarrow (x + y) : z$   
 $0 + 4 \rightarrow 4 \quad 1 + 4 \rightarrow 5 \quad \dots \quad 9 + 4 \rightarrow 1 : 3$   
 $0 + 5 \rightarrow 5 \quad 1 + 5 \rightarrow 6 \quad \dots \quad 9 + 5 \rightarrow 1 : 4$   
 $0 + 6 \rightarrow 6 \quad 1 + 6 \rightarrow 7 \quad \dots \quad 9 + 6 \rightarrow 1 : 5$   
 $0 + 7 \rightarrow 7 \quad 1 + 7 \rightarrow 8 \quad \dots \quad 9 + 7 \rightarrow 1 : 6$   
 $0 + 8 \rightarrow 8 \quad 1 + 8 \rightarrow 9 \quad \dots \quad 9 + 8 \rightarrow 1 : 7$   
 $0 + 9 \rightarrow 9 \quad 1 + 9 \rightarrow 1 : 0 \quad \dots \quad 9 + 9 \rightarrow 1 : 8$

rewriting       $(2 : 3) + (7 : 7)$

## Example (Addition on Natural Numbers in Decimal Notation)

signature       $0 \ 1 \ \dots \ 9$  (constants)     $+$  : (binary, infix)

terms             $1 + 3 \quad 2 + (7 : 3) \quad (2 : (3 : x)) + ((1 + 7) : 2)$

rewrite rules

$0 + 0 \rightarrow 0$	$1 + 0 \rightarrow 1$	$\dots$	$9 + 0 \rightarrow 9$	$0 : x \rightarrow x$
$0 + 1 \rightarrow 1$	$1 + 1 \rightarrow 2$	$\dots$	$9 + 1 \rightarrow 1 : 0$	$x + (y : z) \rightarrow y : (x + z)$
$0 + 2 \rightarrow 2$	$1 + 2 \rightarrow 3$	$\dots$	$9 + 2 \rightarrow 1 : 1$	$(x : y) + z \rightarrow x : (y + z)$
$0 + 3 \rightarrow 3$	$1 + 3 \rightarrow 4$	$\dots$	$9 + 3 \rightarrow 1 : 2$	$x : (y : z) \rightarrow (x + y) : z$
$0 + 4 \rightarrow 4$	$1 + 4 \rightarrow 5$	$\dots$	$9 + 4 \rightarrow 1 : 3$	
$0 + 5 \rightarrow 5$	$1 + 5 \rightarrow 6$	$\dots$	$9 + 5 \rightarrow 1 : 4$	
$0 + 6 \rightarrow 6$	$1 + 6 \rightarrow 7$	$\dots$	$9 + 6 \rightarrow 1 : 5$	
$0 + 7 \rightarrow 7$	$1 + 7 \rightarrow 8$	$\dots$	$9 + 7 \rightarrow 1 : 6$	
$0 + 8 \rightarrow 8$	$1 + 8 \rightarrow 9$	$\dots$	$9 + 8 \rightarrow 1 : 7$	
$0 + 9 \rightarrow 9$	$1 + 9 \rightarrow 1 : 0$	$\dots$	$9 + 9 \rightarrow 1 : 8$	

rewriting       $(2 : 3) + (7 : 7) \quad x \mapsto 2 \quad y \mapsto 3 \quad z \mapsto 7 : 7$

## Example (Addition on Natural Numbers in Decimal Notation)

signature  $0\ 1\ \dots\ 9$  (constants)  $+$   $:$  (binary, infix)

terms  $1 + 3$   $2 + (7 : 3)$   $(2 : (3 : x)) + ((1 + 7) : 2)$

rewrite rules

$0 + 0 \rightarrow 0$	$1 + 0 \rightarrow 1$	$\dots$	$9 + 0 \rightarrow 9$	$0 : x \rightarrow x$
$0 + 1 \rightarrow 1$	$1 + 1 \rightarrow 2$	$\dots$	$9 + 1 \rightarrow 1 : 0$	$x + (y : z) \rightarrow y : (x + z)$
$0 + 2 \rightarrow 2$	$1 + 2 \rightarrow 3$	$\dots$	$9 + 2 \rightarrow 1 : 1$	$(x : y) + z \rightarrow x : (y + z)$
$0 + 3 \rightarrow 3$	$1 + 3 \rightarrow 4$	$\dots$	$9 + 3 \rightarrow 1 : 2$	$x : (y : z) \rightarrow (x + y) : z$
$0 + 4 \rightarrow 4$	$1 + 4 \rightarrow 5$	$\dots$	$9 + 4 \rightarrow 1 : 3$	
$0 + 5 \rightarrow 5$	$1 + 5 \rightarrow 6$	$\dots$	$9 + 5 \rightarrow 1 : 4$	
$0 + 6 \rightarrow 6$	$1 + 6 \rightarrow 7$	$\dots$	$9 + 6 \rightarrow 1 : 5$	
$0 + 7 \rightarrow 7$	$1 + 7 \rightarrow 8$	$\dots$	$9 + 7 \rightarrow 1 : 6$	
$0 + 8 \rightarrow 8$	$1 + 8 \rightarrow 9$	$\dots$	$9 + 8 \rightarrow 1 : 7$	
$0 + 9 \rightarrow 9$	$1 + 9 \rightarrow 1 : 0$	$\dots$	$9 + 9 \rightarrow 1 : 8$	

rewriting  $(2 : 3) + (7 : 7)$   $x \mapsto 2 : 3$   $y \mapsto 7$   $z \mapsto 7$

## Example (Addition on Natural Numbers in Decimal Notation)

signature      0 1 ... 9 (constants)    + : (binary, infix)

terms            1 + 3    2 + (7 : 3)    (2 : (3 : x)) + ((1 + 7) : 2)

rewrite rules     $0 + 0 \rightarrow 0$      $1 + 0 \rightarrow 1$     ...     $9 + 0 \rightarrow 9$                      $0 : x \rightarrow x$   
 $0 + 1 \rightarrow 1$      $1 + 1 \rightarrow 2$     ...     $9 + 1 \rightarrow 1 : 0$      $x + (y : z) \rightarrow y : (x + z)$   
 $0 + 2 \rightarrow 2$      $1 + 2 \rightarrow 3$     ...     $9 + 2 \rightarrow 1 : 1$      $(x : y) + z \rightarrow x : (y + z)$   
 $0 + 3 \rightarrow 3$      $1 + 3 \rightarrow 4$     ...     $9 + 3 \rightarrow 1 : 2$      $x : (y : z) \rightarrow (x + y) : z$   
 $0 + 4 \rightarrow 4$      $1 + 4 \rightarrow 5$     ...     $9 + 4 \rightarrow 1 : 3$   
 $0 + 5 \rightarrow 5$      $1 + 5 \rightarrow 6$     ...     $9 + 5 \rightarrow 1 : 4$   
 $0 + 6 \rightarrow 6$      $1 + 6 \rightarrow 7$     ...     $9 + 6 \rightarrow 1 : 5$   
 $0 + 7 \rightarrow 7$      $1 + 7 \rightarrow 8$     ...     $9 + 7 \rightarrow 1 : 6$   
 $0 + 8 \rightarrow 8$      $1 + 8 \rightarrow 9$     ...     $9 + 8 \rightarrow 1 : 7$   
 $0 + 9 \rightarrow 9$      $1 + 9 \rightarrow 1 : 0$     ...     $9 + 9 \rightarrow 1 : 8$

rewriting         $(2 : 3) + (7 : 7) \rightarrow 7 : ((2 : 3) + 7)$

## Example (Addition on Natural Numbers in Decimal Notation)

signature  $0\ 1\ \dots\ 9$  (constants)  $+$   $:$  (binary, infix)

terms  $1 + 3$   $2 + (7 : 3)$   $(2 : (3 : x)) + ((1 + 7) : 2)$

rewrite rules

$0 + 0 \rightarrow 0$	$1 + 0 \rightarrow 1$	$\dots$	$9 + 0 \rightarrow 9$	$0 : x \rightarrow x$
$0 + 1 \rightarrow 1$	$1 + 1 \rightarrow 2$	$\dots$	$9 + 1 \rightarrow 1 : 0$	$x + (y : z) \rightarrow y : (x + z)$
$0 + 2 \rightarrow 2$	$1 + 2 \rightarrow 3$	$\dots$	$9 + 2 \rightarrow 1 : 1$	$(x : y) + z \rightarrow x : (y + z)$
$0 + 3 \rightarrow 3$	$1 + 3 \rightarrow 4$	$\dots$	$9 + 3 \rightarrow 1 : 2$	$x : (y : z) \rightarrow (x + y) : z$
$0 + 4 \rightarrow 4$	$1 + 4 \rightarrow 5$	$\dots$	$9 + 4 \rightarrow 1 : 3$	
$0 + 5 \rightarrow 5$	$1 + 5 \rightarrow 6$	$\dots$	$9 + 5 \rightarrow 1 : 4$	
$0 + 6 \rightarrow 6$	$1 + 6 \rightarrow 7$	$\dots$	$9 + 6 \rightarrow 1 : 5$	
$0 + 7 \rightarrow 7$	$1 + 7 \rightarrow 8$	$\dots$	$9 + 7 \rightarrow 1 : 6$	
$0 + 8 \rightarrow 8$	$1 + 8 \rightarrow 9$	$\dots$	$9 + 8 \rightarrow 1 : 7$	
$0 + 9 \rightarrow 9$	$1 + 9 \rightarrow 1 : 0$	$\dots$	$9 + 9 \rightarrow 1 : 8$	

rewriting  $(2 : 3) + (7 : 7) \rightarrow 7 : ((2 : 3) + 7)$

## Example (Addition on Natural Numbers in Decimal Notation)

signature       $0\ 1\ \dots\ 9$  (constants)     $+$  : (binary, infix)

terms             $1 + 3$      $2 + (7 : 3)$      $(2 : (3 : x)) + ((1 + 7) : 2)$

rewrite rules     $0 + 0 \rightarrow 0$      $1 + 0 \rightarrow 1$      $\dots$      $9 + 0 \rightarrow 9$                      $0 : x \rightarrow x$   
 $0 + 1 \rightarrow 1$      $1 + 1 \rightarrow 2$      $\dots$      $9 + 1 \rightarrow 1 : 0$      $x + (y : z) \rightarrow y : (x + z)$   
 $0 + 2 \rightarrow 2$      $1 + 2 \rightarrow 3$      $\dots$      $9 + 2 \rightarrow 1 : 1$      $(x : y) + z \rightarrow x : (y + z)$   
 $0 + 3 \rightarrow 3$      $1 + 3 \rightarrow 4$      $\dots$      $9 + 3 \rightarrow 1 : 2$      $x : (y : z) \rightarrow (x + y) : z$   
 $0 + 4 \rightarrow 4$      $1 + 4 \rightarrow 5$      $\dots$      $9 + 4 \rightarrow 1 : 3$   
 $0 + 5 \rightarrow 5$      $1 + 5 \rightarrow 6$      $\dots$      $9 + 5 \rightarrow 1 : 4$   
 $0 + 6 \rightarrow 6$      $1 + 6 \rightarrow 7$      $\dots$      $9 + 6 \rightarrow 1 : 5$   
 $0 + 7 \rightarrow 7$      $1 + 7 \rightarrow 8$      $\dots$      $9 + 7 \rightarrow 1 : 6$   
 $0 + 8 \rightarrow 8$      $1 + 8 \rightarrow 9$      $\dots$      $9 + 8 \rightarrow 1 : 7$   
 $0 + 9 \rightarrow 9$      $1 + 9 \rightarrow 1 : 0$      $\dots$      $9 + 9 \rightarrow 1 : 8$

rewriting         $(2 : 3) + (7 : 7) \rightarrow^* 7 : (2 : (3 + 7))$

## Example (Addition on Natural Numbers in Decimal Notation)

signature  $0\ 1\ \dots\ 9$  (constants)  $+$   $:$  (binary, infix)

terms  $1 + 3$   $2 + (7 : 3)$   $(2 : (3 : x)) + ((1 + 7) : 2)$

rewrite rules

$0 + 0 \rightarrow 0$	$1 + 0 \rightarrow 1$	$\dots$	$9 + 0 \rightarrow 9$	$0 : x \rightarrow x$
$0 + 1 \rightarrow 1$	$1 + 1 \rightarrow 2$	$\dots$	$9 + 1 \rightarrow 1 : 0$	$x + (y : z) \rightarrow y : (x + z)$
$0 + 2 \rightarrow 2$	$1 + 2 \rightarrow 3$	$\dots$	$9 + 2 \rightarrow 1 : 1$	$(x : y) + z \rightarrow x : (y + z)$
$0 + 3 \rightarrow 3$	$1 + 3 \rightarrow 4$	$\dots$	$9 + 3 \rightarrow 1 : 2$	$x : (y : z) \rightarrow (x + y) : z$
$0 + 4 \rightarrow 4$	$1 + 4 \rightarrow 5$	$\dots$	$9 + 4 \rightarrow 1 : 3$	
$0 + 5 \rightarrow 5$	$1 + 5 \rightarrow 6$	$\dots$	$9 + 5 \rightarrow 1 : 4$	
$0 + 6 \rightarrow 6$	$1 + 6 \rightarrow 7$	$\dots$	$9 + 6 \rightarrow 1 : 5$	
$0 + 7 \rightarrow 7$	$1 + 7 \rightarrow 8$	$\dots$	$9 + 7 \rightarrow 1 : 6$	
$0 + 8 \rightarrow 8$	$1 + 8 \rightarrow 9$	$\dots$	$9 + 8 \rightarrow 1 : 7$	
$0 + 9 \rightarrow 9$	$1 + 9 \rightarrow 1 : 0$	$\dots$	$9 + 9 \rightarrow 1 : 8$	

rewriting  $(2 : 3) + (7 : 7) \rightarrow^* 7 : (2 : (3 + 7))$

## Example (Addition on Natural Numbers in Decimal Notation)

signature       $0\ 1\ \dots\ 9$  (constants)     $+$  : (binary, infix)

terms             $1 + 3$      $2 + (7 : 3)$      $(2 : (3 : x)) + ((1 + 7) : 2)$

rewrite rules     $0 + 0 \rightarrow 0$      $1 + 0 \rightarrow 1$      $\dots$      $9 + 0 \rightarrow 9$                      $0 : x \rightarrow x$   
 $0 + 1 \rightarrow 1$      $1 + 1 \rightarrow 2$      $\dots$      $9 + 1 \rightarrow 1 : 0$      $x + (y : z) \rightarrow y : (x + z)$   
 $0 + 2 \rightarrow 2$      $1 + 2 \rightarrow 3$      $\dots$      $9 + 2 \rightarrow 1 : 1$      $(x : y) + z \rightarrow x : (y + z)$   
 $0 + 3 \rightarrow 3$      $1 + 3 \rightarrow 4$      $\dots$      $9 + 3 \rightarrow 1 : 2$      $x : (y : z) \rightarrow (x + y) : z$   
 $0 + 4 \rightarrow 4$      $1 + 4 \rightarrow 5$      $\dots$      $9 + 4 \rightarrow 1 : 3$   
 $0 + 5 \rightarrow 5$      $1 + 5 \rightarrow 6$      $\dots$      $9 + 5 \rightarrow 1 : 4$   
 $0 + 6 \rightarrow 6$      $1 + 6 \rightarrow 7$      $\dots$      $9 + 6 \rightarrow 1 : 5$   
 $0 + 7 \rightarrow 7$      $1 + 7 \rightarrow 8$      $\dots$      $9 + 7 \rightarrow 1 : 6$   
 $0 + 8 \rightarrow 8$      $1 + 8 \rightarrow 9$      $\dots$      $9 + 8 \rightarrow 1 : 7$   
 $0 + 9 \rightarrow 9$      $1 + 9 \rightarrow 1 : 0$      $\dots$      $9 + 9 \rightarrow 1 : 8$

rewriting         $(2 : 3) + (7 : 7) \rightarrow^* 7 : (2 : (1 : 0))$

## Example (Addition on Natural Numbers in Decimal Notation)

signature      0 1 ... 9 (constants) + : (binary, infix)

terms             $1 + 3$     $2 + (7 : 3)$     $(2 : (3 : x)) + ((1 + 7) : 2)$

rewrite rules

$0 + 0 \rightarrow 0$	$1 + 0 \rightarrow 1$	$\dots$	$9 + 0 \rightarrow 9$	$0 : x \rightarrow x$
$0 + 1 \rightarrow 1$	$1 + 1 \rightarrow 2$	$\dots$	$9 + 1 \rightarrow 1 : 0$	$x + (y : z) \rightarrow y : (x + z)$
$0 + 2 \rightarrow 2$	$1 + 2 \rightarrow 3$	$\dots$	$9 + 2 \rightarrow 1 : 1$	$(x : y) + z \rightarrow x : (y + z)$
$0 + 3 \rightarrow 3$	$1 + 3 \rightarrow 4$	$\dots$	$9 + 3 \rightarrow 1 : 2$	$x : (y : z) \rightarrow (x + y) : z$
$0 + 4 \rightarrow 4$	$1 + 4 \rightarrow 5$	$\dots$	$9 + 4 \rightarrow 1 : 3$	
$0 + 5 \rightarrow 5$	$1 + 5 \rightarrow 6$	$\dots$	$9 + 5 \rightarrow 1 : 4$	
$0 + 6 \rightarrow 6$	$1 + 6 \rightarrow 7$	$\dots$	$9 + 6 \rightarrow 1 : 5$	
$0 + 7 \rightarrow 7$	$1 + 7 \rightarrow 8$	$\dots$	$9 + 7 \rightarrow 1 : 6$	
$0 + 8 \rightarrow 8$	$1 + 8 \rightarrow 9$	$\dots$	$9 + 8 \rightarrow 1 : 7$	
$0 + 9 \rightarrow 9$	$1 + 9 \rightarrow 1 : 0$	$\dots$	$9 + 9 \rightarrow 1 : 8$	

rewriting       $(2 : 3) + (7 : 7) \rightarrow^* 7 : (2 : (1 : 0))$

## Example (Addition on Natural Numbers in Decimal Notation)

signature  $0 \ 1 \ \dots \ 9$  (constants)  $+$   $:$  (binary, infix)

terms  $1 + 3 \quad 2 + (7 : 3) \quad (2 : (3 : x)) + ((1 + 7) : 2)$

rewrite rules

$0 + 0 \rightarrow 0$	$1 + 0 \rightarrow 1$	$\dots$	$9 + 0 \rightarrow 9$	$0 : x \rightarrow x$
$0 + 1 \rightarrow 1$	$1 + 1 \rightarrow 2$	$\dots$	$9 + 1 \rightarrow 1 : 0$	$x + (y : z) \rightarrow y : (x + z)$
$0 + 2 \rightarrow 2$	$1 + 2 \rightarrow 3$	$\dots$	$9 + 2 \rightarrow 1 : 1$	$(x : y) + z \rightarrow x : (y + z)$
$0 + 3 \rightarrow 3$	$1 + 3 \rightarrow 4$	$\dots$	$9 + 3 \rightarrow 1 : 2$	$x : (y : z) \rightarrow (x + y) : z$
$0 + 4 \rightarrow 4$	$1 + 4 \rightarrow 5$	$\dots$	$9 + 4 \rightarrow 1 : 3$	
$0 + 5 \rightarrow 5$	$1 + 5 \rightarrow 6$	$\dots$	$9 + 5 \rightarrow 1 : 4$	
$0 + 6 \rightarrow 6$	$1 + 6 \rightarrow 7$	$\dots$	$9 + 6 \rightarrow 1 : 5$	
$0 + 7 \rightarrow 7$	$1 + 7 \rightarrow 8$	$\dots$	$9 + 7 \rightarrow 1 : 6$	
$0 + 8 \rightarrow 8$	$1 + 8 \rightarrow 9$	$\dots$	$9 + 8 \rightarrow 1 : 7$	
$0 + 9 \rightarrow 9$	$1 + 9 \rightarrow 1 : 0$	$\dots$	$9 + 9 \rightarrow 1 : 8$	

rewriting  $(2 : 3) + (7 : 7) \rightarrow^* 7 : ((2 + 1) : 0)$

## Example (Addition on Natural Numbers in Decimal Notation)

signature  $0\ 1\ \dots\ 9$  (constants)  $+$   $:$  (binary, infix)

terms  $1 + 3$   $2 + (7 : 3)$   $(2 : (3 : x)) + ((1 + 7) : 2)$

rewrite rules

$0 + 0 \rightarrow 0$	$1 + 0 \rightarrow 1$	$\dots$	$9 + 0 \rightarrow 9$	$0 : x \rightarrow x$
$0 + 1 \rightarrow 1$	$1 + 1 \rightarrow 2$	$\dots$	$9 + 1 \rightarrow 1 : 0$	$x + (y : z) \rightarrow y : (x + z)$
$0 + 2 \rightarrow 2$	$1 + 2 \rightarrow 3$	$\dots$	$9 + 2 \rightarrow 1 : 1$	$(x : y) + z \rightarrow x : (y + z)$
$0 + 3 \rightarrow 3$	$1 + 3 \rightarrow 4$	$\dots$	$9 + 3 \rightarrow 1 : 2$	$x : (y : z) \rightarrow (x + y) : z$
$0 + 4 \rightarrow 4$	$1 + 4 \rightarrow 5$	$\dots$	$9 + 4 \rightarrow 1 : 3$	
$0 + 5 \rightarrow 5$	$1 + 5 \rightarrow 6$	$\dots$	$9 + 5 \rightarrow 1 : 4$	
$0 + 6 \rightarrow 6$	$1 + 6 \rightarrow 7$	$\dots$	$9 + 6 \rightarrow 1 : 5$	
$0 + 7 \rightarrow 7$	$1 + 7 \rightarrow 8$	$\dots$	$9 + 7 \rightarrow 1 : 6$	
$0 + 8 \rightarrow 8$	$1 + 8 \rightarrow 9$	$\dots$	$9 + 8 \rightarrow 1 : 7$	
$0 + 9 \rightarrow 9$	$1 + 9 \rightarrow 1 : 0$	$\dots$	$9 + 9 \rightarrow 1 : 8$	

rewriting  $(2 : 3) + (7 : 7) \rightarrow^* 7 : ((2 + 1) : 0)$

## Example (Addition on Natural Numbers in Decimal Notation)

signature       $0 \ 1 \ \dots \ 9$  (constants)     $+$      $:$  (binary, infix)

terms             $1 + 3 \quad 2 + (7 : 3) \quad (2 : (3 : x)) + ((1 + 7) : 2)$

rewrite rules

$0 + 0 \rightarrow 0$	$1 + 0 \rightarrow 1$	$\dots$	$9 + 0 \rightarrow 9$	$0 : x \rightarrow x$
$0 + 1 \rightarrow 1$	$1 + 1 \rightarrow 2$	$\dots$	$9 + 1 \rightarrow 1 : 0$	$x + (y : z) \rightarrow y : (x + z)$
$0 + 2 \rightarrow 2$	$1 + 2 \rightarrow 3$	$\dots$	$9 + 2 \rightarrow 1 : 1$	$(x : y) + z \rightarrow x : (y + z)$
$0 + 3 \rightarrow 3$	$1 + 3 \rightarrow 4$	$\dots$	$9 + 3 \rightarrow 1 : 2$	$x : (y : z) \rightarrow (x + y) : z$
$0 + 4 \rightarrow 4$	$1 + 4 \rightarrow 5$	$\dots$	$9 + 4 \rightarrow 1 : 3$	
$0 + 5 \rightarrow 5$	$1 + 5 \rightarrow 6$	$\dots$	$9 + 5 \rightarrow 1 : 4$	
$0 + 6 \rightarrow 6$	$1 + 6 \rightarrow 7$	$\dots$	$9 + 6 \rightarrow 1 : 5$	
$0 + 7 \rightarrow 7$	$1 + 7 \rightarrow 8$	$\dots$	$9 + 7 \rightarrow 1 : 6$	
$0 + 8 \rightarrow 8$	$1 + 8 \rightarrow 9$	$\dots$	$9 + 8 \rightarrow 1 : 7$	
$0 + 9 \rightarrow 9$	$1 + 9 \rightarrow 1 : 0$	$\dots$	$9 + 9 \rightarrow 1 : 8$	

rewriting       $(2 : 3) + (7 : 7) \rightarrow^* 7 : (3 : 0)$

## Example (Addition on Natural Numbers in Decimal Notation)

signature  $0\ 1\ \dots\ 9$  (constants)  $+ :$  (binary, infix)

terms  $1 + 3\ 2 + (7 : 3)\ (2 : (3 : x)) + ((1 + 7) : 2)$

rewrite rules

$0 + 0 \rightarrow 0$	$1 + 0 \rightarrow 1$	$\dots$	$9 + 0 \rightarrow 9$	$0 : x \rightarrow x$
$0 + 1 \rightarrow 1$	$1 + 1 \rightarrow 2$	$\dots$	$9 + 1 \rightarrow 1 : 0$	$x + (y : z) \rightarrow y : (x + z)$
$0 + 2 \rightarrow 2$	$1 + 2 \rightarrow 3$	$\dots$	$9 + 2 \rightarrow 1 : 1$	$(x : y) + z \rightarrow x : (y + z)$
$0 + 3 \rightarrow 3$	$1 + 3 \rightarrow 4$	$\dots$	$9 + 3 \rightarrow 1 : 2$	$x : (y : z) \rightarrow (x + y) : z$
$0 + 4 \rightarrow 4$	$1 + 4 \rightarrow 5$	$\dots$	$9 + 4 \rightarrow 1 : 3$	
$0 + 5 \rightarrow 5$	$1 + 5 \rightarrow 6$	$\dots$	$9 + 5 \rightarrow 1 : 4$	
$0 + 6 \rightarrow 6$	$1 + 6 \rightarrow 7$	$\dots$	$9 + 6 \rightarrow 1 : 5$	
$0 + 7 \rightarrow 7$	$1 + 7 \rightarrow 8$	$\dots$	$9 + 7 \rightarrow 1 : 6$	
$0 + 8 \rightarrow 8$	$1 + 8 \rightarrow 9$	$\dots$	$9 + 8 \rightarrow 1 : 7$	
$0 + 9 \rightarrow 9$	$1 + 9 \rightarrow 1 : 0$	$\dots$	$9 + 9 \rightarrow 1 : 8$	

rewriting  $(2 : 3) + (7 : 7) \rightarrow^* 7 : (3 : 0)$

## Example (Addition on Natural Numbers in Decimal Notation)

signature       $0 \ 1 \ \dots \ 9$  (constants)     $+$  : (binary, infix)

terms             $1 + 3 \quad 2 + (7 : 3) \quad (2 : (3 : x)) + ((1 + 7) : 2)$

rewrite rules     $0 + 0 \rightarrow 0 \quad 1 + 0 \rightarrow 1 \quad \dots \quad 9 + 0 \rightarrow 9 \quad 0 : x \rightarrow x$   
 $0 + 1 \rightarrow 1 \quad 1 + 1 \rightarrow 2 \quad \dots \quad 9 + 1 \rightarrow 1 : 0 \quad x + (y : z) \rightarrow y : (x + z)$   
 $0 + 2 \rightarrow 2 \quad 1 + 2 \rightarrow 3 \quad \dots \quad 9 + 2 \rightarrow 1 : 1 \quad (x : y) + z \rightarrow x : (y + z)$   
 $0 + 3 \rightarrow 3 \quad 1 + 3 \rightarrow 4 \quad \dots \quad 9 + 3 \rightarrow 1 : 2 \quad x : (y : z) \rightarrow (x + y) : z$   
 $0 + 4 \rightarrow 4 \quad 1 + 4 \rightarrow 5 \quad \dots \quad 9 + 4 \rightarrow 1 : 3$   
 $0 + 5 \rightarrow 5 \quad 1 + 5 \rightarrow 6 \quad \dots \quad 9 + 5 \rightarrow 1 : 4$   
 $0 + 6 \rightarrow 6 \quad 1 + 6 \rightarrow 7 \quad \dots \quad 9 + 6 \rightarrow 1 : 5$   
 $0 + 7 \rightarrow 7 \quad 1 + 7 \rightarrow 8 \quad \dots \quad 9 + 7 \rightarrow 1 : 6$   
 $0 + 8 \rightarrow 8 \quad 1 + 8 \rightarrow 9 \quad \dots \quad 9 + 8 \rightarrow 1 : 7$   
 $0 + 9 \rightarrow 9 \quad 1 + 9 \rightarrow 1 : 0 \quad \dots \quad 9 + 9 \rightarrow 1 : 8$

rewriting         $(2 : 3) + (7 : 7) \rightarrow^* (7 + 3) : 0$

## Example (Addition on Natural Numbers in Decimal Notation)

signature  $0\ 1\ \dots\ 9$  (constants)  $+ :$  (binary, infix)

terms  $1 + 3\ 2 + (7 : 3)\ (2 : (3 : x)) + ((1 + 7) : 2)$

rewrite rules

$0 + 0 \rightarrow 0$	$1 + 0 \rightarrow 1$	$\dots$	$9 + 0 \rightarrow 9$	$0 : x \rightarrow x$
$0 + 1 \rightarrow 1$	$1 + 1 \rightarrow 2$	$\dots$	$9 + 1 \rightarrow 1 : 0$	$x + (y : z) \rightarrow y : (x + z)$
$0 + 2 \rightarrow 2$	$1 + 2 \rightarrow 3$	$\dots$	$9 + 2 \rightarrow 1 : 1$	$(x : y) + z \rightarrow x : (y + z)$
$0 + 3 \rightarrow 3$	$1 + 3 \rightarrow 4$	$\dots$	$9 + 3 \rightarrow 1 : 2$	$x : (y : z) \rightarrow (x + y) : z$
$0 + 4 \rightarrow 4$	$1 + 4 \rightarrow 5$	$\dots$	$9 + 4 \rightarrow 1 : 3$	
$0 + 5 \rightarrow 5$	$1 + 5 \rightarrow 6$	$\dots$	$9 + 5 \rightarrow 1 : 4$	
$0 + 6 \rightarrow 6$	$1 + 6 \rightarrow 7$	$\dots$	$9 + 6 \rightarrow 1 : 5$	
$0 + 7 \rightarrow 7$	$1 + 7 \rightarrow 8$	$\dots$	$9 + 7 \rightarrow 1 : 6$	
$0 + 8 \rightarrow 8$	$1 + 8 \rightarrow 9$	$\dots$	$9 + 8 \rightarrow 1 : 7$	
$0 + 9 \rightarrow 9$	$1 + 9 \rightarrow 1 : 0$	$\dots$	$9 + 9 \rightarrow 1 : 8$	

rewriting  $(2 : 3) + (7 : 7) \rightarrow^* (7 + 3) : 0$

## Example (Addition on Natural Numbers in Decimal Notation)

signature       $0 \ 1 \ \dots \ 9$  (constants)     $+$      $:$  (binary, infix)

terms             $1 + 3$      $2 + (7 : 3)$      $(2 : (3 : x)) + ((1 + 7) : 2)$

rewrite rules     $0 + 0 \rightarrow 0$      $1 + 0 \rightarrow 1$      $\dots$      $9 + 0 \rightarrow 9$                      $0 : x \rightarrow x$   
 $0 + 1 \rightarrow 1$      $1 + 1 \rightarrow 2$      $\dots$      $9 + 1 \rightarrow 1 : 0$      $x + (y : z) \rightarrow y : (x + z)$   
 $0 + 2 \rightarrow 2$      $1 + 2 \rightarrow 3$      $\dots$      $9 + 2 \rightarrow 1 : 1$      $(x : y) + z \rightarrow x : (y + z)$   
 $0 + 3 \rightarrow 3$      $1 + 3 \rightarrow 4$      $\dots$      $9 + 3 \rightarrow 1 : 2$      $x : (y : z) \rightarrow (x + y) : z$   
 $0 + 4 \rightarrow 4$      $1 + 4 \rightarrow 5$      $\dots$      $9 + 4 \rightarrow 1 : 3$   
 $0 + 5 \rightarrow 5$      $1 + 5 \rightarrow 6$      $\dots$      $9 + 5 \rightarrow 1 : 4$   
 $0 + 6 \rightarrow 6$      $1 + 6 \rightarrow 7$      $\dots$      $9 + 6 \rightarrow 1 : 5$   
 $0 + 7 \rightarrow 7$      $1 + 7 \rightarrow 8$      $\dots$      $9 + 7 \rightarrow 1 : 6$   
 $0 + 8 \rightarrow 8$      $1 + 8 \rightarrow 9$      $\dots$      $9 + 8 \rightarrow 1 : 7$   
 $0 + 9 \rightarrow 9$      $1 + 9 \rightarrow 1 : 0$      $\dots$      $9 + 9 \rightarrow 1 : 8$

rewriting         $(2 : 3) + (7 : 7) \rightarrow^* (1 : 0) : 0$

## Example (Addition on Natural Numbers in Decimal Notation)

signature       $0 \ 1 \ \dots \ 9$  (constants)     $+$  : (binary, infix)

terms             $1 + 3 \quad 2 + (7 : 3) \quad (2 : (3 : x)) + ((1 + 7) : 2)$

rewrite rules     $0 + 0 \rightarrow 0 \quad 1 + 0 \rightarrow 1 \quad \dots \quad 9 + 0 \rightarrow 9 \quad 0 : x \rightarrow x$   
 $0 + 1 \rightarrow 1 \quad 1 + 1 \rightarrow 2 \quad \dots \quad 9 + 1 \rightarrow 1 : 0 \quad x + (y : z) \rightarrow y : (x + z)$   
 $0 + 2 \rightarrow 2 \quad 1 + 2 \rightarrow 3 \quad \dots \quad 9 + 2 \rightarrow 1 : 1 \quad (x : y) + z \rightarrow x : (y + z)$   
 $0 + 3 \rightarrow 3 \quad 1 + 3 \rightarrow 4 \quad \dots \quad 9 + 3 \rightarrow 1 : 2 \quad x : (y : z) \rightarrow (x + y) : z$   
 $0 + 4 \rightarrow 4 \quad 1 + 4 \rightarrow 5 \quad \dots \quad 9 + 4 \rightarrow 1 : 3$   
 $0 + 5 \rightarrow 5 \quad 1 + 5 \rightarrow 6 \quad \dots \quad 9 + 5 \rightarrow 1 : 4$   
 $0 + 6 \rightarrow 6 \quad 1 + 6 \rightarrow 7 \quad \dots \quad 9 + 6 \rightarrow 1 : 5$   
 $0 + 7 \rightarrow 7 \quad 1 + 7 \rightarrow 8 \quad \dots \quad 9 + 7 \rightarrow 1 : 6$   
 $0 + 8 \rightarrow 8 \quad 1 + 8 \rightarrow 9 \quad \dots \quad 9 + 8 \rightarrow 1 : 7$   
 $0 + 9 \rightarrow 9 \quad 1 + 9 \rightarrow 1 : 0 \quad \dots \quad 9 + 9 \rightarrow 1 : 8$

rewriting         $(2 : 3) + (7 : 7) \rightarrow^* (1 : 0) : 0$                       normal form

## Example (Group Theory)

signature       $e$  (constant)     $-$  (unary, postfix)     $\cdot$  (binary, infix)

## Example (Group Theory)

signature       $e$  (constant)    $^-$  (unary, postfix)    $\cdot$  (binary, infix)

axioms       $e \cdot x \approx x$     $x^- \cdot x \approx e$     $(x \cdot y) \cdot z \approx x \cdot (y \cdot z)$     $\mathcal{E}$

## Example (Group Theory)

signature       $e$  (constant)    $^-$  (unary, postfix)    $\cdot$  (binary, infix)

axioms             $e \cdot x \approx x$      $x^- \cdot x \approx e$      $(x \cdot y) \cdot z \approx x \cdot (y \cdot z)$      $\mathcal{E}$

theorems         $e^- \approx_{\mathcal{E}} e$      $(x \cdot y)^- \approx_{\mathcal{E}} y^- \cdot x^-$



## Example (Group Theory)

signature       $e$  (constant)     $^-$  (unary, postfix)     $\cdot$  (binary, infix)

axioms             $e \cdot x \approx x$      $x^- \cdot x \approx e$      $(x \cdot y) \cdot z \approx x \cdot (y \cdot z)$      $\mathcal{E}$

theorems         $e^- \approx_{\mathcal{E}} e$      $(x \cdot y)^- \approx_{\mathcal{E}} y^- \cdot x^-$

rewrite rules

$$e \cdot x \rightarrow x \qquad x \cdot e \rightarrow x \qquad \mathcal{R}$$

$$x^- \cdot x \rightarrow e \qquad x \cdot x^- \rightarrow e$$

$$(x \cdot y) \cdot z \rightarrow x \cdot (y \cdot z) \qquad x^{--} \rightarrow x$$

$$e^- \rightarrow e \qquad (x \cdot y)^- \rightarrow y^- \cdot x^-$$

$$x^- \cdot (x \cdot y) \rightarrow y \qquad x \cdot (x^- \cdot y) \rightarrow y$$

## Example (Group Theory)

signature      $e$  (constant)      $-$  (unary, postfix)      $\cdot$  (binary, infix)

axioms      $e \cdot x \approx x$       $x^- \cdot x \approx e$       $(x \cdot y) \cdot z \approx x \cdot (y \cdot z)$       $\mathcal{E}$

theorems      $e^- \approx_{\mathcal{E}} e$       $(x \cdot y)^- \approx_{\mathcal{E}} y^- \cdot x^-$

rewrite rules      $e \cdot x \rightarrow x$       $x \cdot e \rightarrow x$       $\mathcal{R}$

$x^- \cdot x \rightarrow e$       $x \cdot x^- \rightarrow e$

$(x \cdot y) \cdot z \rightarrow x \cdot (y \cdot z)$       $x^{--} \rightarrow x$

$e^- \rightarrow e$       $(x \cdot y)^- \rightarrow y^- \cdot x^-$

$x^- \cdot (x \cdot y) \rightarrow y$       $x \cdot (x^- \cdot y) \rightarrow y$

①  $s \approx t$  is valid in  $\mathcal{E}$  ( $s \approx_{\mathcal{E}} t$ )      $\iff$       $s$  and  $t$  have same normal form with respect to  $\mathcal{R}$



## Example (Group Theory)

signature      $e$  (constant)      $^-$  (unary, postfix)      $\cdot$  (binary, infix)

axioms      $e \cdot x \approx x$       $x^- \cdot x \approx e$       $(x \cdot y) \cdot z \approx x \cdot (y \cdot z)$       $\mathcal{E}$

theorems      $e^- \approx_{\mathcal{E}} e$       $(x \cdot y)^- \approx_{\mathcal{E}} y^- \cdot x^-$

rewrite rules      $e \cdot x \rightarrow x$       $x \cdot e \rightarrow x$       $\mathcal{R}$

$x^- \cdot x \rightarrow e$       $x \cdot x^- \rightarrow e$

$(x \cdot y) \cdot z \rightarrow x \cdot (y \cdot z)$       $x^{--} \rightarrow x$

$e^- \rightarrow e$       $(x \cdot y)^- \rightarrow y^- \cdot x^-$

$x^- \cdot (x \cdot y) \rightarrow y$       $x \cdot (x^- \cdot y) \rightarrow y$

①  $s \approx t$  is valid in  $\mathcal{E}$  ( $s \approx_{\mathcal{E}} t$ )      $\iff$       $s$  and  $t$  have same normal form with respect to  $\mathcal{R}$

②  $\mathcal{R}$  admits no infinite computations

① & ②      $\implies$       $\mathcal{E}$  has decidable validity problem

## Example (Combinatory Logic)

signature      S K I (constants)

## Example (Combinatory Logic)

signature      S K I (constants)   · (application, binary, infix)

## Example (Combinatory Logic)

signature       $S \ K \ I$  (constants)    $\cdot$  (application, binary, infix)

terms           $S \ ((K \cdot I) \cdot I) \cdot S \ (x \cdot z) \cdot (y \cdot z)$

## Example (Combinatory Logic)

signature       $S \ K \ I$  (constants)     $\cdot$  (application, binary, infix)

terms             $S \ ((K \cdot I) \cdot I) \cdot S \ (x \cdot z) \cdot (y \cdot z)$

rewrite rules

$$I \cdot x \rightarrow x$$

$$(K \cdot x) \cdot y \rightarrow x$$

$$((S \cdot x) \cdot y) \cdot z \rightarrow (x \cdot z) \cdot (y \cdot z)$$

## Example (Combinatory Logic)

signature       $S \ K \ I$  (constants)     $\cdot$  (application, binary, infix)

terms             $S \ ((K \cdot I) \cdot I) \cdot S \ (x \cdot z) \cdot (y \cdot z)$

rewrite rules

$$I \cdot x \rightarrow x$$

$$(K \cdot x) \cdot y \rightarrow x$$

$$((S \cdot x) \cdot y) \cdot z \rightarrow (x \cdot z) \cdot (y \cdot z)$$

rewriting

$$((S \cdot K) \cdot K) \cdot x$$

## Example (Combinatory Logic)

signature       $S \ K \ I$  (constants)     $\cdot$  (application, binary, infix)

terms             $S \ ((K \cdot I) \cdot I) \cdot S \ (x \cdot z) \cdot (y \cdot z)$

rewrite rules

$$I \cdot x \rightarrow x$$

$$(K \cdot x) \cdot y \rightarrow x$$

$$((S \cdot x) \cdot y) \cdot z \rightarrow (x \cdot z) \cdot (y \cdot z)$$

rewriting

$$((S \cdot K) \cdot K) \cdot x \rightarrow (K \cdot x) \cdot (K \cdot x)$$

## Example (Combinatory Logic)

signature       $S \ K \ I$  (constants)     $\cdot$  (application, binary, infix)

terms           $S \ ((K \cdot I) \cdot I) \cdot S \ (x \cdot z) \cdot (y \cdot z)$

rewrite rules

$$I \cdot x \rightarrow x$$

$$(K \cdot x) \cdot y \rightarrow x$$

$$((S \cdot x) \cdot y) \cdot z \rightarrow (x \cdot z) \cdot (y \cdot z)$$

rewriting

$$((S \cdot K) \cdot K) \cdot x \rightarrow (K \cdot x) \cdot (K \cdot x) \rightarrow x$$

## Example (Combinatory Logic)

signature       $S \ K \ I$  (constants)     $\cdot$  (application, binary, infix)

terms           $S \ ((K \cdot I) \cdot I) \cdot S \ (x \cdot z) \cdot (y \cdot z)$

rewrite rules

$$I \cdot x \rightarrow x$$

$$(K \cdot x) \cdot y \rightarrow x$$

$$((S \cdot x) \cdot y) \cdot z \rightarrow (x \cdot z) \cdot (y \cdot z)$$

rewriting

$$((S \cdot K) \cdot K) \cdot x \rightarrow (K \cdot x) \cdot (K \cdot x) \rightarrow x$$

inventor

**Moses Schönfinkel** (1924)



## Example (Combinatory Logic)

signature       $S \ K \ I$  (constants)     $\cdot$  (application, binary, infix)

terms             $S \ ((K \cdot I) \cdot I) \cdot S \ (x \cdot z) \cdot (y \cdot z)$

rewrite rules

$$I \cdot x \rightarrow x$$

$$(K \cdot x) \cdot y \rightarrow x$$

$$((S \cdot x) \cdot y) \cdot z \rightarrow (x \cdot z) \cdot (y \cdot z)$$

rewriting

$$((S \cdot K) \cdot K) \cdot x \rightarrow (K \cdot x) \cdot (K \cdot x) \rightarrow x$$

inventor

**Moses Schönfinkel** (1924)



Combinatory Logic is Turing-complete

# Outline

## 1. Organisation

## 2. Examples

## 3. Terms

Contexts

Substitutions

## 4. Exercises

## 5. Further Reading

## Definitions (Terms)

- ▶ **signature**       $\mathcal{F}$       function symbols with arities

## Definitions (Terms)

- ▶ signature  $\mathcal{F}$  function symbols with arities
- ▶ **variables**  $\mathcal{V}$   $\mathcal{F} \cap \mathcal{V} = \emptyset$  infinitely many

## Definitions (Terms)

- ▶ signature  $\mathcal{F}$  function symbols with arities
- ▶ variables  $\mathcal{V}$   $\mathcal{F} \cap \mathcal{V} = \emptyset$  infinitely many
- ▶ **terms**  $\mathcal{T}(\mathcal{F}, \mathcal{V})$  smallest set such that
  - ▶  $\mathcal{V} \subseteq \mathcal{T}(\mathcal{F}, \mathcal{V})$

## Definitions (Terms)

- ▶ signature  $\mathcal{F}$  function symbols with arities
- ▶ variables  $\mathcal{V}$   $\mathcal{F} \cap \mathcal{V} = \emptyset$  infinitely many
- ▶ **terms**  $\mathcal{T}(\mathcal{F}, \mathcal{V})$  smallest set such that
  - ▶  $\mathcal{V} \subseteq \mathcal{T}(\mathcal{F}, \mathcal{V})$
  - ▶ if  $f \in \mathcal{F}$  has arity 0 then  $f \in \mathcal{T}(\mathcal{F}, \mathcal{V})$

## Definitions (Terms)

- ▶ signature  $\mathcal{F}$  function symbols with arities
- ▶ variables  $\mathcal{V}$   $\mathcal{F} \cap \mathcal{V} = \emptyset$  infinitely many
- ▶ **terms**  $\mathcal{T}(\mathcal{F}, \mathcal{V})$  smallest set such that
  - ▶  $\mathcal{V} \subseteq \mathcal{T}(\mathcal{F}, \mathcal{V})$
  - ▶ if  $f \in \mathcal{F}$  has arity 0 then  $f \in \mathcal{T}(\mathcal{F}, \mathcal{V})$
  - ▶ if  $f \in \mathcal{F}$  has arity  $n \geq 1$  and  $t_1, \dots, t_n \in \mathcal{T}(\mathcal{F}, \mathcal{V})$  then  $f(t_1, \dots, t_n) \in \mathcal{T}(\mathcal{F}, \mathcal{V})$

## Definitions (Terms)

- ▶ signature  $\mathcal{F}$  function symbols with arities
- ▶ variables  $\mathcal{V}$   $\mathcal{F} \cap \mathcal{V} = \emptyset$  infinitely many
- ▶ **terms**  $\mathcal{T}(\mathcal{F}, \mathcal{V})$  smallest set such that
  - ▶  $\mathcal{V} \subseteq \mathcal{T}(\mathcal{F}, \mathcal{V})$
  - ▶ if  $f \in \mathcal{F}$  has arity  $n \geq 0$  and  $t_1, \dots, t_n \in \mathcal{T}(\mathcal{F}, \mathcal{V})$  then  $f(t_1, \dots, t_n) \in \mathcal{T}(\mathcal{F}, \mathcal{V})$  assuming  $f() = f$

## Definitions (Terms)

- ▶ signature  $\mathcal{F}$  function symbols with arities
- ▶ variables  $\mathcal{V}$   $\mathcal{F} \cap \mathcal{V} = \emptyset$  infinitely many
- ▶ terms  $\mathcal{T}(\mathcal{F}, \mathcal{V})$  smallest set such that
  - ▶  $\mathcal{V} \subseteq \mathcal{T}(\mathcal{F}, \mathcal{V})$
  - ▶ if  $f \in \mathcal{F}$  has arity  $n \geq 0$  and  $t_1, \dots, t_n \in \mathcal{T}(\mathcal{F}, \mathcal{V})$  then  $f(t_1, \dots, t_n) \in \mathcal{T}(\mathcal{F}, \mathcal{V})$  assuming  $f() = f$
- ▶ **ground terms**  $\mathcal{T}(\mathcal{F})$  smallest set such that
  - ▶ if  $f \in \mathcal{F}$  has arity  $n \geq 0$  and  $t_1, \dots, t_n \in \mathcal{T}(\mathcal{F})$  then  $f(t_1, \dots, t_n) \in \mathcal{T}(\mathcal{F})$

►  $\mathcal{V}\text{ar}(\cdot)$

$$\mathcal{V}\text{ar}(t) = \begin{cases} \{t\} & \text{if } t \in \mathcal{V} \\ \bigcup_{i=1}^n \mathcal{V}\text{ar}(t_i) & \text{if } t = f(t_1, \dots, t_n) \end{cases}$$

►  $\mathcal{V}\text{ar}(\cdot)$

$$\mathcal{V}\text{ar}((2 : x) + ((1 : x) : y)) = \{x, y\}$$

$$\mathcal{V}\text{ar}(t) = \begin{cases} \{t\} & \text{if } t \in \mathcal{V} \\ \bigcup_{i=1}^n \mathcal{V}\text{ar}(t_i) & \text{if } t = f(t_1, \dots, t_n) \end{cases}$$

►  $\mathcal{V}\text{ar}(\cdot)$

$$\mathcal{V}\text{ar}((2 : x) + ((1 : x) : y)) = \{x, y\}$$

$$\mathcal{V}\text{ar}(t) = \begin{cases} \{t\} & \text{if } t \in \mathcal{V} \\ \bigcup_{i=1}^n \mathcal{V}\text{ar}(t_i) & \text{if } t = f(t_1, \dots, t_n) \end{cases}$$

►  $\mathcal{F}\text{un}(\cdot)$

$$\mathcal{F}\text{un}(t) = \begin{cases} \emptyset & \text{if } t \in \mathcal{V} \\ \{f\} \cup \bigcup_{i=1}^n \mathcal{F}\text{un}(t_i) & \text{if } t = f(t_1, \dots, t_n) \end{cases}$$

## Definitions (Operations on Terms)

►  $\mathcal{V}\text{ar}(\cdot)$

$$\mathcal{V}\text{ar}((2 : x) + ((1 : x) : y)) = \{x, y\}$$

$$\mathcal{V}\text{ar}(t) = \begin{cases} \{t\} & \text{if } t \in \mathcal{V} \\ \bigcup_{i=1}^n \mathcal{V}\text{ar}(t_i) & \text{if } t = f(t_1, \dots, t_n) \end{cases}$$

►  $\mathcal{F}\text{un}(\cdot)$

$$\mathcal{F}\text{un}((2 : x) + ((1 : x) : y)) = \{2, :, +, 1\}$$

$$\mathcal{F}\text{un}(t) = \begin{cases} \emptyset & \text{if } t \in \mathcal{V} \\ \{f\} \cup \bigcup_{i=1}^n \mathcal{F}\text{un}(t_i) & \text{if } t = f(t_1, \dots, t_n) \end{cases}$$

## Definitions (Operations on Terms)

►  $\mathcal{V}\text{ar}(\cdot)$

$$\mathcal{V}\text{ar}((2 : x) + ((1 : x) : y)) = \{x, y\}$$

$$\mathcal{V}\text{ar}(t) = \begin{cases} \{t\} & \text{if } t \in \mathcal{V} \\ \bigcup_{i=1}^n \mathcal{V}\text{ar}(t_i) & \text{if } t = f(t_1, \dots, t_n) \end{cases}$$

►  $\mathcal{F}\text{un}(\cdot)$

$$\mathcal{F}\text{un}((2 : x) + ((1 : x) : y)) = \{2, :, +, 1\}$$

$$\mathcal{F}\text{un}(t) = \begin{cases} \emptyset & \text{if } t \in \mathcal{V} \\ \{f\} \cup \bigcup_{i=1}^n \mathcal{F}\text{un}(t_i) & \text{if } t = f(t_1, \dots, t_n) \end{cases}$$

► **size**  $|\cdot|$

$$|t| = \begin{cases} 1 & \text{if } t \in \mathcal{V} \\ 1 + \sum_{i=1}^n |t_i| & \text{if } t = f(t_1, \dots, t_n) \end{cases}$$

## Definitions (Operations on Terms)

►  $\mathcal{V}\text{ar}(\cdot)$

$$\mathcal{V}\text{ar}((2 : x) + ((1 : x) : y)) = \{x, y\}$$

$$\mathcal{V}\text{ar}(t) = \begin{cases} \{t\} & \text{if } t \in \mathcal{V} \\ \bigcup_{i=1}^n \mathcal{V}\text{ar}(t_i) & \text{if } t = f(t_1, \dots, t_n) \end{cases}$$

►  $\mathcal{F}\text{un}(\cdot)$

$$\mathcal{F}\text{un}((2 : x) + ((1 : x) : y)) = \{2, :, +, 1\}$$

$$\mathcal{F}\text{un}(t) = \begin{cases} \emptyset & \text{if } t \in \mathcal{V} \\ \{f\} \cup \bigcup_{i=1}^n \mathcal{F}\text{un}(t_i) & \text{if } t = f(t_1, \dots, t_n) \end{cases}$$

► **size**  $|\cdot|$

$$|(2 : x) + ((1 : x) : y)| = ?$$

$$|t| = \begin{cases} 1 & \text{if } t \in \mathcal{V} \\ 1 + \sum_{i=1}^n |t_i| & \text{if } t = f(t_1, \dots, t_n) \end{cases}$$

## Definitions (Operations on Terms)

►  $\mathcal{V}\text{ar}(\cdot)$

$$\mathcal{V}\text{ar}((2 : x) + ((1 : x) : y)) = \{x, y\}$$

$$\mathcal{V}\text{ar}(t) = \begin{cases} \{t\} & \text{if } t \in \mathcal{V} \\ \bigcup_{i=1}^n \mathcal{V}\text{ar}(t_i) & \text{if } t = f(t_1, \dots, t_n) \end{cases}$$

►  $\mathcal{F}\text{un}(\cdot)$

$$\mathcal{F}\text{un}((2 : x) + ((1 : x) : y)) = \{2, :, +, 1\}$$

$$\mathcal{F}\text{un}(t) = \begin{cases} \emptyset & \text{if } t \in \mathcal{V} \\ \{f\} \cup \bigcup_{i=1}^n \mathcal{F}\text{un}(t_i) & \text{if } t = f(t_1, \dots, t_n) \end{cases}$$

► **size**  $|\cdot|$

$$|(2 : x) + ((1 : x) : y)| = 9$$

$$|t| = \begin{cases} 1 & \text{if } t \in \mathcal{V} \\ 1 + \sum_{i=1}^n |t_i| & \text{if } t = f(t_1, \dots, t_n) \end{cases}$$

► **root( $\cdot$ )**       $\text{root}(t) = \begin{cases} t & \text{if } t \in \mathcal{V} \\ f & \text{if } t = f(t_1, \dots, t_n) \end{cases}$


## Definitions (Operations on Terms)

►  $\text{root}(\cdot)$

$$\text{root}(t) = \begin{cases} t & \text{if } t \in \mathcal{V} \\ f & \text{if } t = f(t_1, \dots, t_n) \end{cases}$$

## Example

$(2 : x) + ((1 : x) : y)$




## Definitions (Operations on Terms)

►  $\text{root}(\cdot)$

$$\text{root}(t) = \begin{cases} t & \text{if } t \in \mathcal{V} \\ f & \text{if } t = f(t_1, \dots, t_n) \end{cases}$$

## Example

$(2 : x) + ((1 : x) : y)$




$\text{root}((2 : x) + ((1 : x) : y)) = +$

## Definitions (Operations on Terms)

- ▶  $\text{root}(\cdot)$        $\text{root}(t) = \begin{cases} t & \text{if } t \in \mathcal{V} \\ f & \text{if } t = f(t_1, \dots, t_n) \end{cases}$
- ▶  $\text{height}(\cdot)$        $\text{height}(t) = \begin{cases} 0 & \text{if } t \in \mathcal{V} \text{ or } t \text{ is constant} \\ 1 + \max_{1 \leq i \leq n} \text{height}(t_i) & \text{if } t = f(t_1, \dots, t_n) \text{ with } n \geq 1 \end{cases}$

## Example

$(2 : x) + ((1 : x) : y)$




$\text{root}((2 : x) + ((1 : x) : y)) = +$

## Definitions (Operations on Terms)

- ▶  $\text{root}(\cdot)$        $\text{root}(t) = \begin{cases} t & \text{if } t \in \mathcal{V} \\ f & \text{if } t = f(t_1, \dots, t_n) \end{cases}$
- ▶  $\text{height}(\cdot)$        $\text{height}(t) = \begin{cases} 0 & \text{if } t \in \mathcal{V} \text{ or } t \text{ is constant} \\ 1 + \max_{1 \leq i \leq n} \text{height}(t_i) & \text{if } t = f(t_1, \dots, t_n) \text{ with } n \geq 1 \end{cases}$

## Example

$(2 : x) + ((1 : x) : y)$



$\text{root}((2 : x) + ((1 : x) : y)) = +$


$\text{height}((2 : x) + ((1 : x) : y)) = ?$

## Definitions (Operations on Terms)

- ▶  $\text{root}(\cdot)$        $\text{root}(t) = \begin{cases} t & \text{if } t \in \mathcal{V} \\ f & \text{if } t = f(t_1, \dots, t_n) \end{cases}$
- ▶  $\text{height}(\cdot)$        $\text{height}(t) = \begin{cases} 0 & \text{if } t \in \mathcal{V} \text{ or } t \text{ is constant} \\ 1 + \max_{1 \leq i \leq n} \text{height}(t_i) & \text{if } t = f(t_1, \dots, t_n) \text{ with } n \geq 1 \end{cases}$

## Example

$(2 : x) + ((1 : x) : y)$



$\text{root}((2 : x) + ((1 : x) : y)) = +$

$\text{height}((2 : x) + ((1 : x) : y)) = 3$

## Definitions (Subterms)

▶  $s \trianglelefteq t$   $s$  is **subterm** of  $t$

▶  $s = t$

## Definitions (Subterms)

- ▶  $s \trianglelefteq t$   $s$  is **subterm** of  $t$ 
  - ▶  $s = t$  or
  - ▶  $t = f(t_1, \dots, t_n)$  and  $s \trianglelefteq t_i$  for some  $1 \leq i \leq n$

## Definitions (Subterms)

- ▶  $s \sqsubseteq t$   $s$  is subterm of  $t$ 
  - ▶  $s = t$  or
  - ▶  $t = f(t_1, \dots, t_n)$  and  $s \sqsubseteq t_i$  for some  $1 \leq i \leq n$
- ▶  $s \triangleleft t$   $s$  is **proper** subterm of  $t$ 
  - ▶  $t = f(t_1, \dots, t_n)$  and  $s \sqsubseteq t_i$  for some  $1 \leq i \leq n$

## Definitions (Subterms)

- ▶  $s \sqsubseteq t$   $s$  is subterm of  $t$ 
  - ▶  $s = t$  or
  - ▶  $t = f(t_1, \dots, t_n)$  and  $s \sqsubseteq t_i$  for some  $1 \leq i \leq n$
- ▶  $s \triangleleft t$   $s$  is proper subterm of  $t$ 
  - ▶  $t = f(t_1, \dots, t_n)$  and  $s \sqsubseteq t_i$  for some  $1 \leq i \leq n$

## Example

term  $(2 : x) + ((1 : x) : y)$  has subterms

2

## Definitions (Subterms)

- ▶  $s \sqsubseteq t$   $s$  is subterm of  $t$ 
  - ▶  $s = t$  or
  - ▶  $t = f(t_1, \dots, t_n)$  and  $s \sqsubseteq t_i$  for some  $1 \leq i \leq n$
- ▶  $s \triangleleft t$   $s$  is proper subterm of  $t$ 
  - ▶  $t = f(t_1, \dots, t_n)$  and  $s \sqsubseteq t_i$  for some  $1 \leq i \leq n$

## Example

term  $(2 : x) + ((1 : x) : y)$  has subterms

2     $x$

## Definitions (Subterms)

- ▶  $s \sqsubseteq t$   $s$  is subterm of  $t$ 
  - ▶  $s = t$  or
  - ▶  $t = f(t_1, \dots, t_n)$  and  $s \sqsubseteq t_i$  for some  $1 \leq i \leq n$
- ▶  $s \triangleleft t$   $s$  is proper subterm of  $t$ 
  - ▶  $t = f(t_1, \dots, t_n)$  and  $s \sqsubseteq t_i$  for some  $1 \leq i \leq n$

## Example

term  $(2 : x) + ((1 : x) : y)$  has subterms

2     $x$      $2 : x$

## Definitions (Subterms)

- ▶  $s \sqsubseteq t$   $s$  is subterm of  $t$ 
  - ▶  $s = t$  or
  - ▶  $t = f(t_1, \dots, t_n)$  and  $s \sqsubseteq t_i$  for some  $1 \leq i \leq n$
- ▶  $s \triangleleft t$   $s$  is proper subterm of  $t$ 
  - ▶  $t = f(t_1, \dots, t_n)$  and  $s \sqsubseteq t_i$  for some  $1 \leq i \leq n$

## Example

term  $(2 : x) + ((1 : x) : y)$  has subterms

2    x    2 : x    1

## Definitions (Subterms)

- ▶  $s \sqsubseteq t$   $s$  is subterm of  $t$ 
  - ▶  $s = t$  or
  - ▶  $t = f(t_1, \dots, t_n)$  and  $s \sqsubseteq t_i$  for some  $1 \leq i \leq n$
- ▶  $s \triangleleft t$   $s$  is proper subterm of  $t$ 
  - ▶  $t = f(t_1, \dots, t_n)$  and  $s \sqsubseteq t_i$  for some  $1 \leq i \leq n$

## Example

term  $(2 : x) + ((1 : x) : y)$  has subterms

2    x    2 : x    1    1 : x

## Definitions (Subterms)

- ▶  $s \sqsubseteq t$   $s$  is subterm of  $t$ 
  - ▶  $s = t$  or
  - ▶  $t = f(t_1, \dots, t_n)$  and  $s \sqsubseteq t_i$  for some  $1 \leq i \leq n$
- ▶  $s \triangleleft t$   $s$  is proper subterm of  $t$ 
  - ▶  $t = f(t_1, \dots, t_n)$  and  $s \sqsubseteq t_i$  for some  $1 \leq i \leq n$

## Example

term  $(2 : x) + ((1 : x) : y)$  has subterms

2   x   2 : x   1   1 : x   y

## Definitions (Subterms)

- ▶  $s \sqsubseteq t$   $s$  is subterm of  $t$ 
  - ▶  $s = t$  or
  - ▶  $t = f(t_1, \dots, t_n)$  and  $s \sqsubseteq t_i$  for some  $1 \leq i \leq n$
- ▶  $s \triangleleft t$   $s$  is proper subterm of  $t$ 
  - ▶  $t = f(t_1, \dots, t_n)$  and  $s \sqsubseteq t_i$  for some  $1 \leq i \leq n$

## Example

term  $(2 : x) + ((1 : x) : y)$  has subterms

2    x    2 : x    1    1 : x    y    (1 : x) : y

## Definitions (Subterms)

- ▶  $s \sqsubseteq t$   $s$  is subterm of  $t$ 
  - ▶  $s = t$  or
  - ▶  $t = f(t_1, \dots, t_n)$  and  $s \sqsubseteq t_i$  for some  $1 \leq i \leq n$
- ▶  $s \triangleleft t$   $s$  is proper subterm of  $t$ 
  - ▶  $t = f(t_1, \dots, t_n)$  and  $s \sqsubseteq t_i$  for some  $1 \leq i \leq n$

## Example

term  $(2 : x) + ((1 : x) : y)$  has subterms

2    x    2 : x    1    1 : x    y    (1 : x) : y    (2 : x) + ((1 : x) : y)

# Outline

## 1. Organisation

## 2. Examples

## 3. Terms

Contexts

Substitutions

## 4. Exercises

## 5. Further Reading

## Definitions (Context)

- ▶ **context** is term with one **hole**

## Definitions (Context)

- ▶ **context** is term with one **hole**: element of  $\mathcal{T}(\mathcal{F} \cup \{\square\}, \mathcal{V})$  that contains exactly one occurrence of special constant  $\square$

## Definitions (Context)

- ▶ **context** is term with one hole: element of  $\mathcal{T}(\mathcal{F} \cup \{\square\}, \mathcal{V})$  that contains exactly one occurrence of special constant  $\square$

## Examples

- ▶  $\square$      $s(0) + s(s(\square))$      $\square + x$

## Definitions (Context)

- ▶ context is term with one hole: element of  $\mathcal{T}(\mathcal{F} \cup \{\square\}, \mathcal{V})$  that contains exactly one occurrence of special constant  $\square$
- ▶  $C[t]$  denotes result of replacing hole in context  $C$  by term  $t$

## Examples

- ▶  $\square$      $s(0) + s(s(\square))$      $\square + x$

## Definitions (Context)

- ▶ context is term with one hole: element of  $\mathcal{T}(\mathcal{F} \cup \{\square\}, \mathcal{V})$  that contains exactly one occurrence of special constant  $\square$
- ▶  $C[t]$  denotes result of replacing hole in context  $C$  by term  $t$

## Examples

- ▶  $\square \quad s(0) + s(s(\square)) \quad \square + x$
- ▶  $\square[s(0)] = s(0) \quad (\square + x)[0 + x] = (0 + x) + x$

## Definitions (Context)

- ▶ context is term with one hole: element of  $\mathcal{T}(\mathcal{F} \cup \{\square\}, \mathcal{V})$  that contains exactly one occurrence of special constant  $\square$
- ▶  $C[t]$  denotes result of replacing hole in context  $C$  by term  $t$
- ▶ binary relation  $R$  on terms is **closed under contexts** if for all terms  $s, t$

$$s R t \implies C[s] R C[t] \text{ for all contexts } C$$

## Examples

- ▶  $\square \quad s(0) + s(s(\square)) \quad \square + x$
- ▶  $\square[s(0)] = s(0) \quad (\square + x)[0 + x] = (0 + x) + x$

## Definitions (Context)

- ▶ context is term with one hole: element of  $\mathcal{T}(\mathcal{F} \cup \{\square\}, \mathcal{V})$  that contains exactly one occurrence of special constant  $\square$
- ▶  $C[t]$  denotes result of replacing hole in context  $C$  by term  $t$
- ▶ binary relation  $R$  on terms is **closed under contexts** if for all terms  $s, t$

$$s R t \implies C[s] R C[t] \text{ for all contexts } C$$

## Examples

- ▶  $\square \quad s(0) + s(s(\square)) \quad \square + x$
- ▶  $\square[s(0)] = s(0) \quad (\square + x)[0 + x] = (0 + x) + x$
- ▶ subterm relation is not closed under contexts:  $0 \sqsubseteq s(0)$  but  $0 + 0 \not\sqsubseteq s(0) + 0$

►  $s \trianglelefteq t \iff t = C[s]$  for some context  $C$

## Lemmata

- ▶  $s \trianglelefteq t \iff t = C[s]$  for some context  $C$
- ▶  $s \triangleleft t \iff t = C[s]$  for some context  $C \neq \square$

# Outline

## 1. Organisation

## 2. Examples

## 3. Terms

Contexts

Substitutions

## 4. Exercises

## 5. Further Reading

## Definitions (Substitution)

- ▶ **substitution** is mapping  $\sigma: \mathcal{V} \rightarrow \mathcal{T}(\mathcal{F}, \mathcal{V})$  such that its domain

$$\text{Dom}(\sigma) = \{x \in \mathcal{V} \mid \sigma(x) \neq x\}$$

is finite

## Definitions (Substitution)

- ▶ substitution is mapping  $\sigma: \mathcal{V} \rightarrow \mathcal{T}(\mathcal{F}, \mathcal{V})$  such that its domain

$$\text{Dom}(\sigma) = \{x \in \mathcal{V} \mid \sigma(x) \neq x\}$$

is finite

- ▶ (unique) substitution with empty domain is denoted by  $\varepsilon$

## Definitions (Substitution)

- ▶ substitution is mapping  $\sigma: \mathcal{V} \rightarrow \mathcal{T}(\mathcal{F}, \mathcal{V})$  such that its domain

$$\text{Dom}(\sigma) = \{x \in \mathcal{V} \mid \sigma(x) \neq x\}$$

is finite

- ▶ (unique) substitution with empty domain is denoted by  $\varepsilon$
- ▶ **application** of substitution  $\sigma$  to term  $t$

$$t\sigma = \begin{cases} \sigma(t) & \text{if } t \in \mathcal{V} \\ f(t_1\sigma, \dots, t_n\sigma) & \text{if } t = f(t_1, \dots, t_n) \end{cases}$$

## Definitions (Substitution)

- ▶ substitution is mapping  $\sigma: \mathcal{V} \rightarrow \mathcal{T}(\mathcal{F}, \mathcal{V})$  such that its domain

$$\text{Dom}(\sigma) = \{x \in \mathcal{V} \mid \sigma(x) \neq x\}$$

is finite

- ▶ (unique) substitution with empty domain is denoted by  $\varepsilon$
- ▶ application of substitution  $\sigma$  to term  $t$

$$t\sigma = \begin{cases} \sigma(t) & \text{if } t \in \mathcal{V} \\ f(t_1\sigma, \dots, t_n\sigma) & \text{if } t = f(t_1, \dots, t_n) \end{cases}$$

## Example

$$t = x + s(y + z)$$

$$\sigma = \{x \mapsto s(y), y \mapsto x + s(0)\}$$

## Definitions (Substitution)

- ▶ substitution is mapping  $\sigma: \mathcal{V} \rightarrow \mathcal{T}(\mathcal{F}, \mathcal{V})$  such that its domain

$$\text{Dom}(\sigma) = \{x \in \mathcal{V} \mid \sigma(x) \neq x\}$$

is finite

- ▶ (unique) substitution with empty domain is denoted by  $\varepsilon$
- ▶ application of substitution  $\sigma$  to term  $t$

$$t\sigma = \begin{cases} \sigma(t) & \text{if } t \in \mathcal{V} \\ f(t_1\sigma, \dots, t_n\sigma) & \text{if } t = f(t_1, \dots, t_n) \end{cases}$$

## Example

$$t = x + s(y + z)$$

$$\sigma = \{x \mapsto s(y), y \mapsto x + s(0)\}$$

$$t\sigma = s(y) + s((x + s(0)) + z)$$

## Definition (Closure under Substitutions)

relation  $R$  on terms is **closed under substitutions** if for all terms  $s, t$

$$s R t \implies s\sigma R t\sigma \text{ for all substitutions } \sigma$$

## Definition (Closure under Substitutions)

relation  $R$  on terms is closed under substitutions if for all terms  $s, t$

$$s R t \implies s \sigma R t \sigma \text{ for all substitutions } \sigma$$

## Lemma

subterm relation is closed under substitutions

## Definition (Closure under Substitutions)

relation  $R$  on terms is closed under substitutions if for all terms  $s, t$

$$s R t \implies s \sigma R t \sigma \text{ for all substitutions } \sigma$$

## Lemma

subterm relation is closed under substitutions

## Definition (Matching Problem)

instance: terms  $s, t$

question:  $\exists$  substitution  $\sigma: s \sigma = t$ ?

## Definition (Closure under Substitutions)

relation  $R$  on terms is closed under substitutions if for all terms  $s, t$

$$s R t \implies s \sigma R t \sigma \text{ for all substitutions } \sigma$$

## Lemma

subterm relation is closed under substitutions

## Definition (Matching Problem)

instance: terms  $s, t$

question:  $\exists$  substitution  $\sigma: s \sigma = t?$  ( $t$  is **instance** of  $s$ ,  $t \succeq s$ )

## Examples

①  $e^- \cdot e^{-}$  is instance of  $x \cdot x^-$

## Examples

①  $e^- \cdot e^{-}$  is instance of  $x \cdot x^-$ :  $(x \cdot x^-)\sigma = e^- \cdot e^{-}$  for  $\sigma = \{x \mapsto e^-\}$

## Examples

- 1  $e^- \cdot e^{-}$  is instance of  $x \cdot x^-$ :  $(x \cdot x^-)\sigma = e^- \cdot e^{-}$  for  $\sigma = \{x \mapsto e^-\}$
- 2  $e^- \cdot (e^- \cdot e^-)$  is no instance of  $x^- \cdot (x \cdot y)$

## Examples

- 1  $e^- \cdot e^{-}$  is instance of  $x \cdot x^-$ :  $(x \cdot x^-)\sigma = e^- \cdot e^{-}$  for  $\sigma = \{x \mapsto e^-\}$
- 2  $e^- \cdot (e^- \cdot e^-)$  is no instance of  $x^- \cdot (x \cdot y)$

## Remarks

- ▶ term  $t$  can be rewritten if subterm of  $t$  is instance of left-hand side of rewrite rule

## Examples

- 1  $e^- \cdot e^{-}$  is instance of  $x \cdot x^-$ :  $(x \cdot x^-)\sigma = e^- \cdot e^{-}$  for  $\sigma = \{x \mapsto e^-\}$
- 2  $e^- \cdot (e^- \cdot e^-)$  is no instance of  $x^- \cdot (x \cdot y)$

## Remarks

- ▶ term  $t$  can be rewritten if subterm of  $t$  is instance of left-hand side of rewrite rule
- ▶ matching problem is decidable (in linear time)

## Definition (Matching Problem)

instance: terms  $s, t$

question:  $\exists$  substitution  $\sigma: s\sigma = t$ ?

## Matching Algorithm

① start with  $\{s \mapsto t\}$

## Definition (Matching Problem)

instance: terms  $s, t$

question:  $\exists$  substitution  $\sigma: s\sigma = t?$

## Matching Algorithm

- ① start with  $\{s \mapsto t\}$
- ② repeatedly apply following transformation rules

$$\{f(s_1, \dots, s_n) \mapsto f(t_1, \dots, t_n)\} \uplus S \implies \{s_1 \mapsto t_1, \dots, s_n \mapsto t_n\} \cup S$$

## Definition (Matching Problem)

instance: terms  $s, t$

question:  $\exists$  substitution  $\sigma: s\sigma = t?$

## Matching Algorithm

① start with  $\{s \mapsto t\}$

② repeatedly apply following transformation rules

$$\{f(s_1, \dots, s_n) \mapsto f(t_1, \dots, t_n)\} \uplus S \implies \{s_1 \mapsto t_1, \dots, s_n \mapsto t_n\} \cup S$$

$$\{f(s_1, \dots, s_n) \mapsto g(t_1, \dots, t_m)\} \uplus S \implies \perp \quad \text{if } f \neq g$$

## Definition (Matching Problem)

instance: terms  $s, t$

question:  $\exists$  substitution  $\sigma: s\sigma = t?$

## Matching Algorithm

① start with  $\{s \mapsto t\}$

② repeatedly apply following transformation rules

$$\{f(s_1, \dots, s_n) \mapsto f(t_1, \dots, t_n)\} \uplus S \implies \{s_1 \mapsto t_1, \dots, s_n \mapsto t_n\} \cup S$$

$$\{f(s_1, \dots, s_n) \mapsto g(t_1, \dots, t_m)\} \uplus S \implies \perp \quad \text{if } f \neq g$$

$$\{f(s_1, \dots, s_n) \mapsto x\} \uplus S \implies \perp$$

## Definition (Matching Problem)

instance: terms  $s, t$

question:  $\exists$  substitution  $\sigma: s\sigma = t$ ?

## Matching Algorithm

① start with  $\{s \mapsto t\}$

② repeatedly apply following transformation rules

$$\{f(s_1, \dots, s_n) \mapsto f(t_1, \dots, t_n)\} \uplus S \implies \{s_1 \mapsto t_1, \dots, s_n \mapsto t_n\} \cup S$$

$$\{f(s_1, \dots, s_n) \mapsto g(t_1, \dots, t_m)\} \uplus S \implies \perp \quad \text{if } f \neq g$$

$$\{f(s_1, \dots, s_n) \mapsto x\} \uplus S \implies \perp$$

$$\{x \mapsto t\} \uplus S \implies \perp \quad \text{if } S \text{ contains } x \mapsto u \text{ with } t \neq u$$

## Examples

①  $s(y) + s((x + s(0)) + z)$  is instance of  $x + s(y + z)$

## Examples

①  $s(y) + s((x + s(0)) + z)$  is instance of  $x + s(y + z)$ :

$$\{x + s(y + z) \mapsto s(y) + s((x + s(0)) + z)\}$$

①  $s(y) + s((x + s(0)) + z)$  is instance of  $x + s(y + z)$ :

$$\{x + s(y + z) \mapsto s(y) + s((x + s(0)) + z)\}$$

$$\implies \{x \mapsto s(y), s(y + z) \mapsto s((x + s(0)) + z)\}$$

①  $s(y) + s((x + s(0)) + z)$  is instance of  $x + s(y + z)$ :

$$\{x + s(y + z) \mapsto s(y) + s((x + s(0)) + z)\}$$

$$\implies \{x \mapsto s(y), s(y + z) \mapsto s((x + s(0)) + z)\}$$

$$\implies \{x \mapsto s(y), y + z \mapsto (x + s(0)) + z\}$$

①  $s(y) + s((x + s(0)) + z)$  is instance of  $x + s(y + z)$ :

$$\{x + s(y + z) \mapsto s(y) + s((x + s(0)) + z)\}$$

$$\implies \{x \mapsto s(y), s(y + z) \mapsto s((x + s(0)) + z)\}$$

$$\implies \{x \mapsto s(y), y + z \mapsto (x + s(0)) + z\}$$

$$\implies \{x \mapsto s(y), y \mapsto x + s(0), z \mapsto z\}$$

## Examples

①  $s(y) + s((x + s(0)) + z)$  is instance of  $x + s(y + z)$ :

$$\{x + s(y + z) \mapsto s(y) + s((x + s(0)) + z)\}$$

$$\implies \{x \mapsto s(y), s(y + z) \mapsto s((x + s(0)) + z)\}$$

$$\implies \{x \mapsto s(y), y + z \mapsto (x + s(0)) + z\}$$

$$\implies \{x \mapsto s(y), y \mapsto x + s(0), z \mapsto z\}$$

②  $(e \cdot x)^- \cdot ((e \cdot e) \cdot x)$  is no instance of  $x^- \cdot (x \cdot y)$ :

$$\{x^- \cdot (x \cdot y) \mapsto (e \cdot x)^- \cdot ((e \cdot e) \cdot x)\}$$

## Examples

①  $s(y) + s((x + s(0)) + z)$  is instance of  $x + s(y + z)$ :

$$\{x + s(y + z) \mapsto s(y) + s((x + s(0)) + z)\}$$

$$\implies \{x \mapsto s(y), s(y + z) \mapsto s((x + s(0)) + z)\}$$

$$\implies \{x \mapsto s(y), y + z \mapsto (x + s(0)) + z\}$$

$$\implies \{x \mapsto s(y), y \mapsto x + s(0), z \mapsto z\}$$

②  $(e \cdot x)^- \cdot ((e \cdot e) \cdot x)$  is no instance of  $x^- \cdot (x \cdot y)$ :

$$\{x^- \cdot (x \cdot y) \mapsto (e \cdot x)^- \cdot ((e \cdot e) \cdot x)\}$$

$$\implies \{x^- \mapsto (e \cdot x)^-, x \cdot y \mapsto (e \cdot e) \cdot x\}$$

## Examples

①  $s(y) + s((x + s(0)) + z)$  is instance of  $x + s(y + z)$ :

$$\begin{aligned} & \{x + s(y + z) \mapsto s(y) + s((x + s(0)) + z)\} \\ & \implies \{x \mapsto s(y), s(y + z) \mapsto s((x + s(0)) + z)\} \\ & \implies \{x \mapsto s(y), y + z \mapsto (x + s(0)) + z\} \\ & \implies \{x \mapsto s(y), y \mapsto x + s(0), z \mapsto z\} \end{aligned}$$

②  $(e \cdot x)^- \cdot ((e \cdot e) \cdot x)$  is no instance of  $x^- \cdot (x \cdot y)$ :

$$\begin{aligned} & \{x^- \cdot (x \cdot y) \mapsto (e \cdot x)^- \cdot ((e \cdot e) \cdot x)\} \\ & \implies \{x^- \mapsto (e \cdot x)^-, x \cdot y \mapsto (e \cdot e) \cdot x\} \\ & \implies \{x \mapsto e \cdot x, x \cdot y \mapsto (e \cdot e) \cdot x\} \end{aligned}$$

## Examples

①  $s(y) + s((x + s(0)) + z)$  is instance of  $x + s(y + z)$ :

$$\begin{aligned} & \{x + s(y + z) \mapsto s(y) + s((x + s(0)) + z)\} \\ & \implies \{x \mapsto s(y), s(y + z) \mapsto s((x + s(0)) + z)\} \\ & \implies \{x \mapsto s(y), y + z \mapsto (x + s(0)) + z\} \\ & \implies \{x \mapsto s(y), y \mapsto x + s(0), z \mapsto z\} \end{aligned}$$

②  $(e \cdot x)^- \cdot ((e \cdot e) \cdot x)$  is no instance of  $x^- \cdot (x \cdot y)$ :

$$\begin{aligned} & \{x^- \cdot (x \cdot y) \mapsto (e \cdot x)^- \cdot ((e \cdot e) \cdot x)\} \\ & \implies \{x^- \mapsto (e \cdot x)^-, x \cdot y \mapsto (e \cdot e) \cdot x\} \\ & \implies \{x \mapsto e \cdot x, x \cdot y \mapsto (e \cdot e) \cdot x\} \\ & \implies \{x \mapsto e \cdot x, x \mapsto e \cdot e, y \mapsto x\} \end{aligned}$$

## Examples

①  $s(y) + s((x + s(0)) + z)$  is instance of  $x + s(y + z)$ :

$$\begin{aligned} & \{x + s(y + z) \mapsto s(y) + s((x + s(0)) + z)\} \\ & \implies \{x \mapsto s(y), s(y + z) \mapsto s((x + s(0)) + z)\} \\ & \implies \{x \mapsto s(y), y + z \mapsto (x + s(0)) + z\} \\ & \implies \{x \mapsto s(y), y \mapsto x + s(0), z \mapsto z\} \end{aligned}$$

②  $(e \cdot x)^- \cdot ((e \cdot e) \cdot x)$  is no instance of  $x^- \cdot (x \cdot y)$ :

$$\begin{aligned} & \{x^- \cdot (x \cdot y) \mapsto (e \cdot x)^- \cdot ((e \cdot e) \cdot x)\} \\ & \implies \{x^- \mapsto (e \cdot x)^-, x \cdot y \mapsto (e \cdot e) \cdot x\} \\ & \implies \{x \mapsto e \cdot x, x \cdot y \mapsto (e \cdot e) \cdot x\} \\ & \implies \{x \mapsto e \cdot x, x \mapsto e \cdot e, y \mapsto x\} \\ & \implies \perp \end{aligned}$$

## Definition

- ▶ **solution** of matching problem  $\{s_1 \mapsto t_1, \dots, s_n \mapsto t_n\}$  is substitution  $\sigma$  such that  $s_i\sigma = t_i$  for all  $1 \leq i \leq n$

## Definition

- ▶ solution of matching problem  $\{s_1 \mapsto t_1, \dots, s_n \mapsto t_n\}$  is substitution  $\sigma$  such that  $s_i\sigma = t_i$  for all  $1 \leq i \leq n$
- ▶  $\perp$  has no solutions

## Definition

- ▶ solution of matching problem  $\{s_1 \mapsto t_1, \dots, s_n \mapsto t_n\}$  is substitution  $\sigma$  such that  $s_i\sigma = t_i$  for all  $1 \leq i \leq n$
- ▶  $\perp$  has no solutions

## Lemmata

- ▶ there are no infinite derivations  $T_1 \implies T_2 \implies \dots$

## Definition

- ▶ solution of matching problem  $\{s_1 \mapsto t_1, \dots, s_n \mapsto t_n\}$  is substitution  $\sigma$  such that  $s_i\sigma = t_i$  for all  $1 \leq i \leq n$
- ▶  $\perp$  has no solutions

## Lemmata

- ▶ there are no infinite derivations  $T_1 \implies T_2 \implies \dots$
- ▶ if  $S \implies^* T$  then  $S$  and  $T$  have same solutions

## Definition

- ▶ solution of matching problem  $\{s_1 \mapsto t_1, \dots, s_n \mapsto t_n\}$  is substitution  $\sigma$  such that  $s_i\sigma = t_i$  for all  $1 \leq i \leq n$
- ▶  $\perp$  has no solutions

## Lemmata

- ▶ there are no infinite derivations  $T_1 \implies T_2 \implies \dots$
- ▶ if  $S \implies^* T$  then  $S$  and  $T$  have same solutions
- ▶ if  $S \implies^* T$  is maximal derivation and  $T \neq \perp$  then  $T$  is solution of  $S$

## Definition

- ▶ solution of matching problem  $\{s_1 \mapsto t_1, \dots, s_n \mapsto t_n\}$  is substitution  $\sigma$  such that  $s_i\sigma = t_i$  for all  $1 \leq i \leq n$
- ▶  $\perp$  has no solutions

## Lemmata

- ▶ there are no infinite derivations  $T_1 \implies T_2 \implies \dots$
- ▶ if  $S \implies^* T$  then  $S$  and  $T$  have same solutions
- ▶ if  $S \implies^* T$  is maximal derivation and  $T \neq \perp$  then  $T$  is solution of  $S$

## Definition

term is **linear** if it does not contain multiple occurrences of same variable

# Outline

1. Organisation
2. Examples
3. Terms
- 4. Exercises**
5. Further Reading

## Homework Exercises for March 9

- |                  |   |
|------------------|---|
| ① Exercise 2.    | 1 |
| ② Exercise 3(b). | 2 |
| ③ Exercise 2.5.  | 2 |
| ④ Exercise 2.15. | 1 |
| ⑤ Exercise 2.16. | 1 |

## Homework Exercises for March 9

- ① Exercise 2. ①
- ② Exercise 3(b). ②
- ③ Exercise 2.5. ②
- ④ Exercise 2.15. ①
- ⑤ Exercise 2.16. ①
- ⑥ Exercise 4. ☆☆☆

## Starred Exercises

... are optional; solutions give bonus points

# Outline

1. Organisation
2. Examples
3. Terms
4. Exercises
- 5. Further Reading**

- ▶ Section 2.1 (except Definition 2.1.14 — Example 2.1.17 and Definition 2.1.19)

## Lecture Notes

- ▶ Section 2.1 (except Definition 2.1.14 — Example 2.1.17 and Definition 2.1.19)

## Important Concepts

- ▶ closure under contexts
- ▶ closure under substitutions
- ▶ context
- ▶ ground term
- ▶ instance
- ▶ matching algorithm
- ▶ substitution
- ▶ term